

L 05290-57

ACC NR: AR6021347

0

The confused antecedent  $S_{\alpha}$  is a non- $S_1$  noun to which the personal pronoun replacing  $S_2$  may be erroneously referred. The noun  $S_2$  may be replaced with a 3rd person pronoun in the following cases: 1) If between  $S_1$  and  $S_2$  there do not exist any one of the following relations: a)  $S_1$  and  $S_2$  are in the same clause and are syntactically connected; b)  $S_1$  is to the left of the first word of the first principal clause which entirely precedes the clause with  $S_2$ ; c) one of the initial nouns enters into homonymous locution from which the other noun is excluded. 2) If either  $S_{\alpha}$  is absent or, given the presence of  $S_{\alpha}$ , between each  $S_{\alpha}$  and  $S_2$  there exists at least one of the relations not permissible for  $S_1$  and  $S_2$  (cf. a, b, c), taking  $S_{\alpha}$  as  $S_1$  or, lastly, in the presence of  $S_{\alpha}$  the following relationship exists between  $S_1$ ,  $S_2$  and each "dangerous"  $S_{\alpha}$ :  $S_{\alpha}$  and  $S_2$  are present in different clauses while  $S_1$  and  $S_2$  are both in the same clause and are not separated by  $S_{\alpha}$ . The presented result of the verification of these rules with respect to 400 pairs of  $S_1$  and  $S_2$  show that the rules, while not always valid, produce correct results in 89% of cases. Ways of further refining the rules are pointed out. O. Kulagina. [Translation of abstract]

SUB CODE: 05, 09

Card

2/2

*egh*

22977

18-3100

1087, 1454, 1208

S/180/61/000/003/005/012  
E193/E183

**AUTHORS:** Darvoyd, T.I., Vigdorovich, V.N., and Iordanskaya, N.A.

**TITLE:** Purification of thallium by the crystallization methods

**PERIODICAL:** Izvestiya Akademii nauk SSSR, Otdeleniye tekhnicheskikh nauk, Metallurgiya i toplivo, 1961, No.3, pp. 55-62

**TEXT:** Growing demand for high purity thallium in the semiconductor, atomic energy, and optical industries prompted the present author to undertake a systematic study of refining of this metal by the zone melting and crystal pulling techniques. The possibilities of these techniques were first evaluated on the basis of the analysis of the Tl-rich ends of the constitution diagrams of the relevant binary alloy systems. The results of this analysis are presented in Fig.2. Metals with a relatively high solid solubility in Tl are grouped in the left-hand side of the diagram showing their position in the periodic table of the elements; those whose solid solubility in Tl is extremely low are grouped on the right-hand side. Where possible, the distribution coefficients  $K$  were determined from the appropriate constitution diagrams and these are quoted under the symbol of the given metal; the numbered Card 1/9

22977

S/180/61/000/003/005/012  
E193/E183

Purification of thallium by the crystallization methods

arrows indicate groups of metals which (1) form with Tl systems of relatively simple type, (2) are insoluble in liquid Tl, and (3) are characterized by  $K > 1$ . It was inferred from the results of this analysis that most of the impurities likely to be present in thallium (with the exception of metals that are close neighbours of thallium in the periodic table) should be capable of being removed by the crystallization methods, the object of the experimental work carried out by the present author being to check this prediction. The experiments were conducted on Tl specimens with known impurity content, some of which had been preliminarily refined by the alkaline or electrolytic methods. The crystal pulling experiments were conducted in vacuum ( $10^{-4}$  mm Hg); both the crucible and the crystal were rotated (in opposite directions) at 25 and 50 revs/min respectively, the rate of crystal pulling varying between 0.4 and 2 mm/min. The zone refining tests were carried out in O-free, dry nitrogen on bars 150-180 mm long and weighing 20-30 or 150 g. The width of the molten zone was approximately 15 mm, the rates of zone traverse employed being

Card 2/9

22977

S/180/61/000/003/005/012  
E193/E183

Purification of thallium by the crystallization methods

0.5, 1.0 and 2.0 mm/min. Electromagnetic stirring was used in some experiments and the distribution of impurities in the refined bars was determined after 5, 10 and 20 passes; depending on the type of impurity, chemical, spectrographic and radioactive tracer techniques of analysis were used. In the analysis of the results obtained, the behaviour of Cu, Ag, Zn, Sn, Fe, Ni, Mn, S, and Pb is discussed. Some of the typical results are reproduced graphically. Thus, in Fig.4 the Cu concentration ( $C \times 10^4$  wt.%) in the zone refined bar of T1 is plotted against the distance (in % of the bar length,  $l$ ) from the starting end. The four curves relate to bars, examined after 10 (curves 1 and 3) and 20 (curves 2 and 4) passes and refined at the zone traverse rates of 1.0 (curves 1 and 2) or 0.5 (curves 3 and 4) mm/min, the initial Cu content being shown by the broken line - - - -. Fig.6 shows the distribution of sulphur in a bar obtained by the crystal pulling technique (pulling rate 0.5 mm/min); here, the S concentration ( $C \times 10^3$  wt.%) is plotted against the distance from the starting end, measured as the ratio,  $g$ , of the weight of the analysed to the

Card 3/ 9

X

22977

S/180/61/000/003/005/012  
E193/E183

X

**Purification of thallium by the crystallization methods**

total length of the bar. Curves 1, 2 and 3 relate to bars obtained after the molten metal had been held at the temperature for 6, 7 and 11 hours respectively. Finally, the effect of electromagnetic stirring is illustrated in Fig.8, showing the distribution of Cu in a zone-refined bar. Here, log C is plotted against the distance (% l) from the starting end of the bar, obtained with (curves 1 and 2) or without (curves 3 and 4) the application of stirring, at the zone traverse rates of 0.5 (curves 1 and 3) and 1.0 (curves 2 and 4) mm/min. The initial Cu concentration is shown by the broken line. It was concluded that in many cases the zone refining and/or crystal pulling experiments yielded results better than those predicted from the theoretical considerations. This improvement in the segregation coefficient was attributed to the effect of secondary factors. Thus, for instance, the removal of Cd, Hg, and S was assisted by volatilization, that of Cu and Sn by oxydation. Iron which is insoluble in Tl cannot be separated by the methods studied, and filtration has to be used in this case. This is quite an effective method, as has been shown by the results of

Card 4/ 9

22977

S/180/61/000/003/005/012  
E193/E183

**Purification of thallium by the crystallization methods**

experiments in which the thallium samples, containing  $1.8 \times 10^{-4}$  and  $> 10^{-3}$  % Fe, were filtered through porous graphite, after which the Fe concentration was reduced to less than  $5 \times 10^{-5}$  and  $10^{-4}$  %. The concentration of lead in thallium cannot be reduced by the zone refining techniques, and this metal has to be removed by other (alkaline, electrolytic) methods. The effectiveness of zone refining of thallium is greatly increased by the application of electromagnetic stirring.

A.A. Il'inskaya, I.M. Blokh, N.P. Men'shova, V.G. Goryushina, M.A. Notkina, Ye.Ya. Biryukova, V.A. Nazarenko, B.S. Tsivina, N.K. Davidovich and L.I. Gosteva are mentioned for their contributions.

There are 8 figures and 13 references: 10 Soviet and 3 non-Soviet. The English language references read as follows:

Ref.6: K.D. Alexopoulos. Acta crystallogr., 1955, v.8, part 4, p.235

Ref.8: M. Hansen, Lt Anderko. Constitution of binary alloys.

McGraw-Hill Publishing Company, N.Y. - Toronto - London, 1958.

Card 5/9

22977

Purification of thallium by the .... S/180/61/000/003/005/012  
E193/E183

Ref.9: J.L. Haughton, A. Prince. The constitutional diagrams of  
alloys: a bibliography. The Institute of Metals, London,  
1956.

ASSOCIATION: Giredmet/In-t tsvetnykh metallov im. Kalinina  
(Giredmet/Institute of Non-ferrous Metals imeni  
Kalinin)

SUBMITTED: October 8, 1960

Card 6/9

S/080/62/035/010/004/012  
D204/D307

AUTHORS: Vigdorovich, V.N., Darvoyd, T.I., Iordanskaya, N.A.  
and Mamayev, Yu.O.

TITLE: A study of the distribution of Ag admixtures in the  
crystallization methods of the purification of  
thallium

PERIODICAL: Zhurnal prikladnoy khimii, v. 35, no. 10, 1962,  
2165-2170

TEXT: The above subject was investigated in continuation  
of earlier work concerned with the study of phenomena associated  
with the purification of Tl from various metallic admixtures by  
crystallization methods, to determine the effectiveness of purifica-  
tion in relation to the initial concentration of the impurity and  
to the rate of purification, the amounts of Ag being varied between  
0.25 and  $5 \times 10^{-6}\%$ . The Tl crystals were extracted from the melt,  
contained in a graphite crucible, under a pressure of  $10^{-4}$  mm Hg,  
and were 100 - 200 mm long and 8 - 10 mm in diameter. The rates of

Card 1/2



S/080/62/035/010/004/012  
D204/D307

A study of the distribution ...

extraction,  $f$ , were made 0.5, 1.0, and 2.0 mm/min, the crucible being revolved at 25 rpm and the extracting wire at 50 rpm in the opposite direction. The metallic rods were zone-crystallized, under  $O_2$ -free, dry  $N_2$ , and the distributions of Ag along the rods were determined after 5 passes, chemically (for  $< 10^{-3}\%$  Ag) and by an isotope method (for  $\geq 10^{-3}\%$  Ag). L.A. Radushkevich and I.V. Vlasovaya assisted in these determinations. Effective distribution coefficients,  $k$ , (defined by  $k = C/C_0 (1 - g)^{k-1}$ , where  $C_0$  is the initial concentration of Ag and  $C$  is that at a distance  $g$  from the point at which crystallization front was started) calculated from data obtained by these 2 methods, were in fair agreement. The results are discussed, showing that  $k$  decreased with decreasing  $C_0$ , and was lower for higher values of  $f$ . The effect of  $f$  on  $k$  also became greater with decreasing  $C_0$ . In practice, complete purification of Tl from Ag admixtures, by extracting a crystal from the melt and zone-purification, is only effective when  $C_0$  is low, ( $\leq 10^{-4}\%$  Ag); the efficiency of the process may be increased by lowering the rate of crystallization, e.g. to 0.5 mm/min. There are 4 figures and 1 table. ✓

SUBMITTED: April 24, 1961

Card 2/2

IORDANSKAYA, N.I.

Changes in the extra- and intramural nervous system in cardiospasm.  
Kaz. med. zhur. no. 2:49-51 Mr-Apr '61. (MIRA 14:4)

1. Klinika obshchey khirurgii (zav. - prof. A.A. Polyantsev)  
Stalingradskogo meditsinskogo instituta i khirurgicheskoye  
otdeleniye oblastnoy bol'nitsy (glavnyy vrach - A.I. Gusev).  
(CARDIOSPASM) (NERVOUS SYSTEM)

JORDANSKAYA, N. I.

Functional disorder of the vagus nerves in cardiospasm. Vest. khir.  
no.2:24-28 '62. (MIRA 15:2)

1. Iz kliniki obshchey khirurgii (zav. - prof. A. A. Polyantsev)  
Volgogradskogo meditsinskogo instituta i oblastnoy klinicheskoy  
bol'nitsy. Adres avtora: Volgograd, Angarskaya ul., oblastnaya  
bol'nitsa. klinika obshchey khirurgii.

(CARDIOSPASM) (VAGUS NERVE--DISEASES)

IORDANSKIY, N.N.

Some functional characteristics of a crocodile skull. Nauch.  
dokl. vys. shkoly; biol. nauki no.3:42-46 '63. (MIRA 16:9)

1. Rekomendovana kafedroy zoologii pozvonochnykh Moskovskogo  
gosudarstvennogo universiteta im. M.V.Lomonosova.  
(Crocodiles) (Skull)

IORDANSKAYA, N.N.; SEREBRAYAKOV, I.G.

Morphogenesis of a vital form of brushwood exemplified by the  
warty spindle tree *Evonymus verrucosa* Scop. Bot.zhur. 39 no.5:  
768-773 S-O '54. (MIRA 7:11)

1. Zvenigorodskaya biostantsiya Moskovskogo gosudarstvennogo  
universiteta.  
(Spindle tree)

**IORDANSKAYA, N.N.**

**Phreatophytes of the Chernyye Zemli and some aspects of their  
ecology. Biul.MOIP.Otd.geol.31 no.3:117-118 My-Je '56.  
(Chernyye Zemli—Botany) (MLBA 9:12)**

IORDANSKAYA, N.N.

~~SOME DATA ON phreatophyte root systems in the Chernyye Zemli [with  
summary in English].~~ ~~Biul. MOIP. otd. biol. 63 no.1:79-87 Ja-F '58.~~  
(MIRA 11:5)

(CHERNYYE ZEMLI--WATER, UNDERGROUND)

USSR/Pharmacology. Toxicology. Chemotherapeutic Preparations.

C

A) Antibiotics

Abs Jour : Ref Zhur - Biol., No II, 1958, No 52086

Author : ~~Jordanskaya, N. Ye.~~, Matynkina A.A., Khachaturova T.I.  
Inst : Uzbek Tuberculosis Institute  
Title : The Immediate Therapeutic Effect of the Preparation Larusan

Orig Pub : Sb. tr. Uzb. n-i. tuberk. in-t, 1957, 3, 70-75

Abstract : Larusan (I) was administered to 54 patients (adults) with various forms of pulmonary tuberculosis (the majority with fibro-cavernous processes) in doses of 0.2 g, 3 times daily. It was demonstrated that I lowered toxemia, and in many cases led to improvement of the local process. Toxic side-effects (giddiness, headaches, excitement, precordial pains) were observed in 4 patients treated with I. As compared with phthivazid, I was less effective. -- V.I. Yel'nik.

Card : 1/1



IODANSKAYA, N.Ye.; MATYNKINA, O.A.; KHACHATUROVA, T.I.

Immediate therapeutic affect from the drug, Iarusan. Sbor. trud.  
Uz. nauch.-issl. tub. inst. 3:82-86 '57. (MIRA 14:5)  
(ISONICOTINIC ACID) (TUBERCULOSIS)

IODANSKAYA, N.Ye.

Immediate results of antibacterial therapy combined with novocaine  
block in pulmonary tuberculosis patients. Sbor. trud. Uz. nauch.-  
issl. tub. inst. 3:105-110 '57. (MIRA 14:5)  
(TUBERCULOSIS) (NOVOCAINE)

**IX** IORDANSKAYA, N.Ye., Cand Med Sci -- (diss) "Novocain  
block~~ed~~ in symptomatic therapy of pulmonary tuberculosis."  
Tashkent, 1959, 16 pp (Min of Health UzSSR. Tashkent State Med  
Inst) 250 copies (KL, 33-59, 121)

NESMEYANOVA, T.N.; IORDANSKAYA, Ye.I.; BRAZOVSKAYA, F.A.

Effect of various doses of pyrogenal on the formation of a  
brain scar. Biul. eksp. biol. i med. 56 no.9:115-119 S '63.

(MIRA 17:10)

1. Iz Instituta vysshey nervnoy deyatel'nosti i neyrofiziologii  
AN SSSR. Predstavlena deystvitel'nym chlenom AMN SSSR A.V. Le-  
bedinskim.

USSR/Medicine - Physiology

FD-2697

Card 1/1

Pub. 33-6/28

Author : Shatenshteyn, D. I.; Iordanskaya, Ye. N.

Title : Towards the physiology of the motor analysor of man

Periodical : Fiziol. zhur. 41, 35-42, Jan-Feb 1955

Abstract : Investigated the functional state of the central terminal of the motor analysor in man and the development of states of excitation and inhibition in it during work. Ergograms. Nine references, all USSR (6 since 1940)

Institution : Laboratory of Physiology of Labor of the Institute of Hygiene of Labor and Occupational Diseases of the Academy of Medical Sciences USSR

Submitted : December 24, 1953

IORDANSKAYA, Ye.N.

Size of a conditioned motor reflex in man as a function of intensity of conditioned auditory stimuli [with summary in English]. Zhur. vys.nerv.deiat. 8 no.1:28-35 Ja-F '58. (MIRA 11:3)

1. Institut biofiziki AN SSSR, Moskva.

(REFLEX CONDITIONED,

eff. of intensity of sound stimuli on level of motor reflex (Rus)

BRAZOVSKAYA, F.A.; NESMEYANOVA, T.N.; IORDANSKAYA, Ye.N.

Scar formation in the central nervous system under the influence  
of pyrogenal. *Biul. eksp. biol. i med.* 50 no. 11:121-123 N '60.  
(MIRA 13:12)

1. Iz fiziologicheskoy laboratorii Akademii nauk SSSR, Moskva.  
(PYROGENS) (SPINAL CORD) (CICATRICES)

NESMEYANOVA, T.N.; BRAZOVSKAYA, F.A.; IODANSKAYA, Ye.N.

Case of partial regeneration of nerve conductors in sectioned spinal cord in dogs. Fiziol.zhur. 46 no.2:202-209 F '60. (MIRA 14:5)

1. From the Physiological Laboratory, U.S.S.R. Academy of Sciences, Moscow.

(NERVOUS SYSTEM--DEGENERATION AND REGENERATION)



BRAZOVSKAYA, F. A.; NESMEYANOVA, T. N.; IORDANSKAYA, Ye. N. (Moskva)

Effect of pyrogenal on the formation of the cicatrix after  
sectioning of the spinal cord. Vop. neirokhirurgii no.3:6-9  
'62. (MIRA 15:7)

1. Fiziologicheskaya laboratoriya Akademii nauk SSSR.

(SPINAL CORD SURGERY) (CICATRICES)  
(PYROGENAL)

IORDANSKIY, A., red. toma; ATROSHCHENKO, I., tekhn. red.

[Science and mankind, 1963] Nauka i chelovechestvo,  
1963. Moskva, Izd-vo "Znanie," 1963. 522 p.  
(MIRA 17:1)

IORDANSKIY, A.B.

Radioautographic study of chromosome reproduction in Vicia  
faba. Dokl. AN SSSR 158 no.1:192-195 S-0 '64 (MIRA 17:8)

1. Institut radiatsionnoy i fiziko-khimicheskoy biologii AN  
SSSR. Predstavleno akademikom A.N. Belozerskim.

ICORDANSKIY, A.B.

Interchromatid exchanges in the chromosomes of beans. *Cytologia*  
(MIRA 18:8)  
6 no.6:732-741 N-5 '64.

1. Laboratoriya obshchey i kosmicheskoy karyologii Instituta  
radiatsionnoy i fiziko-khimicheskoy biologii AN SSSR, Moskva.

BOGDANOV, Yu.F.; IORDANSKIY, A.B.; GINDILIS, V.M.

Problem of multistrand chromosome model. Genetika no.5:82-100  
N '65. (MIRA 19:1)

1. Institut molekulyarnoy biologii AN SSSR, Moskva. Submitted  
August 25, 1965.

5  
IORDANSKIY, A.B.

Autoradiographic analysis of sister interchromatid exchanges  
in the third division following exposure to H<sup>3</sup>-thymidine.  
TSitologiya 7 no.5:673-675 S-0 '65. (MIRA 18:12)

1. Laboratoriya obshchey i kosmicheskoy kariologii Instituta  
radiatsionnoy i fiziko-khimicheskoy biologii AN SSSR, Moskva.  
Submitted March 19, 1965.

LUKASHEV, Konstantin Ignat'yevich; IORDANSKIY, A.D., red.

[Atoms and our planet] Atomy i nasha planeta. Moskva,  
Znanie, 1965. 91 p. (Narodnyi universitet: Estestvenno-  
nauchnyi fakul'tet, no.6) (MIRA 18:7)

1. Vitse-prezident AN Belorusskoy SSR (for Lukashev).

IORDANSKIY, A.D.; red. vypuska; ATROSHCHENKO, L., tekhn. red.

[Science and mankind] Nauka i chelovechestvo, 1962. Moskva,  
Izd-vo "Znanie," 1962. 404 p. (MIRA 16:6)  
(Science--Yearbooks)



PROCESSES AND PROPERTIES INDEX

12

6-Ethoxyquinoline. A. N. Iordanskii. Russ. AZ. 327 Dec. 31, 1927. To produce film is added 20% HCl and the mixt. is heated with Act1.

AS 35.5 A METALLURGICAL LITERATURE CLASSIFICATION

CLASSIFICATION	INDEX
1	2
3	4
5	6
7	8
9	10
11	12
13	14
15	16
17	18
19	20
21	22
23	24
25	26
27	28
29	30
31	32
33	34
35	36
37	38
39	40
41	42
43	44
45	46
47	48
49	50
51	52
53	54
55	56
57	58
59	60
61	62
63	64
65	66
67	68
69	70
71	72
73	74
75	76
77	78
79	80
81	82
83	84
85	86
87	88
89	90
91	92
93	94
95	96
97	98
99	100

1ST AND 2ND SERIES      3RD AND 4TH SERIES

PROCESSES AND PROPERTIES INDEX

25

*Ch*

**Doth. A. N. Jordanik. Russ. 56,321, Dec. 31, 1939.**  
Tertiary amine of heterocyclic bases are fused with aromatic aldehydes in the presence of the usual condensing agents in vacuo at a temp. of 110-115°, with removal of the water formed during the reaction.

COMMON ELEMENTS

COMMON VARIABLES INDEX

ABB-51A METALLURGICAL LITERATURE CLASSIFICATION

6-277,577,74872

1ST AND 2ND SERIES      3RD AND 4TH SERIES

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----

IORDANSKIY, A. et al.

"Colorphotographu on Triple-Emulsion Light-sensitive Maters". Goskinoizdat,  
State Publishing House of Cinematographic Literature, M., 1949.

IORDANSKIY, A. <sup>N</sup> et al

Color Photography on Three-Layer Photosensitive Materials. Goskinoizdat (1949)

IODANSKIY, A. N. and CHELTZOV, V. S.

"Color in the Cinema", (tztvet v kino), published by the State Publishing House for Cinematography, Moscow, 1950.

SO: D-52286, 9 July 1954.

2

CA

In memory of G. I. Aronov, K. V. Astakhov, A. N. Jordanishvili, I. I. Levtchenko, and N. N. Sveshnikov. *Uspehi* *Khim.* 38, 645-6 (1969).—Obituary, with list of publications and portrait. N. Thon

1957

IORDANSKIY, A. N.

177T32

USSR/Chemistry - Photography

Mar 51

"Brief Communication: The Yield of Dyestuffs in Color Development," A. N. Iordanskiy, G. I. Arbuzov, All-Union Sci Res Cine Photo Inst

"Zhur Prik Khim" Vol XXIV, No 3, pp 337-340

Showed by expt that in color development 1 mol of dyestuff that composes purple partial image is formed for each 4 atoms of reduced (metallic) AG, supporting theoretical ideas on dyestuff formation in process based in part on work by A. Ye. Paray-Koshits and G. I. Arbuzov. Showed for emulsions applied in practice true yield of dyestuff is independent of av grain size.

177T32

CA

5

The yield of dyes in color development. A. N. Jordan'skii

and G. I. Arbusov. *J. Applied Chem. U.S.S.R.* 24, 373-7 (1951) (Engl. translation); (Russ. ed., 337-40) cf. *C.A.* 45, 10105M. -- The mol. ratio of azomethine dye (I) formed to Ag produced, on development of 3-layer color film, is predicted as 1:4 (Tull, *C.A.* 33, 0163D), and is described as a relative ratio by the Cheltsov ratio (II),  $\Delta D_{\lambda} / \Delta D_{Ag}$ , where  $D_{\lambda}$  and  $D_{Ag}$  are corresponding optical ds. of I (at  $\lambda_{max}$ ) and of Ag image. The  $D_{\lambda}$  ( $\lambda_{max}$ )--concn. relation was detd. for a purple I (III), formed in substance and examd. in 5% gelatin films at  $\lambda = 620 m\mu$ , and was found to obey Beer's law at least for  $D_{620}$  of 0.00-3.00. These results gave the amt. of dye,  $q'$ , in mol. cm.<sup>-1</sup> for  $D_{620} = 1$ . The absorption curve of III formed by development was found to be the same. Several un sensitized, standard emulsions, varying in mean grain size (IV) (3 NH<sub>2</sub> types, including sound and spectral, and one non-NH<sub>2</sub>), were used. After const. exposure, a variable development time (0.5-3.5 min.) with standard developer gave a film which was cut in half. III in one half was bleached with dil. H<sub>2</sub>SO<sub>4</sub> to give  $D_{Ag}$ ; Ag in the other half was removed with K<sub>2</sub>Fe(CN)<sub>6</sub> and Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> to give  $D_{Ag}$ . For  $D_{620} = f(D_{Ag})$ , which resembled the log illumination vs.  $D_{Ag}$  curve, from the linear portion with maximal II,  $\Delta D_{620}$  was found for  $\Delta D_{Ag} = 0.4$ , giving II = 6.5-4.3 for a IV radius of 0.40-0.31. The abs. mol. ratio  $\phi = \Delta Q / \Delta C$ , where  $\Delta Q = q' \cdot \Delta D_{620}$ , and  $\Delta C = \rho' \cdot \Delta D_{Ag}$  ( $\rho'$  is the photometric equiv. as g. atoms Ag cm.<sup>-1</sup> for each emulsion type), was 0.240-0.260, unaffected by IV, and agreeing with theory. W. B. McCormack



IORDANSKIY, A.N.

USSR/Chemistry - Photography

1 May 52

"Dyestuff Yield in Color Development," S. A. Bondard, A. N. Iordanskiy, V. S. Chel'tsov

"Dok Ak Nauk SSSR" Vol LXXXIV, No 1, pp 81-84

The relationship between the amts of silver and dyestuff formed during color development with dyestuff components of various classes was studied. As typical components, the following were chosen: for yellow derive of anilide of an acylacetic acid; purple, a compd of the pyrazalone series; blue a deriv of 1,2-hydroxymaphthalene carbonic acid contg a sulfonic acid group in the 4-position. The

22476

relationship between the optical density of the dyestuff and its surface concn in the photographic layer was detd and found to be a linear function. In order to det the yield of dyestuff, which was found to be const throughout the development process, the relationship between the optical density of the depth of color and the surface concn of metallic silver formed during the development process was experimentally established. Presented by Acad A. N. Terenin 1 Mar 52.

22476

JORDANSKIY, A-N

U S S R .

Field of dye in color development. V. S. Chel'tsov, A. S. Jordanskiy, M. V. Krasheninnikova, and S. A. Bogard. *Uspekhi Nauch. Fot., Akad. Nauk S.S.S.R., Otdel. Khim. Nauk* 2, 48-55(1954).—The relative photographic yield of the dye was d<sub>rel</sub>, rather than the mol. yield. The yield was expressed as  $D_{\lambda}/D_{\lambda_0}$ , the ratio of optical d. of the dye (found for monochromatic light with wave length corresponding to max. of absorption) to optical d. of correspond-

ing Ag image. The influence of concns. of developing agents, diffusing components, and Na<sub>2</sub>SO<sub>3</sub> was studied. The effect of different components and influence of developing were studied. With increase of developing time, the coeff. of contrast of the dye image increased faster than that for the Ag image. Relative photographic yield depended on the properties of the emulsion and developing conditions of the Ag image.  
Euzilla Mayer

AB 25/4

IORDANSKIY, A. N., et al. and CHELZOV, V. S.

"On the Inter-Relation of the Optical Density of Silver and Dyestuff in Color Development," a paper given at the International Conference on Scientific Photography, Cologne, 24-27 Sep 1956

E-3072367

ICORDANSKIY, A.M.

Influence of the structure of multilayer color photographic materials on their resolving power and on the sharpness of the image. Zhur.nauch. i prikl.fot.i kin. 1 no.1:52-55 Ja-F '56. (MLRA 9:10)

I.Vsesoyuznyy nauchno-issledovatel'skiy kino-foteinstitut.  
(Color cinematography)

REF ID: A66111

✓ 909

20

771.534.554 : 771.356

**Spectrozoal Photography.** A. N. JORDANSKII. *Zh. nauch. priklad. Fotogr. Kinematogr.*, Jan.-Feb. 1957, 2, 28-34. [In Russian].—A method is described for calculating the difference in optical density of photographic images of two objects ( $\Delta D\lambda^{a/b}$ ), [where  $\Delta D\lambda^{a/b}$  is the difference between the two densities  $D^a$  and  $D^b$ ]. Results are given for the calculation of the quantity  $\Delta D\lambda^{a/b}$  for a number of objects typical of a summer landscape in relation to "forest plantations of deciduous species in the summer period". It is shown to be impossible to distinguish clearly all the objects studied in the summer landscape by photographing them with the aid of any monochromatic radiation within the range 400-840m $\mu$ . It is shown that one group of objects shows the best image in relation to the foliage on photographing with the aid of red radiation, while the other group requires the aid of infra-red radiation. It is shown that, for separate objects referred to the group in which the best image in relation to the foliage appears on photographing with the aid of infra-red radiation, an increase in the wavelength corresponding to more than 790 and 820 m $\mu$  leads to deterioration in their appearance. The principle of spectrozoal photography and the formulation of its basic requirements are discussed. S.C.G.

Translation of Author's Abstract.

*Handwritten initials: RD and a signature*

*Handwritten notes: 4, 4E4C, 4E2D*

*IORDANSKIY, A.N.*

USSR/Chemical Technology - Chemical Products and Their  
Application. Photographic Materials.

I-6

Abs Jour : Ref Zhur - Khimiya, No 1, 1958, 2425

Author : Rozental', L.V., Iordanskiy, A.N.

Inst : -

Title : Black Antihalo Counterlayer of Color Motion Picture Films.

Orig Pub : Tekhnika kino i televideniya, 1957, No 7, 63-72

Abstract : Description of the properties of a black antihalo counter-  
layer consisting of a dispersion of finely dispersed car-  
bon black having high colloidal stability, in cellulose  
acetophthalate. It is reported that deposition of such a  
counterlayer on the backing results in a sharp increase  
of the resolving power of the film without lowering its  
photographic, physical and mechanical characteristics.

Card 1/1

AUTHOR: Iordanskiy, A.N. SOV 77-3-4-7/23

TITLE: Spectrozoal Photography (Spektrozoal'naya fotografiya);  
II. Chromatic and Achromatic Density Detail as a Guide to the  
Mutual Exposure of Spectrozoal Images (Khromaticheskaya i akhro-  
maticheskaya detali potemneniya - mera vzaimnogo vyyavleniya  
spe trozoal'nykh izobrazheniy)

PERIODICAL: Zhurnal nauchnoy i prikladnoy fotografii i kinematografii, 1958,  
Vol 3, Nr 4, pp 275-278 (USSR)

ABSTRACT: The author discusses the way a change in the spectral brightness  
factors of a pair of objects is reflected in a change in the op-  
tical density difference of their photographic images for both  
achromatic and chromatic film. He proposes the use of chromatic  
and achromatic density detail as a gage to the difference in the  
two-color spectral images of the objects and works out a mathema-  
tical method for calculating the values of the density detail,  
making use of the spectral brightness factors of the objects being  
photographed for two monochromatic radiations and two spectral  
zones. There are 2 figures and 2 Soviet references.

Card 1/2

SOV 77-3-4-7/23

Spectrozoal Photography; II Chromatic and Achromatic Density Detail as a Guide  
to the Mutual Exposure of Spectrozoal Images

ASSOCIATION: Vsesoyuznyy nauchno-issledovatel'skiy kinofotoinstitut (All-  
Union Research Institute for Photography and Cinematography)

SUBMITTED: June 14, 1957

1. Photographic film--Performance
2. Photographic film--Properties
3. Mathematics--Applications

Card 2/2



CHEL'TSOV, V.S., kand.khim.nauk; BONGARD, S.A., kand.khim.nauk;  
IORDANSKIY, A.N., kand.tekhn.nauk

Present-day methods of producing color photographs. Khim.nauk 1  
prom. 3 no.5:576-587 '58. (MIRA 11:11)  
(Color photography--Three-color process)

ORDANSKIY A. N.  
p. 3.

PHASE I BOOK EXPLOITATION

SOV/3815  
SOV/7-M-7

Akademiya nauk SSSR. Laboratoriya aerometodov

Trudy, tom 7: Materialy VII Vsesoyuznogo mezhdruvedomstvennogo soveshchaniya po aeros"yemke, 25 noyabrya - 1 dekabrya 1956 g. (Transactions of the Laboratory of Aerial Methods, Academy of Sciences USSR, Vol. 7: Materials of the 7th All-Union Interdepartmental Conference on Aerial Surveying) Moscow, 1959. 331 p. 1,400 copies printed.

Editorial Board: A.V. Glagolev, V.G. Zdanovich, N.G. Kell' (Resp. Ed.), D.M. Kudritskiy, K.S. Lyalikov, and G.G. Samoylovich; Ed. of Publishing House: D.M. Kudritskiy; Tech. Ed.: M.Ye. Zindel'.

**PURPOSE:** This collection of articles is intended for photogrammetrists. The articles will be of interest to all governmental and industrial agencies concerned with aerial photography.

**COVERAGE:** This is the first volume of a 2-volume work containing reports read at the All-Union Conference on Photogrammetry which took place in Leningrad from November 25 to December 1, 1956, under the auspices of the Laboratory of Aerial Photography Methods of the Academy of Sciences USSR. These reports

Card 1/15

Transactions of the Laboratory (Cont.)

80V/3815

describe the principles and applications of photo interpretation in the fields of soil science, forestry, geology, hydrology, industrial development, etc. Individual reports discuss the equipment used and techniques employed. References accompany each article.

TABLE OF CONTENTS:

Kell', N.G. Preface	3
Bilanov, A.I. [Glavnoye upravleniye geodezii i kartografii - Main Administration of Geodesy and Cartography]. Organization, Planning, and Execution of Aerial Survey Operations by the Main Administration of Geodesy and Cartography of the Ministry of Internal Affairs, USSR	5
Mikhaylov, V.Ya. [Tsentral'nyy nauchno-issledovatel'skiy institut geodezii, aerofotos''yemki i kartografii - Central Scientific-Research Institute of Geodesy, Photogrammetry, and Cartography]. Present State and Future Prospects of the Development of Scientific Programs in Aerial Photography	10

Card 2/15

Transactions of the Laboratory (Cont.)

SOV/3815

Lyalikov, K.S. [Laboratoriya aerometodov - Laboratory of Aerial-Surveying Methods].

19

Ways of Improving Aerial Photography

Iordanskiy, A.N. [Nauchno-issledovatel'skiy kinofotoinstitut - Scientific-Research Institute of Photography and Cinematography].

25

Spectrozoal Photography and Spectrozoal Films [Color Photography]

Veydenbakh, V.A. [Gosudarstvennyy opticheskiy institut imeni S.I. Vavilova - State Institute of Optics imeni S.I. Vavilov].

32

Speed Methods of Processing Aerial Photographic Materials

Feygel'son, Ye.M., and N.S. Malkevich [Institut fiziki atmosfery - Institute of Atmospheric Physics].

37

Computation of Light Intensity and Haze Coefficients in Anisotropic Dispersion

Card 3/15

S/081/61/000/022/057/076  
B101/B147

AUTHORS: Kilinskiy, I. M., Iordanskiy, A. N.

TITLE: Influence of the yellow color filter layer on the resolving power and effective color sensitivity of color film layers

PERIODICAL: Referativnyy zhurnal. Khimiya, no. 22, 1961, 381, abstract 22L338 (Tr. Vses. n.-i. kinofotoin-ta, no. 29, 1959, 59-61)

TEXT: The yellow filter layer containing colloidal Ag hardly reduces the resolving power of the green- and red sensitive layers of the color film, but slightly reduces its effective sensitivity to light. It is advisable to replace the layer with the colloidal Ag by a light filter having a higher transmissivity for green and red light. [Abstracter's note: Complete translation.]



Card 1/1

KILINSKIY, I.M.; IORDANSKIY, A.N.

Effect of silver halide concentration of the emulsion layer on its resolving capacity dependent on the nature of the developing agent. Zhur.nauch.i prikl.fot. i kin. 5 no.2:108-113 Mr-Ap '60.  
(MIRA 14:5)

1. Vsesoyuznyy nauchno-issledovatel'skiy kinofotoinstitut (NIKFI).  
(Photography—Developing and developers)

VILENSKIY, Yu.B.; IORDANSKIY, A.N.; BUDARINA, N.N.

Some problems of the improvement of color reproduction and  
sharpness of positive color films. Usp. nauch. fot. 8:13-20  
'62. (MIRA 17:7)

S/058/63/000/003/046/104  
A062/A101

**AUTHORS:** Kilinskiy, I. M., Vilenskiy, Yu. B., Iordanskiy, A. N.

**TITLE:** On the improvement of light-sensitivity, resolving power and quality of color reproduction in color negative motion-picture films

**PERIODICAL:** Referativnyy zhurnal, Fizika, no. 3, 1963, 87, abstract 3D587 ("Uspekhi nauchn. fotogr.", 1962, v. 8, 3 - 12)

**TEXT:** The article describes new color films, produced by NIKFI and the Shostkin chemical plant. The increase of light sensitivity has been attained owing to a rational choice of the form of change in the quantity of excessive bromide in the ripening process of the emulsion. The results of work on sensitization of color photography materials, filter layer structure etc. are described. It is shown that an increase of sharpness in color images can be attained by a reduction of light scattering in the elementary layers, and an improvement of the color reproduction - by introducing into these layers masking components. Peculiarities of the treatment of films with internal masking are described.

[Abstracter's note: Complete translation]

Card 1/1

D. Balabukha



S/058/63/000/003/047/104  
A062/A101

AUTHORS: Vilenskiy, Yu. B., Iordanskiy, A. N., Budarina, N. N.

TITLE: Some problems in the improvement of color reproduction and sharpness in color positive films

PERIODICAL: Referativnyy zhurnal, Fizika, no. 3, 1963, 87, abstract 3D588 ("Uspekhi nauchn. fotogr.", 1962, v. 8, 13 - 20)

TEXT: Some problems in the improvement of color reproduction and image sharpness are considered, related to the properties of color positive materials. For improving the color separation it is proposed to use AgCl emulsions and more selective dyes, and for increasing the sharpness - to displace the components with respect to the sensitizers in the emulsion layers. A series of motion-picture materials, both from this country and from abroad, which meet these requirements are described.

D. Balabukha

[Abstracter's note: Complete translation]

Card 1/1

IORDANSKIY, A.N.

New spectrosonal negative films. Zhur. nauch. i prikl. fot.  
i kin. 9 no.3:210-211 My-Je '64. (MIRA 18:11)

1. Vsesoyuznyy nauchno-issledovatel'skiy kinofotoinstitut  
(NIKFI). Submitted February 6, 1964.

IODANSKIY, D.

Machine and speech. Un.tekh. 5 no.3 25-28 Mr '61.

(MIRA 14:6)

(Cybernetics) (Speech)

GADOMSKI, Yan; IORDANSKIY, D. [translator]

Colored stars. IUn.tekh. 5 no.4:54 Ap '61.  
(Stars)

(MIRA 14:3)

L 39892-06 MT(d)/EMP(k)/EMP(h)/EMP(l)/EMP(y) 14/GR-2  
ACC NR: AP6017694 SOURCE CODE: UR/0103/65/026/012/2289/2291

AUTHOR: Iordanskiy, D. I. 12B

ORG: none

TITLE: Problems of the control of large systems (Scientific conference held in Poland)

SOURCE: AN SSSR. Avtomatika i telemekhanika, v. 26, no. 12, 1965, 2289-2291

TOPIC TAGS: scientific conference, automatic control system, nervous system

ABSTRACT: The conference was held in November 1964 in Yablonna near Warsaw. 96 Polish specialists plus others from Bulgaria, Hungary and Yugoslavia attended. Reports were heard on the general problems involved in the control of large systems, the structure of large systems, the classification of large systems, adaptive optimization of hierarchically controlled large systems, reliability in large systems, the nervous system as an example of a large system, and biological receptors and associated nerve networks as an example of the inputs to a large system. Particular attention was paid to large-scale production processes. [JPRS]

SUB CODE: 13, 06 / SUBM DATE: none

Card 1/1 N/S

UDC: 62-50:061.3

L 37107-66 EWP(c)/EWP(k)/EWT(d)/EWP(h)/EWP(l)/EWP(v) BC/JT/JXT(BF)/GD  
ACC NR: AT6012883 SOURCE CODE: UR/0000/65/000/000/0020/0027

AUTHOR: Iordanskiy, D. I.

ORG: None

TITLE: Some problems of studying staffs controlling large systems

SOURCE: Sistema chelovek i avtomat (Man-automaton systems). Moscow, Izd-vo Nauka, 1965, 20-27

TOPIC TAGS: automatic control system, bionics, man machine communication, information theory

ABSTRACT: The author defines a large system as one containing people. Large systems must include people or be totally made up of people, and are divided into regular and irregular. An example is given depicting a regular system such as an automobile plant. Examples of irregular systems are mine construction, house building, and hydroelectric station construction. The function of staffs working in large irregular systems are not considered. A staff or collective is defined as the entire aggregate of people controlling a given large system. An example of such a staff or collective is the entire staff of a plant, hospital, or design bureau. The advantages of studying staffs in Communist and

Card 1/3

47  
B+1

L 37107-66

ACC NR: AT6012883

Socialist countries are discussed. The staff is studied as a set. The effect of personnel on the function of large systems is studied. The staff is studied as a certain set apart from all other elements of a large system. It is essential to keep in mind that all the parts constitute a single whole. A set is given and a certain complex system K. Together they constitute a large system with n inputs  $X = \{x_1, x_2, \dots, x_n\}$  which are the controlling

actions of people, and t outputs  $Y = \{y_1, y_2, \dots, y_t\}$  which are the operational indexes of the system K yields the mapping  $X \rightarrow Y$ . The staff A can be divided according to purpose.

Thus the staff of a large system which is the set A can be divided into groups such as  $A_1, A_2, \dots, A_k$ . These groups include people performing identical or near identical functions.

Such a classification is useful in studying the part played by people in large systems as a function of their tasks. The groups  $A_1, A_2, \dots, A_k$  are called nonintersecting subsets.

Overlapping of the boundaries is permissible. Various elements can also cross over.

Thus

$$A = \bigcup_{i=1}^k A_i, A_i \cap A_j = \emptyset \text{ for all } i \neq j$$

Card 2/3

L 37107-66

ACC NR: AT6012883

It is shown that  $A_i = \{a_{i1}, a_{i2}, \dots, a_{ip}\}$ , where  $a_{ij} \in A$  are the elements of the initial set—staff

An example is given based on several parallel assembly lines where identical units are assembled. Each line includes workers performing identical operations. A graphic representation of this is given. An expression is given for the correlation coefficient. A dispersion of the respective magnitudes is presented. Two criteria should be considered in studying large systems: the extent to which enumerated and unenumerated characteristics of staffs are reflected in the operational indexes indices of large systems and the extent to which work conditions and characteristics of the systems themselves affect the material and psychological needs of the staff members. Orig. art. has: 6 figures and 10 formulas.

SUB CODE: 05/  
09/ SUBM DATE: 02Aug65 / ORIG REF: 002  
06/

*me*  
Card 3/3



SOV-98-58-2-6/21

AUTHORS: Karpov, A.N., and Iordanskiy, I.Ye., Engineers

TITLE: The Reconstruction of the Shores of the Tsimlyanskoye Water Reservoir (Pererabotka beregov Tsimlyanskogo vodokhranilishcha)

PERIODICAL: Gidrotekhnicheskoye stroitel'stvo, 1958, Nr 2, p 27 (USSR) *FEB*

ABSTRACT: To obtain factual material on the rebuilding of the shores of large water reservoirs, the profiles of shores consisting of various rock formations were studied at the Tsimlyanskoye Water Reservoir in 1953. The measuring of the shores at the selected sections was carried out for 3 years. Figures 1 and 2 show the results of the observations, which have led to the preliminary conclusion that it is possible to forecast the amount of erosion of sandy shores. Little is known about the form of shores consisting of rocks which convert

Card 1/2

SOV-98-58-2-6/21

The Reconstruction of the Shores of the Tsimlyanskoye Water Reservoir

easily into a suspension state.  
There are 2 diagrams.

1. Inland waterways--USSR
2. Beaches--Erosion

Card 2/2

JORDANSKIY, S.V.

40-4-3/24

AUTHOR: JORDANSKIY, S.V. (Moscow)

TITLE: On the Stability of a Plane Stationary Shock Wave (Ob ustoychivosti ploskoy statsionarnoy udarnoy volny).

PERIODICAL: Prikladnaya Mat.i Mekh., 1957, Vol.21, Nr 4, pp.465-472 (USSR)

ABSTRACT: A plane piston is assumed to move with constant velocity in the direction of the negative x-axis in a homogeneous medium. The author considers the perturbations of the hydrodynamic parameters behind the front of the arising shock wave. In linear approximation there hold the equations

$$(1) \frac{\partial \vec{u}}{\partial t} = -v \text{ grad } \pi, \quad \frac{\partial \pi}{\partial t} + \frac{c^2}{v} \text{ div } \vec{u} = 0, \quad \frac{\partial \epsilon}{\partial t} = 0$$

in a coordinate system moving with the shock wave, where  $v$  is the resting specific volume and  $\pi, \vec{u}, \epsilon$  are small perturbations of pressure, velocity and entropy behind the shock-wave front;  $c$  is the velocity of sound. Let the equation of the front surface be

$$(2) \quad x = -vt + \xi(y, z, t)$$

where  $\xi$  denotes the perturbation of the surface. From (1) it follows

CARD 1/3

On the Stability of a Plane Stationary Shock Wave

40-4-3/24

$$\frac{\partial^2 \eta}{\partial t^2} - c^2 \frac{\partial^2 \eta}{\partial x^2} + k^2 c^2 \eta = 0$$

The boundary conditions are given by D'yakov's linear approximations (Zh.E.T.F., 27, 3, 1954). For  $\zeta$  it holds:

$$(3) \quad \frac{d\zeta}{ds} = \frac{d\zeta}{ds} \Big|_{\Gamma} - \mu \int_0^{\infty} U(s-w) J_0(s-w) \left( \frac{d^2 \zeta}{dw^2} + \nu \zeta \right) dw + F +$$

$$+ \int_0^{\infty} U(s-a-b) \left\{ \alpha f_1 J_0'(f) \frac{d\zeta}{dw} - \mu J_0(f) \left( \frac{d^2 \zeta}{dw^2} + \nu \zeta \right) \right\} dw$$

Here it is  $\alpha = \frac{c-v}{c+v}$ ,  $s = \sqrt{\alpha} k \xi_0$ ,  $\mu = \frac{c}{2v}(1-j)$ ,  $\zeta = ct - x$ ,

$\nu = \frac{v v_0}{c^2 - v^2} \frac{1+j}{1-j}$ ,  $L = \frac{2lk}{\sqrt{\alpha}}$  (l describes the position of the

piston for  $t=0$ ),  $\Gamma = \alpha s - L$ ,  $f = \sqrt{(s-a)^2 - b^2}$ ,  $f_1 = -\frac{1}{2} \sqrt{\frac{s-a-b}{s-a+b}} +$

$+\frac{1}{2\alpha^2} \sqrt{\frac{s-a+b}{s-a-b}}$ ,  $a = \frac{1+\alpha^2}{2\alpha} w + L \frac{1-\alpha}{2}$ ,  $b = \frac{1-\alpha^2}{2\alpha} w + L \frac{1+\alpha}{2}$ ,

CARD 2/3

$U(\eta) = 1$  for  $\eta > 0$ ,  $U(\eta) = 0$  for  $\eta < 0$ , F is a certain

On the Stability of a Plane Stationary Shock Wave

40-4-3/24

known function of the initial values,  $J_0$  is the Bessel function. For a gas extending at infinity behind the shock wave it is  $l=\infty$  and the initial perturbations vanish for  $x > h$ . Then instead of (3) it holds:

$$(4) \quad \frac{d\zeta}{ds} = -\mu \int_0^{\infty} U(s-w) J_0(s-w) \left( \frac{d^2\zeta}{dw^2} + \nu\zeta \right) dw + F$$

From this with the aid of Laplace transformations the author obtains the stability conditions and an asymptotic law for the attenuation of the perturbations. The influence of the reflection on the surface of the piston is equally considered.

SUBMITTED: March 12, 1957

AVAILABLE: Library of Congress

CARD 3/3

10(4), 24(3)

SOV/20-121-4-10/54

AUTHOR: Iordanskiy, S. V.

TITLE: Zemplen's Theorem in Magnetic Hydrodynamics (Teorema Tsemplena v magnitnoy gidrodinamike)

PERIODICAL: Doklady Akademii nauk SSSR, 1958, Vol 121, Nr 4, pp 610-612 (USSR)

ABSTRACT: The discontinuities of the shock wave type (where the matter passes through the discontinuity surface and where the thermodynamic quantities are varied) are very interesting. L. D. Landau and Ye. M. Lifshits show in their book (Ref 2) that only compression waves are possible under such conditions (just as in the case of ordinary shock waves). This paper proves this assumption for any values of the discontinuities. First the equations for the discontinuities of the shock wave type are given in an explicit form. The author investigates only substances for which there is

$$\left(\frac{\partial^2 p}{\partial v^2}\right)_S = 0, \left(\frac{\partial p}{\partial S}\right)_V > 0, \quad p \text{ denotes the pressure, } V - \text{ the specific volume,}$$

The above mentioned equations describe any pos-

Card 1/2

Zemplen's Theorem in Magnetic Hydrodynamics

SOV/20-121-4-10/54

sible kind of shock waves in magnetic hydrodynamics. Next, the author deduces an equation for the curves  $p_2(v_2)$  of the Hugoniot (Gyugonio) type. This equation has 3 solutions which, in the case of weak discontinuities, are real solutions and correspond to 3 different Hugoniot curves  $p_2(v_2)$ . But in the case  $H_1 \neq 0$  this equation has only one real radical if  $p_2$  is sufficiently high. The entropy  $S_2$  increases in a monotonous way along any of these Hugoniot curves and there is always  $p_2 > p_1$ . The proof of this assertion is analogous to the corresponding proof of ordinary gas dynamics. The Hugoniot function is then given and discussed.  $dS_2$  does not change its sign along the Hugoniot curve if certain conditions (which are given by the author) are satisfied. The entropy  $S_2$  and the Hugoniot potential have their maximum in the same points. There are 3 references, 3 of which are Soviet.

PRESENTED: March 29, 1958, by M. A. Lavrent'yev, Academician  
SUBMITTED: March 27, 1958  
Card 2/2

FOR DANUSKIY, S.V.

Тема 1: ВОЛН ВЗАИМОДЕЙСТВИЯ

001/762

Ученые магнетронной гидродинимики и динамике плазмы (Problems of Magnetron Hydrodynamics and Plasma Dynamics) Transactions of a Conference Held in the USSR Academy of Sciences, 1959. 343 p. Reprints only. Moscow, 1,000 copies printed.

Sponsoring Agency: Akademiya Nauk Latvriyevoy SSR, Institut Fiziki.

Editorial Board: B.A. Frank-Kamenetskiy, Doctor of Physics and Mathematics, Professor; A.I. Vol'pert, Doctor of Technical Sciences, Professor; I.M. Elm, Doctor of Physics and Mathematics; V.Za. Volyn, Candidate of Physics and Mathematics; V.G. Vitok, Candidate of Physics and Mathematics; Yu.M. Kravtsov, and V.Ia. Kravchenko.

M.I. A. Svyetlov; Tech. M.I. A. Svyetlov

Summary: This book is intended for physicists working in the field of magnetron hydrodynamics and plasma dynamics. It contains the transactions of a conference held in Riga, June 1959, on problems in applied and theoretical magnetohydrodynamics. The articles of the conference were the investigation of the basic trends in theoretical and applied magnetohydrodynamics, establishing contact between the people doing research in different branches of magnetohydrodynamics, and promoting the participation of theoretical physicists in problems in applied magnetohydrodynamics. More than 150 persons from different parts of the Soviet Union took part in the conference, and 55 papers were read. Similar conferences are held regularly in the future; the next such conference is scheduled to be held in Riga in June 1960. In this present collection of articles, the authors present the results of their work. The book is divided into two parts: the first part deals with problems in theoretical magnetohydrodynamics and plasma dynamics, and consists of 35 articles on such aspects of the problem as the solution of magnetohydrodynamics in astrophysics (D.A. Frank-Kamenetskiy), magnetohydrodynamics and the investigation of cosmic-ray variations (L.I. Borzov), neutralization of plasmas in a magnetic field (G.V. Gerasimov and E.I. Oshinov), stability of shock waves and magnetohydrodynamics (A.Y. Akhiezer). The second part, consisting of 35 articles, deals with problems of experimental magnetohydrodynamics, including the application of jet-like simulation for investigation of electromagnetic processes in liquid metals (M. Kikini) and the simulation of electromagnetic processes in liquid metals (M. Kikini) in the laboratory of Science of Scientific Academy USSR. Several articles are devoted to induction lamps, electromagnetic crosslinks, electromagnetic stirrers for molten metals, and their application in the metallurgical industry including schematic diagrams of their power-supply systems. References are given at the end of most of the articles.

Svyetlov, M.I. On the Stability of Shock Waves in Magnetohydrodynamics	127
Polurin, B.V., and G.Ia. Lyubchenko. The Impossibility of Baroclinic Shock Waves in Magnetohydrodynamics	132
Lordashvili, B.V. Szepfen's Theorem in Magnetohydrodynamics	133
Kiselev, M.I., and V.I. Isakovich. Oblique Shock Wave in Plasma With Finite Conductivity and Certain Other Problems in Shock-Wave Theory	135
Shubelkin, I.I. Thermodynamic Equilibrium of Strong Discontinuity Surfaces	136
Akhiezer, A.I., and A.G. Sitenko. On the Theory of Hydromagnetic Wave Excitation	137
Borsov, A.I. Generation of Magnetic Sound Waves	141
Sitenko, A.G., and Yu.A. Kirichkin. Dissipation of Hydromagnetic Waves in Turbulent Fluctuations	143
Card 6/12	11



10 (2)

AUTHOR:

Iordanskiy, S. V.

SOV/20-125-6-9/61

TITLE:

On the Asymptotic Form of the Axially Symmetric Expanding Wave  
in a Heavy Fluid (Ob asimptotike osesimmetrichnoy  
raskhodyashcheyasya volny v tyazheloy zhidkosti)

PERIODICAL:

Doklady Akademii nauk SSSR, 1959, Vol 125, Nr 6,  
pp 1211-1214 (USSR)

ABSTRACT:

The present paper deals with the asymptotic form of an  
outgoing wave at large distances from the symmetry axis on the  
assumption of axial symmetry. The depth of the fluid is  
assumed to be finite, and the motion of the fluid as being a  
potential motion. For the velocity potential it holds that

$\frac{\partial^2 \psi}{\partial z^2} + \Delta \psi = 0$ , where the z-axis is perpendicular in an upward

direction. The boundary conditions on the bottom, which is  
assumed to be plane, is  $\frac{\partial \psi}{\partial z} = 0$  with  $z = 0$ . On the free surface

the following conditions must be satisfied:

$\frac{\partial \psi}{\partial t} + \frac{1}{2} (\nabla \psi)^2 + \frac{1}{2} \left( \frac{\partial \psi}{\partial z} \right)^2 + g\zeta = F(t)$  (Bernoulli-equation) and

Card 1/3

On the Asymptotic Form of the Axially Symmetric  
Expanding Wave in a Heavy Fluid

SOV/20-125-6-9/61

$\frac{\partial \xi}{\partial t} + \nabla \xi \cdot \nabla \psi = \frac{\partial \psi}{\partial z}$  (kinematic condition). Here  $\nabla$  denotes the gradient in the  $x, y$ -plane. The required equations are developed by expansion of  $\psi$  with respect to powers of  $z$ . By using the condition which holds for the bottom,

$$\psi = \psi_0(x, y, t) - \frac{1}{2} z^2 \Delta \psi_0 + \frac{1}{24} z^4 \Delta \Delta \psi_0 + \dots$$

is obtained by means of the Laplace equation. This representation for  $\psi$  is correct if the characteristic length  $L$  in the  $x, y$ -plane is much greater than  $\xi$ , which is assumed in this case. The author then introduces dimensionless variables and obtains an approximated equation for  $\psi$ . He then endeavors to find the waves which satisfy the conditions given here and the form of which is asymptotically stable at  $t \rightarrow \infty$ . On this occasion, the waves in the channel  $\psi = \psi(x, t)$  are first investigated. In first approximation there exist waves of arbitrary shape  $\psi_1 = f(x-t)$ . The author then puts  $\psi = f(x-t) + \psi_2$ , where  $\psi_2$  is a high-order small quantity. In  $\psi_2$  a secular term

Card 2/3

On the Asymptotic Form of the Axially Symmetric  
Expanding Wave in a Heavy Fluid

SOV/20-125-6-9/61

occurs, which increases infinitely with increasing  $\eta = x + t$ . Thus, the terms of higher order may, after a sufficiently long time, completely change the shape of the original current wave. In conclusion, an asymptotic representation for the velocity of the fluid in a wave is written down. In a similar manner it is possible to investigate the asymptotic form of waves in a fluid of variable depth if this depth varies sufficiently slowly. There are 4 Soviet references.

ASSOCIATION: Matemicheskii institut im. V. A. Steklova Akademii nauk  
SSSR (Mathematics Institute imeni V. A. Steklov of the  
Academy of Sciences, USSR)

PRESENTED: December 31, 1958, by M. A. Lavrent'yev, Academician

SUBMITTED: December 27, 1958

Card 3/3

16(1)

SOV/20-127-3-7/71

AUTHOR:

Iordanskiy, S.V.

TITLE:

A Solution of the Cauchy Problem for the Kinetic Equation of Electron Plasma

PERIODICAL:

Doklady Akademii nauk SSSR, 1959, Vol 127, Nr 3, pp 509-512 (USSR)

ABSTRACT:

According to Landau [Ref 1] the kinetic equation for an electron plasma without collisions (the magnetic field is equal to zero) can be written in the form

$$(1) \quad \frac{\partial n}{\partial t} + v \frac{\partial n}{\partial x} - \frac{e}{m} E(x,t) \frac{\partial n}{\partial v} = 0 ,$$

where E is determined from

$$(2) \quad \frac{\partial E}{\partial x} = -4\pi e \left\{ \int_{-\infty}^{\infty} n(v,x,t) dv - N_0 \right\} ;$$

here e and m are charge and mass of the electron,  $N_0$  the density of the positive ions assumed to be constant. For (1) - (2) the author poses the Cauchy problem

Card 1/3

A Solution of the Cauchy Problem for the Kinetic  
Equation of Electron Plasma

SOV/20-127-3-7/71

$$n|_{t=0} = f(x, v) \quad , \quad \lim_{x \rightarrow -\infty} E = 0$$

where  $f(x, v) > 0$  is a continuous function.

Theorem: The solution of (1) - (2) with these conditions exists for every continuous  $f(x, v)$  which satisfies the conditions

$$\int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} f(x, v) dv - N_0 \right\} dx = 0$$

$$|f(x, v) - N(v)| < K(v) \varphi(x) \quad (0 < N(v) < K(v))$$

where

$$N(v) = \lim_{x \rightarrow \pm \infty} f(x, v) \quad \left( N_0 = \int_{-\infty}^{\infty} N(v) dv \right)$$

$\varphi(x)$  is bounded and  $K(v)$  decreases monotonely with increasing  $|v|$  so that

Card 2/3

A Solution of the Cauchy Problem for the Kinetic  
Equation of Electron Plasma

SOV/20-127-3-7/71

$$\int_{-\infty}^{\infty} v^2 K(v) dv < \infty, \quad \int_{-\infty}^{\infty} \varphi(x) dx < \infty$$

The solution is unique in the class of bounded functions satisfying the Lipschitz condition on the whole  $x$ -axis, vanishing for  $x \rightarrow \pm \infty$  and possessing a continuous partial derivative with respect to  $x$ .

There are 2 Soviet references.

ASSOCIATION: Matematicheskiy institut imeni V.A. Steklova Akademii nauk  
SSSR (Mathematical Institute imeni V.A. Steklov, AS USSR)

PRESENTED: April 10, 1959, by M.A. Lavrent'yev, Academician

SUBMITTED: April 1, 1959

Card 3/3

IORDANSKIY, S.V. (Moskva)

Equations of motion for a liquid containing gas bubbles.  
PMTF no.3:102-110 S-0'60. (MIRA 14:7)  
(Differential equations, Partial)  
(Hydrodynamics)

10.3200

27791  
S/508/60/028/000/002/022  
D237/D305

26.5200

**AUTHORS:** Iordanskiy, S.V., and Shmyglevskiy, Yu.D. (Moscow)

**TITLE:** Sublimation of an axially symmetric blunt body near the stagnation point of incident gas flow

**PERIODICAL:** Akademiya nauk SSSR. Otdeleniye tekhnicheskikh nauk. Inzhenernyy sbornik, v. 28, 1960, 26 - 35

**TEXT:** The authors obtain here the equations of an axially symmetric laminar boundary layer for a 2-component gas at low temperatures with diffusion present. Boundary conditions are derived for the case of sublimation, and the method is given for calculating sublimation flow and velocity near the stagnation point. Finally solid CO<sub>2</sub> in the stream of air is considered as an example. According to L.D. Landau and Ye.M. Livshits (Ref. 2: Mekhanika sploshnykh sred (Mechanics of Continuous Media) Gostekhizdat, M. 1954) the flow of multi-component gas is described by

Card 1/9



Sublimation of an axially ...

27791  
S/508/60/028/000/002/022  
D237/D305

$$\left. \begin{aligned} \frac{\partial \rho w_l}{\partial x_l} &= 0, \\ \frac{\partial \rho c_\alpha w_l}{\partial x_l} + \frac{\partial j_{l\alpha}}{\partial x_l} &= 0, \\ \rho w_k \frac{\partial w_l}{\partial x_k} &= -\frac{\partial p}{\partial x_l} + \frac{\partial \sigma_{lk}}{\partial x_k}, \\ \frac{\partial}{\partial x_l} \left[ \rho \left( \frac{w^2}{2} + h \right) w_l - w_k \sigma_{lk} + q_l \right] &= 0, \end{aligned} \right\} \quad (1.1)$$

$$\rho = \rho(p, T, c_1, c_2, \dots). \quad (1.2)$$

$$j_{i\alpha} = -\rho D_\alpha \left( \frac{\partial c_\alpha}{\partial x_i} + \frac{k_T^{(\alpha)}}{T} \frac{\partial T}{\partial x_i} + \frac{k_p^{(\alpha)}}{p} \frac{\partial p}{\partial x_i} \right), \quad (1.3)$$

$$q_i = [k_T^{(\alpha)} M_\alpha - T M_\alpha' + \mu_\alpha] j_{i\alpha} - \eta \frac{\partial T}{\partial x_i},$$

Card 2/9

Sublimation of an axially ...

27791  
S/508/60/028/000/002/022  
D237/D305

$$d_{1k} = \eta \left( \frac{\partial w_1}{\partial x_k} + \frac{\partial w_k}{\partial x_1} - \frac{2}{3} \delta_{1k} \frac{\partial w_l}{\partial x_l} \right) + \zeta \delta_{1k} \frac{\partial w_l}{\partial x_l}; \quad (1.3)$$

For two-component gas (1.1) and (1.3) are transformed into cylindrical coordinates by  $x_1 = x$ ,  $x_2 = x \cos \vartheta$ ,  $x_3 = r \sin \vartheta$  and the equations of axial flow in  $(x, r)$  plane are derived in an  $(s, n)$  orthogonal coordinate system associated with the surface AB of the body (Fig. 1). The partials are then

$$\frac{\partial}{\partial x} = \frac{R \cos \gamma}{R+n} \frac{\partial}{\partial s} - \sin \gamma \frac{\partial}{\partial n}, \quad \frac{\partial}{\partial r} = \frac{R \sin \gamma}{R+n} \frac{\partial}{\partial s} + \cos \gamma \frac{\partial}{\partial n}.$$

where  $R$  = radius of curvature,  $\gamma$  = angle between tangent to AB and  $x$ -axis at the given point. Tangential and normal velocities  $u$  and  $v$  are given by

$$w_r = u \sin \gamma + v \cos \gamma, \quad w_x = u \cos \gamma - v \sin \gamma.$$

Then for a small velocity of sublimation

Card 3/9

44

Sublimation of an axially ...

27791  
S/508/60/028/000/002/022  
D237/D305

$$\begin{aligned}
 & \frac{\partial r \rho u}{\partial s} + \frac{\partial r \rho v}{\partial n} = 0, \\
 & \rho u \frac{\partial c}{\partial s} + \rho v \frac{\partial c}{\partial n} = \frac{\partial}{\partial n} \rho D \left( \frac{\partial c}{\partial n} + \frac{k_T}{T} \frac{\partial T}{\partial n} \right), \\
 & \rho u \frac{\partial u}{\partial s} + \rho v \frac{\partial u}{\partial n} = -\frac{dp}{ds} + \frac{\partial}{\partial n} \eta \frac{\partial u}{\partial n}, \\
 & \frac{\partial p}{\partial n} = 0 \text{ или } p = p(s), \\
 & \rho u \frac{\partial}{\partial s} \left( h + \frac{u^2}{2} \right) + \rho v \frac{\partial}{\partial n} \left( h + \frac{u^2}{2} \right) = \frac{\partial}{\partial n} \left\{ \eta \frac{\partial}{\partial n} \frac{u^2}{2} + \right. \\
 & \left. + \kappa \frac{\partial T}{\partial n} + \rho D (h_\alpha - h_\beta + k_T M) \left( \frac{\partial c}{\partial n} + \frac{k_T}{T} \frac{\partial T}{\partial n} \right) \right\},
 \end{aligned} \tag{1.4}$$

is obtained, where  $M = M_\alpha + M_\beta$ . For low temperature work in the absence of chemical reactions,

$$p = mR \left( \frac{c}{m_\alpha} + \frac{1-c}{m_\beta} \right) \rho T, \quad h = c h_\alpha + (1-c) h_\beta,$$

*Handwritten mark*

Card 4/9

27791

S/508/60/028/000/002/022  
D237/D305

Sublimation of an axially ...

can be utilized, where  $c$  is independent of  $\xi$  and  $T$ . Boundary conditions when gas  $\alpha$  flows around a body  $\beta$  are for  $\lambda = \infty$ ,

$$f_\lambda(\xi, \infty) = 1 \quad (2.1), \quad T(\xi, \infty) = T_e(\xi), \quad c(\xi, \infty) = c_e(\xi), \quad (2.2)$$

where  $T_e(\xi)$  and  $c_e(\xi)$  are the temperature and concentration of  $\alpha$  and for  $\lambda = 0$

$$f_\lambda(\xi, 0) = 0 \quad (2.3), \quad T(\xi, 0) = T_w(p_e(\xi)) \quad (2.9)$$

and

$$\left. \begin{aligned} & \left[ (2f_\lambda + f)c + \frac{L}{P} \left( c_\lambda + \frac{k_T}{T} T_\lambda \right) \right]_{\lambda=0} = 0, \\ & \left[ (2f_\lambda + f)(Q - ck_T M) + \frac{c_p}{P} T_\lambda \right]_{\lambda=0} = \frac{q_T \sqrt{2\xi}}{r u_p \gamma_w} \end{aligned} \right\} \quad (2.11)$$

where  $Q = [h_\beta(T_w)]_{+0} - [h_\beta(T_w)]_{-0}$  = heat of sublimation of unit mass of  $\beta$  at the temperature  $T_w$ . Flow near the axis of symmetry is Card 5/9

27791

S/508/60/028/000/002/022  
D237/D305

Sublimation of an axially

solved where the solution can be expressed in the form of a power series in  $\sqrt{\xi}$  with coefficients dependent on  $\lambda$ , if  $p_e(\xi)$  can also be expanded in powers of  $\sqrt{\xi}$ . Terms independent of  $\xi$  will then give a solution on the axis of symmetry. In dimensionless magnitudes

$$l = \frac{T}{T_w|_{\xi=0}}, \quad H = \frac{m_a(h_a - h_b)}{mRT_w|_{\xi=0}}, \quad \gamma = \frac{m_a c_p}{mR},$$

$$\Phi = lf_{\lambda\lambda}, \quad F = f_{\lambda}, \quad K = \frac{Ll}{P} \left( c_{\lambda} + \frac{k_T}{T} T_{\lambda} \right)$$

$$E = \frac{l\gamma}{P} t_{\lambda} + K(H + k_T M).$$

$$\Phi_{\lambda} = -l \frac{\Phi}{T} - \frac{1}{2} \left( \frac{p_e}{P} - F^2 \right),$$

$$F_{\lambda} = \frac{\Phi}{T}, \quad f_{\lambda} = F,$$

(3.1)

Card 6/9

27791

S/508/60/028/000/002/022  
D237/D305

Sublimation of an axially ...

$$\begin{aligned}
 t_\lambda &= \frac{P}{L_T} \left\{ E - K \left[ H + k_T \frac{m_p(1-c) + m_s e}{m_p e(1-c)} t \right] \right\}, \\
 c_\lambda &= \frac{PK}{L_T} - \frac{k_T}{t} t_\lambda, \quad K_\lambda = -f c_\lambda, \\
 E_\lambda &= -f(H c_\lambda + \gamma t_\lambda).
 \end{aligned}
 \tag{3.1}$$

is obtained and the boundary conditions (2.1)-(2.3), (2.9) and (2.11) become

$$\left. \begin{aligned}
 F(0) &= 0, \quad t(0) = 1, \quad K(0) = -f(0)c(0), \\
 E(0) &= -f(0)[\bar{Q} + H(0)c(0)], \\
 F(\infty) &= 1, \quad t_\infty = \frac{T_e}{T_w}, \quad c(\infty) = 1,
 \end{aligned} \right\}
 \tag{3.2}$$

where

$$\bar{Q} = \frac{m_\alpha Q}{mRT_w} \Big|_{\xi=0}, \quad Q = \frac{mRT_w^2}{p_e} \frac{dp_e}{dT_w}.$$

Card 7/9

*HT*

27791

S/508/60/028/000/002/022  
D237/D305

Sublimation of an axially ...

For flow without diffusion (3.1) can be used if its 5th and 6th equations are replaced by

$$K(\lambda) = 0, c = \begin{cases} 0 & \text{when } f < 0, \\ 1 & \text{when } f > 0, \end{cases}$$

and for the flow without sublimation (3.1) can be used with the boundary conditions

$$F(0) = 0, t(0) = 1, f(0) = 0, c(0) = 1,$$

$$F(\infty) = 1, t(\infty) = T_e/T_w, c(\infty) = 1,$$

where  $T_w$  = given temperature. The problem of the flow of air  $M = 6.2$  around the body composed of solid  $\text{CO}_2$  is solved as an example. There are 3 figures and 11 references: 3 Soviet-bloc and 8 non-Soviet-bloc. The 4 most recent references to the English-language publications read as follows: J.A. Fay, R.F. Riddel, Theory of Stagnation Point Heat Transfer in Dissociated Air, J. Aeron. Sci. vol. 25, No. 2, 1958; Tables of Thermal Properties of Gases, US. Department of Commerce National Bureau of Standards, Circular 564.

Card 8/9





Cauchy's problem for a kinetic equation ...

<sup>29898</sup>  
S/517/61/060/000/004/009  
B112/B125

generalizes his results for the case of a plasma with more than one components. There are 5 references: 4 Soviet and 1 non-Soviet.

UX

Card 2/2

24, 2190 (1049, 1141)  
26. 2330

S/057/61/031/005/006/020  
B104/B205

AUTHOR: Iordanskiy, S. V.

TITLE: Electron oscillations of plasma between two electrodes

PERIODICAL: Zhurnal tekhnicheskoy fiziki, v. 31, no. 5, 1961, 549-556

TEXT: The author studied the stability of an electron plasma between two plane, infinitely large electrodes, one of which is traversed by a beam of electrons. A similar theoretical study was performed by Bohm et al. (Phys. Rev., 79, 992, 1950). Looney et al. (Phys. Rev., 93, 915, 1954) obtained experimental results which agree more or less with the data found here. In his experiments, the present author proceeded from the assumption of a high oriented electron velocity compared to the thermal velocities in the electron beam and in the plasma. If also the characteristic dimension is supposed to be large compared to the Debye radius, the problem can be treated as a problem of "cold plasma" in hydrodynamic approximation. The author confines himself to one-dimensional oscillations and proceeds from the system

Card 1/8

S/057/61/031/005/006/020  
B104/B205

Electron oscillations...

$$\left. \begin{aligned} \frac{\partial N}{\partial t} + \frac{\partial}{\partial x} NV &= 0, \\ \frac{\partial V}{\partial t} + V \frac{\partial V}{\partial x} &= -\frac{e}{m} \frac{\partial \varphi}{\partial x}, \\ \frac{\partial n}{\partial t} + \frac{\partial}{\partial x} nv &= 0, \\ \frac{\partial v}{\partial t} + v \frac{\partial v}{\partial x} &= -\frac{e}{m} \frac{\partial \varphi}{\partial x}, \\ \frac{\partial^2 \varphi}{\partial x^2} &= -4\pi e (n + N - N_+) \end{aligned} \right\} (1)$$

where  $N$  and  $V$  stand for the density and velocities, respectively, of the electrons in the beam;  $n$  and  $v$  are the analogous quantities in the plasma;  $N_+$  is the given density of positive ions;  $\varphi$  is the potential of the electric field,  $e$  is the electron charge, and  $m$  is the electron mass. In the following, the author considers only the case where a given potential,  $-\varphi_1$ , which is negative with respect to the plasma, is applied to the

Card 2/8

S/057/61/031/005/006/020  
B104/B205

Electron oscillations...

electrodes A and B. In this case, ion layers are formed on the electrodes, in which there are no plasma electrons if  $e\varphi_1 \gg kT_e$ . System (1) has steady solutions so that outside the ion layers the plasma is electrically neutral; the quantities  $N=N_0$ ,  $n=n_0$ ,  $V=V_0$  are constant and  $\partial\varphi/\partial x = v = 0$ . Inside the ion layers, these quantities are a function of  $x$ . Provided the plasma potential is equal to zero ( $mV^2/2 \gg e\varphi_1$ ),  $N$ ,  $n$ , and  $V$  are also constant inside the ion layers. With an ion layer of thickness  $\delta$  and an electrode spacing  $l$ , the problem consists in examining the small perturbations of the steady solutions of (1)... One obtains  $N=N_0+N'$ ,  $V=V_0+V'$ ,  $n=n_0+n'$ ,  $v=v'$ , and  $\varphi=\varphi_0+\varphi'$ . System (1) is linearized, and for the interior of the ion layers one finds the system

$$\left. \begin{aligned} \frac{\partial N'}{\partial t} + \frac{\partial}{\partial x}(N_0 V' + V_0 N') &= 0, \\ \frac{\partial V'}{\partial t} + V_0 \frac{\partial V'}{\partial x} &= -\frac{e}{m} \frac{\partial \varphi'}{\partial x}, \\ \frac{\partial^2 \varphi'}{\partial x^2} &= -4\pi e N' \end{aligned} \right\} \quad (3)$$

Card 3/8

22775

Electron oscillations...

S/057/61/031/005/006/020  
B104/B205

For the interior of the plasma, an analogous system is obtained in the same way. When solving these linearized systems, it is necessary to account for the altered boundary conditions

$$\left. \begin{aligned} \varphi|_{z=0} + \frac{\partial \varphi}{\partial x} \Big|_{z=0} \zeta_A &= \varphi|_{z+\delta} + \frac{\partial \varphi}{\partial x} \Big|_{z+\delta} \zeta_A, \\ \frac{\partial \varphi}{\partial x} \Big|_{z=0} + \frac{\partial^2 \varphi}{\partial x^2} \Big|_{z=0} \zeta_A &= \frac{\partial \varphi}{\partial x} \Big|_{z+\delta} + \frac{\partial^2 \varphi}{\partial x^2} \Big|_{z+\delta} \zeta_A. \end{aligned} \right\} \quad (A)$$

It is shown that the problem concerning the stability of the steady solution results in the determination of eigenvalues  $\lambda$  at which solutions to the above-mentioned linearized systems exist in the form  $e^{\lambda t} f(x)$ . These solutions satisfy the boundary conditions

$$\left. \begin{aligned} \varphi|_{z=0} &= \varphi|_{z+\delta}, \quad \frac{\partial \varphi}{\partial x} \Big|_{z=0} + 4\pi n_0 \zeta_A = \frac{\partial \varphi}{\partial x} \Big|_{z+\delta}, \\ N'|_{z=0} &= N'|_{z+\delta}, \quad V'|_{z=0} = V'|_{z+\delta}, \quad \frac{d\zeta_A}{dt} = \psi' \Big|_{z+\delta}. \end{aligned} \right\} \quad (5)$$

Card 4/8

S/057/61/031/005/006/020  
B104/B205

Electron oscillations...

$$\left. \begin{aligned} \varphi'|_{l-\delta+0} &= \varphi'|_{l-\delta-0}, \quad \frac{\partial \varphi'}{\partial x}|_{l-\delta+0} + 4\pi e n_0 \epsilon_B = \frac{\partial \varphi'}{\partial x}|_{l-\delta-0}, \\ N'|_{l-\delta+0} &= N'|_{l-\delta-0}, \quad V'|_{l-\delta+0} = V'|_{l-\delta-0}, \quad \frac{d\zeta_B}{dt} = \nu'|_{l-\delta-0}. \end{aligned} \right\} (6)$$

and

$$\left. \begin{aligned} \varphi'|_0 &= \varphi'|_l = 0 \quad (\text{заданный потенциал электродов}), \\ N'|_0 &= V'|_0 = 0 \quad (\text{заданный пучок на электроде } A). \end{aligned} \right\} (7)$$

If the desired functions which are multiplied by  $e^{-\lambda t}$ , are indicated by the same letters but without primes, the following systems of ordinary differential equations will be obtained for both regions:

$$\left. \begin{aligned} \lambda N + V_0 \frac{dN}{dx} + N_0 \frac{dV}{dx} &= 0, \\ \lambda V + V_0 \frac{dV}{dx} + \frac{e}{m} \frac{d\varphi}{dx} &= 0, \\ \frac{d^2 \varphi}{dx^2} &= -4\pi e N \end{aligned} \right\} (8)$$

$(0 \leq x \leq \delta, l - \delta \leq x \leq l);$

Card 5/8

Electron oscillations...

S/057/61/051/005/006/020  
B104/B205

and

$$\left. \begin{aligned} \lambda N + V_0 \frac{dN}{dx} + N_0 \frac{dV}{dx} &= 0, \\ \lambda V + V_0 \frac{dV}{dx} + \frac{e}{m} \frac{d\varphi}{dx} &= 0, \\ \lambda v &= -\frac{e}{m} \frac{d\varphi}{dx}, \\ \lambda n + n_0 \frac{dv}{dx} &= 0, \\ \frac{d^2\varphi}{dx^2} &= -4\pi e (N + n) \end{aligned} \right\} \quad (8)$$

$(l - \delta \gg x \gg \delta).$

By eliminating the quantities  $V$ ,  $\varphi$ ,  $n$ , and  $v$  from these systems,  $N$  is given by the expressions

$$\begin{aligned} N &= B_1 e^{i\alpha_1 x} + B_2 e^{i\alpha_2 x} \quad (0 \leq x \leq \delta), \\ N &= A_1 e^{i\alpha_1 x} + A_2 e^{i\alpha_2 x} \quad (l - \delta \leq x \leq l), \\ N &= C_1 e^{i\alpha_1 x} + C_2 e^{i\alpha_2 x} \quad (\delta \leq x \leq l - \delta), \end{aligned}$$

where

$$\alpha_{1,2} = -\frac{\lambda}{V_0} \pm \frac{i\omega}{V_0}, \quad \alpha_{1,2} = -\frac{\lambda}{V_0} \pm \frac{i\omega}{V_0 \sqrt{1 + \frac{\Omega^2}{\lambda^2}}} \quad (12)$$

Card 6/8

S/057/61/031/005/006/020  
B104/B205

Electron oscillations...

Then, it is possible to derive the corresponding relations for the other desired quantities,  $N$ ,  $V$ , and  $\varphi$ , with the aid of which the eigenvalues  $\lambda$  are found from the boundary conditions for the present problem. This is, as a rule, a very cumbersome procedure. For  $N=0$  the eigenvalues are given by  $\lambda = \pm i\Omega\sqrt{2\delta/1}$  (16).  $\lambda = \pm i\Omega$  is an eigenvalue of infinitely multiple degeneracy. For  $N_0 \neq 0$  there are two types of eigenvalues. The first type corresponds to plasma oscillations, and for  $N_0 \rightarrow 0$   $\lambda$  is given by (16). The second type corresponds to internal plasma oscillations, and for  $N_0 \rightarrow 0$   $-\lambda^2$  tends toward the square of the plasma frequencies. The most interesting oscillations occur if

$$\left. \begin{aligned} \lambda^2 + \Omega^2 &= -\frac{\omega^2}{\beta^2}, \\ \alpha_{1,2} &= -\frac{\lambda}{V_0}(1 \mp \beta), \end{aligned} \right\} \quad (17)$$

$\beta$  tends to a constant value if  $\omega \rightarrow 0$ . The existence of plasma boundaries does not alter the principal results obtained in hydrodynamic approxima-

Card 7/8



Electron oscillations...

S/057/61/031/005/006/020  
B104/B205

tion for the instability of steady states in an unbounded plasma in the presence of an electron beam. The further results obtained here are in qualitative agreement with those obtained by Looney et al. A quantitative comparison is not possible since the electron density in the beam was much higher than in the plasma. Yu. L. Klimovich is thanked for a discussion of several interesting problems. There are 3 figures and 4 references: 2 Soviet-bloc and 2 non-Soviet-bloc.

ASSOCIATION: Matematicheskiy institut im. V. A. Steklova Moskva  
(Institute of Mathematics imeni V. A. Steklov, Moscow)

SUBMITTED: July 20, 1960

Card 8/8

IODANSKIY, S.V.

Excitati n of magnetoacoustic waves in a conducting fluid.  
Dokl. AN SSSR 146 no.3:557-560 S '62. (MIRA 15:10)

1. Matematicheskiy institut im. V.A.Steklova AN SSSR. Predstavleno  
akademikom L.I.Sedovym.  
(Magnetohydrodynamics)

14 1200 (2119) (5105)

111751  
S/057/63/033/001/013/017  
B125/B186

AUTHOR: Iordanskiy, S. V.

TITLE: On the resonance excitation of waves in an infinitely conducting liquid

PERIODICAL: Zhurnal tekhnicheskoy fiziki, v. 33, no. 1, 1963, 105 - 114 f

TEXT: The characteristic longitudinal waves close to resonance in an infinitely well conducting liquid are studied in magnetohydrodynamic approximation. This problem is closely analogous to that treated in gas dynamics (G. Bechov, Phys. Fluids, 1, no. 3, 205, 1958).  $H_x, H_y, H_z$  are the components of the magnetic field strength;  $v_x, v_y, v_z$  the components of the velocity vector;  $\rho$  the density of the liquid;  $c^2 = (\partial p / \partial \rho)_s$  the square of the sonic velocity. The smallness of the external exciting field makes it possible to write  $\vec{H} = \vec{H}_0 + \vec{h}$ ,  $\rho = \rho_0 + \rho'$ , with  $|\vec{H}_0| \gg |\vec{h}|$ ,  $\rho_0 \gg \rho'$ , etc. The right-hand sides of the magnetohydrodynamic equations are considered after separation of the linear part of the perturbation. The solution of

Card 1/4

S/057/63/033/001/013/017  
B125/B186

On the resonance excitation of waves ...

the separated linear equations consists of a slow and a fast acoustic wave with velocities

$$u_{1,2} = \frac{1}{2} \left\{ \sqrt{c_0^2 + \frac{H_0^2}{4\pi\rho_0} + \frac{H_{0x}c_0}{\sqrt{\pi\rho_0}}} \pm \sqrt{c_0^2 + \frac{H_0^2}{4\pi\rho_0} - \frac{H_{0x}c_0}{\sqrt{\pi\rho_0}}} \right\} \quad (1.3) \text{ and an } \checkmark$$

Alfven wave with velocity  $u_3 = H_{0x} / \sqrt{4\pi\rho_0}$ . Each of these three waves is characterized by arbitrary functions  $\Lambda_1^\pm, \Lambda_2^\pm, \Lambda_3^\pm$ . The boundary conditions

$$\left. \begin{aligned} \Lambda_i^+ - \mu\Lambda_i^- &= 0 \quad (x=l); \quad \Lambda_i^+ - \mu\Lambda_i^- = \epsilon \cos \omega t \quad (x=0), \\ \sum_k (\alpha_k \Lambda_k^+ + \beta_k \Lambda_k^-) &= 0, \quad (x=l, x=0, k \neq i, j \neq i). \end{aligned} \right\} \quad (1.4)$$

can cause a deviation of the spectrum from the integral multiple of any of the fundamental frequencies. Close to the resonance the next higher corrections need consideration, for which  $|\Lambda_i| \gg \epsilon$  and  $|\Lambda_i| \gg |\Lambda_j|$ ,  $\Lambda_i$  and  $\Lambda_j$  being invariants.

The perturbation traverses the distance  $2l$  in the time  $\tau$ , derived by

$$\begin{aligned} \tau &= \int_{S_1} \frac{dx}{u_i + \alpha L^+ + \beta L^-} - \int_{S_2} \frac{dx}{u_i + \alpha L^- + \beta L^+} \approx \\ &\approx \frac{2l_0}{u_i} + \frac{2\Delta l}{u_i} - \frac{\alpha(1+\mu)l_0}{u_i^2} L^-(t) - \frac{\beta(1+\mu)l_0}{u_i^2} \int_0^{l_0} L^-\left(t + \frac{2\pi x}{u_i j_0}\right) dx. \end{aligned}$$

Card 2/4