

ACC NR: AP6033538 SOURCE CODE: UR/0170/66/011/004/0516/0520

AUTHOR: Lyubchenko, A. P.; Tsarina, I. V.; Sherman, D. G.; Shukhov, A. S.

ORG: Transportation Machinery Plant, Khar'kov (Zavod transportnogo mashino-stroyeniya)

TITLE: Method of determining temperature fields of machinery-part surfaces inaccessible during operation

SOURCE: Inzhenerno-fizicheskiy zhurnal, v. 11, no. 4, 1966, 516-520

TOPIC TAGS: temperature, temperature dependence, temperature measurement, diffusion method, diffusion parameter, temperature field

ABSTRACT: A method is proposed for determining the temperatures and topologies of the temperature fields of objects which are inaccessible during operation. It is based on the use of the critical dependence of the diffusion parameters of materials on temperature. The method was tested on simple and complex multicomponent heterophase alloys over a wide temperature range (the lowest temperature was 200C). The alloys tested were 65G, Kh12M, Kh18N9T, EI283, and AK-4 grades, with partial reference made to parts of internal

Card 1/2

UDC: 536.5

1971, p. 7.

"An Atomic File for 'Sensitized' Individuals," *Doc. A. 6211*, 61, No 7, 1970

Int. J. Radiology, 4, 1970

"A Simple Derivation of the Potential Energy and the Elasticity Theory for a Plane
Having a Finite Number of Nonlinear Intersections," *Proc. R. Soc. Lond.*, 1967

Inst. of Mathematics, A.S. USSR

1. Introduction

For the solution of the first boundary value problem of the theory of elastic plane stress - an example of the theory of elasticity," *Tr. AN SSSR*, no. 1, 1960.

Inst. of Mathematics, AS Uzb.

"On the problem of the static theory of the elasticity of infinitely
connected plates - on the solution of the second fundamental problem of the static
plane theory of elasticity," Dokl. Akad. Nauk, No. 1, 1940.

Prof. G. G. Gerasimov, Moscow

1. 1. 1.

"A Danish newspaper article's author," *Dan. Ad. Blad*, 1, 1940.

Incl. info. from, AD 1001.

1. Introduction

"A New Solution to the Problem of the Flexure of the Elasticity Theory
for an Anisotropic Body," *Dokl. Akad. Nauk, No. 11, 1961*

Inst. Solid State Phys., Acad. Sci. USSR

1971, p. 1.

"Mechanism of the reaction of a blast of air on the internal structure of
a rock," *Ann. N.Y. Acad. Sci.*, 1971, 1971

Inst. Seismology, Acad. Sci. USSR

SHERMAN, D.I.

Sherman, D. I. On the propagation of waves in a fluid layer covering an elastic half-space. Acad. Sci. URSS. Publ. [Trudy] Inst. Séismolog. no. 115, 43 pp. (1945). (Russian)

Dans ce mémoire l'auteur étudie le problème de détermination des ondes superficielles de Rayleigh sur les limites des milieux considérés. Le problème est effectivement résolu dans le cas particulier de présence d'une force périodique superficielle appliquée sur la surface de séparation de deux milieux ou sur la limite supérieure de la couche fluide. Les formules obtenues sont discutées, et les oscillations principale et résiduelle, à des distances suffisamment grandes des points d'application de la force périodique superficielle, sont calculées.

V. A. Kostitzin (Paris).

SMW
K

SHERMAN, B. I.

"Concerning Certain Problems of the Static Theory of Elasticity for the Half-Space \sqrt{z} e^z and for Two Interconnected Half-Spaces \sqrt{z} e^z with Different Elastic Properties." Iz. Ak. Nauk SSSR, Otdel. Tekh. Nauk, No. 9, 1945. Submitted 7 Jun 1945.

Report U-1582, 6 Dec 1951.

SHERMAN, D.

Sherman, D. On the reduction of the plane problem of the theory of potential to an integral equation. Bull. Acad. Sci. URSS. Sér. Math. [Izvestia Akad. Nauk SSSR] 9, 357-362 (1945). (Russian. English summary)

Let S be a finite plane domain whose boundary L consists of a finite number of closed curves of continuous curvature. The author considers the problem of determining a function $u(x, y)$ harmonic in S and satisfying on L a differential condition of the form $a(s)\partial u/\partial x + b(s)\partial u/\partial y + c(s)u = f(s)$, or more generally of the form

$$\sum_{k=0}^m \sum_{j=0}^k a_{kj}(s) \frac{\partial^k u}{\partial x^k \partial y^j} = f(s),$$

where $a_{kj}(s)$ and $f(s)$ are suitably restricted functions of the arc length; he reduces the problem to the solution of a Fredholm equation. *E. F. Beckenbach.*

Source: *Mathematical Reviews*,

Vol. 8, No. 2

SHERMAN, D.

Sherman, D. On some problems of the theory of stationary oscillations. Bull. Acad. Sci. URSS. Sér. Math. [Izvestia Akad. Nauk SSSR] 9, 363-370 (1945). (Russian. English summary)

The method of the paper reviewed above is extended to treat similarly the problem of determining solutions of the differential equation $\Delta u + \lambda^2 u = 0$, with boundary condition the same as in the former problem. . . . E. F. Beckenbach.

Source: Mathematical Reviews,

Vol 8, No. 2

SHERMAN, D. I.

500

Sherman, D. I. Certain problems of the theory of potential. *Appl. Math. Mech.* [Akad. Nauk SSSR. Prikl. Mat. Mech.] 9, 479-488 (1945). (Russian. English summary) [AIF 15441]

The author shows how the potential function $u(x, y)$ which is regular in the finite simply-connected region S and which satisfies the boundary condition $a(s)u_x + b(s)u_y = f(s)$ on the continuously curved boundary L of S can be found as the solution of an equivalent Fredholm equation. Assuming $a^2(s) + b^2(s) = 1$ all along L , two cases arise according to whether in the representation $a = \cos \omega(s)$, $b = \sin \omega(s)$ the function $\omega(s)$ increases by a nonpositive or by a positive multiple of 2π when the contour L is described once in the positive sense. In the first case the boundary-value problem always has a solution, in the second case there are non-trivial solutions of the homogeneous problem.

By the same method Poincaré's problem, where the boundary condition is $du/dn + p(s)du/ds + q(s)u = f(s)$, is reduced to a Fredholm equation, and so are certain singular integral equations.
M. Golomb (Lafayette, Ind.)

Source: *Mathematical Reviews*, 1948, Vol 9, No. 3

22

SHENIN, S. I.

Institute of Mechanics, Academy of Sciences, USSR. "Concerning One Case of Variation of an Elastic Half-Space." Iz. Ak. Nauk SSSR, Otdel. Tekh. Nauk, No. 10-11, 1945. Submitted 16 Jul 1945.

Report U-1582, 6 Dec 1951.

"A description of electric waves," *Ann. Real Lab. Sci., Ser. 4, Vol. 1, No. 1*,
1917, p. 1.

A description of the calculation of the vector components of a
plane wave in a medium in which the medium is anisotropic.

SHERMAN, D. I.

★Schermann, D. I. Oscillation du demi-espace élastique aux déplacements ou aux forces extérieures données à la frontière. Acad. Sci. URSS. Publ. [Trudy] Inst. Seismolog. no. 118, 47 pp. (1946). (Russian)

This paper contains a new method of solution of the three-dimensional problem of propagation of vibrations in an elastic half-space when either the displacements or external forces are specified on the boundary. For simplicity, initial displacements and velocities are assumed to vanish. The method of solution is based on a direct application of Fourier integrals. After a sequence of transformations on scalar and vector potentials both boundary value problems are reduced to a consideration of certain quadruple integrals, which yield a qualitative analysis of the phenomenon of wave propagation. I. S. Sokolnikoff (Los Angeles, Calif.).

Source: Mathematical Reviews,

Vol 8 No.6

SHERMAN, E. I.

1A 1504

USSR/Mathematics
Potential theory

Feb 1946

"The General Problem of the Potential Theorem,"
D. I. Sherman, 14 pp

"Izv Ak Nauk Ser Mat" Vol X, No 2

Determination of a function $u(x,y)$ harmonic in a
finite (simply or multiply connected) domain S in
the plane $z = x + iy$, satisfying certain conditions
on the boundary L of S .

13T94

SHERMAN, D. I.

Sherman, D. I. *Steady elastic oscillations in the case of given displacements on the boundary of the medium.* Appl. Math. Mech. [Akad. Nauk SSSR, Prikl. Mat. Mech.] 10, 617-622 (1946). (Russian. English summary)

The elastic medium fills a finite multiply-connected domain S in the complex plane bounded by nonintersecting and nonenclosing closed curves. Consideration of the oscillations arising from given displacements on the boundary gives rise to a pair of second order partial differential equations in the longitudinal and transverse potentials subject to prescribed boundary conditions involving first partial derivatives of the potentials. Potentials of a form leading to the solution of a system of Fredholm integral equations with kernels containing the frequency λ of oscillations as a parameter are sought. The existence of solutions for $\lambda=0$ is established, and hence, by a theorem of Tamarkin [Ann. of Math. (2) 28, 127-152 (1927)], solutions for almost all λ exist. T. C. Doyle (Hanover, N. H.).

Ann. of Math. (2) 28, 127-152 (1927)

Vol. 28, No. 6

SHERMAN, D. I.

FA 1577

USSR/Oscillations - Theory
Mathematics, Applied

Feb 1947

"The Dirichlet and Neuman Problems in the Theory of
Steady Oscillations," D. I. Sherman, 8 pp

"Prikl Mat i Mekh" Vol XI, No 2

Reduction of the problems of the multi-connected
domain to the Fredholm equations, which differ
somewhat from the hitherto known integral equations
for the same problems and make possible direct
establishment of the existence of the solution.

1577

SHERMAN, D. I.

200

Scherman, D. J. Sur une méthode de résoudre certains problèmes de la théorie de l'élasticité pour les domaines doublement connexes. C. R. (Doklady) Acad. Sci. URSS (N.S.) 55, 697-700 (1947).

Let the elastic medium occupy a doubly connected domain S in the plane $z=x+iy$. The boundary L of S consists of two simple closed curves L_1 and L_2 having no points in common. Suppose L_1 is in the interior of L_2 and denote by S_1 and S_2 two simply connected domains bounded by L_1 and L_2 , respectively. The domain S_1 is then infinite. The determination of stresses throughout the domain S , when forces are prescribed on L , can be reduced to the search for two functions $\varphi(z)$ and $\psi(z)$, regular in S , and satisfying on L the condition $\varphi(t) + \overline{t\varphi'(t)} + \psi(t) = f(t)$, where t is a variable point on L and $f(t)$ is a known function. The author indicates a method of reducing the solution of this problem to an analogous problem for a simply connected domain and is led to one integral equation of Fredholm type for a certain auxiliary function defined over one of the curves L_1 or L_2 . I. S. Sokolnikoff (Los Angeles, Calif.).

Source: Mathematical Reviews, 1948, Vol 9, No. 2 (SMW) [Signature]

SHERMAN, D. I.

PA 58T92

USSR/Physics
Vibration
Mathematics, Applied

May 1947

"Several Particular Cases of a General Problem in the Theory of Vibrations," D. I. Sherman, Inst Mechanics, Acad Sci USSR, 4 pp

"Dok Akad Nauk SSSR, Nova Ser" Vol LVI, No 6

Contains number of mathematical formulas designed to prove that for any function $u(x, y; \lambda)$ satisfying the equation $\Delta u - \lambda^2 u = 0$,

$$u(x, y; \lambda) = \text{Re} \left[\frac{1}{\pi i} \int_{\gamma} v(s, \lambda) \frac{ds}{s} + \sum_{k=0}^{m+n} c_k(\lambda) i^k e^{-i k \theta} I_k(\lambda \rho) \right]$$

58T92

SHERMAN, D.I.

Serman, D. I. On the state of stress in some shrink-fitted members. *Izvestiya Akad. Nauk SSSR. Otd. Tehn. Nauk* 1948, 1371-1388 (1948). (Russian)

The paper contains a solution of the following two-dimensional elastic problem. A long hollow prismatic body whose section by a plane normal to the axis of the prism is a square (with rounded corners) with a circular hole at the center of the square, is shrink-fitted on a solid circular shaft. The elastic properties of the shaft are identical with those of the prism, and the lateral surface of the prism is free of stress. What is the state of stress in the member so formed? If the boundary of the square in the (x, y) -plane is L , and that of the circular hole is γ , the solution of the problem, following Muschelišvili, reduces to the search for four functions $\varphi_j(z)$ and $\psi_j(z)$ ($j=1, 2$) of a complex variable $z=x+iy$, analytic in the regions S_j , where S_1 is a doubly-connected region bounded by L and γ and S_2 is the circular region bounded by γ .

Source: *Mathematical Reviews*,

Vol 10. No. 0

The functions φ_i and ψ_i are determined by the following boundary conditions:

$$\varphi_1(t) + t\overline{\varphi_1'(t)} + \overline{\psi_1(t)} = 0$$

on L ;

$$\begin{aligned} \varphi_1(t) + t\overline{\varphi_1'(t)} + \overline{\psi_1(t)} &= \varphi_2(t) + t\overline{\varphi_2'(t)} + \overline{\psi_2(t)}, \\ \kappa\varphi_1(t) - t\overline{\varphi_1'(t)} - \overline{\psi_1(t)} &= \kappa\varphi_2(t) - t\overline{\varphi_2'(t)} - \overline{\psi_2(t)} + 2kt \end{aligned}$$

on γ , where κ and k are constants determined by the elastic properties of the medium and by the amount of shrink along γ . The author reduces the problem (by means of analytic continuation) to the determination of only two functions $\varphi(z)$ and $\psi(z)$, analytic in the region $S_1 + S_2$, of the form

$$\varphi(z) = \frac{1}{2\pi i} \int_L \frac{\omega(t)}{t-z} dt; \quad \psi(z) = \frac{1}{2\pi i} \int_L \frac{\omega(t) - t\omega'(t)}{t-z} dt,$$

where $\omega(t)$ satisfies a certain integral-differential equation and bars denote conjugate values. The solution of the latter equation is obtained in the form of an infinite series. A calculation of the distribution of normal stress along γ , contained in the paper, illustrates the practical value of the function-theoretic methods of solution of elastic problems.

I. S. Sokolnikoff (Los Angeles, Calif.).

SHERMAN, D. I.

76T45

USSR/Engineering
Wing Theory
Mathematics, Applied

May 1948

"The Prandtl Equation in the Theory of a Wing of
Finite Span," D. I. Sherman, Inst of Mech, Acad Sci
USSR, 6 pp

"Iz Ak Nauk SSSR, Otdel Tekh Nauk" No 5

Presents solutions for Prandtl's integral differential
singular equation for case when function $p(x)$ is
rational. Also presents approximate method for
solving the equation which will hold true for any
value of the function $b(x)$. Submitted 2 Feb 1948.

76T45

MA 11/17/120

USSR/Engineering
Stresses

Sep 48

"The Tension States in Some Pressed Parts," D. I. Sherman, Mech Inst, Acad Sci USSR, 18 pp

"Iz Ak Nauk SSSR, Otdel Tekh Nauk" No 9

A hollow prismatic body whose section is a plane perpendicular to its axis, is of quadrate shape. It is weakened by a symmetrically located circular hole and by mounting on a solid circular shaft. Tightness of fit is given. Elastic properties of both bodies are considered identical. Lateral surface of the prismatic body is assumed to be free

14/49123

USSR/Engineering (Contd)

Sep 48

from external forces. Calculates stress distribution in body. Concludes by considering case, when elastic properties of parts are not identical. Submitted 8 Mar 48.

SHERMAN, D. I.

14/49123

SHERMAN, D. I.

Serman, D. I. On methods of solving certain singular integral equations. Akad. Nauk SSSR. Prikl. Mat. Meh. 12, 423-452 (1948). (Russian)

This is a study of systems of the form

$$(1) \sum_{j=1}^2 \left\{ a_{kj}(t_0) \omega_j(t_0) + b_{kj}(t_0) (\pi i)^{-1} \int_L (t-t_0)^{-1} \omega_j(t) dt \right\} = f_k(t_0)$$

(k=1, 2), where L is a simple, closed, "smooth" curve (in the complex plane of s=x+iy), bounding a finite simply-connected region S; the ω_j(t) are the unknowns and the a_{kj}, b_{kj}, f_k are assigned, suitably differentiable on L; the integrals are in the sense of principal values. On letting c_{kj}=a_{kj}-b_{kj}, d_{kj}=a_{kj}+b_{kj}, one forms determinants Δ₁=|(c_{kj})|, Δ₂=|(d_{kj})|. The extensive literature relating to equations of type (1), and of other similar types, is largely concerned with transformations into regular Fredholm equations of the second kind, when Δ₁, Δ₂ (or other analogous functions) are distinct from zero on L. One of the novel features of this work is that one of the determinants is allowed to vanish at some points of L (the other one is assumed not 0 on L).

Specifically, it is assumed that Δ₁(t) has a simple zero at t=α (on L) and the coefficients are analytic at t=α (the latter condition can be lightened). It is shown that a reduction to regular Fredholm equations is possible and that (1) has a solution ω_j (j=1, 2) continuous on L. Such results are extended to systems

$$(2) \sum_{j=1}^2 \left\{ a_{kj}(t_0) \omega_j(t_0) + (\pi i)^{-1} b_{kj}(t_0) \int_L (t-t_0)^{-1} \omega_j(t) dt + T_j^k \right\} = f_k(t_0)$$

(k=1, 2), where T_j^k=∫_L ω_j(t) K_{kj}(t, t₀) dt; here the K_{kj}(t, t₀) (and the coefficients) are Hölder on L, in t, in t₀, and are analytic at the point t₀=α, at which Δ₁ has a zero of multiplicity m. The system (1) is also studied when the a_{kj}, b_{kj} are constants and L is an open arc. W. J. Trjitzinsky.

Source: Mathematical Reviews;

Vol 10 No. 3

SHERMAN, D. I.

Sherman, D. I. On a case in the theory of singular equations. Doklady Akad. Nauk SSSR (N.S.) 59, 647-650 (1948). (Russian)

The author studies the equation

$$(1) \quad A(t_0)u(t_0) + (\pi i)^{-1}B(t_0) \int (t-t_0)^{-1}w(t)dt + \int w(t)K(t, t_0)dt = f(t_0),$$

where integration is along a simple "smooth" contour L ; t, t_0 are on L ; the coefficients are essentially of a Hölder class on L . In previous literature (1) has been transformed into a regular Fredholm equation of the second kind, predominantly under the supposition that (2) $A^2 - B^2 \neq 0$ on L . The author gives a new method for effecting such a transformation when (2) does not hold. The case actually carried out is the one when $A - B$ has just one simple zero on L . The question of equivalence of the resulting Fredholm equation and of (1) is examined. The results obtained can be extended to systems of equations analogous to (1).

W. J. Trjitzinsky (Urbana, Ill.)

Smw

Source: Mathematical Reviews,

Vol. 9 No. 3

SHERMAN, D. I.

FA 55/49T83

USSR/Physics
Beams - Stress Analysis
Theory of Elasticity
Dec 48

"One Torsion Problem," D. I. Sherman, Inst of
Mech, Acad Sci USSR, 4 pp

"Dok Ak Nauk SSSR" Vol IXIII, No 5

Gives a method to solve special problems in the
theory of elasticity and hydrodynamics relating
to torsion and curvature in hollow prism-shaped
beams, the cross sections of which are areas with
double connections, problems in the theory of
elasticity for similar areas, and other problems

55/49T83

USSR/Physics (Contd)
Dec 48

Including the more general case where an ellipse
is placed asymmetrically with regard to a circle.
Submitted by Acad L. S. Leybenzon 14 Oct 48.

55/49T83

SHERMAN, D. I.

PA 149T42

USSR/Engineering - Mechanics
Elasticity

Sep/Oct 49

"Theory of Steady Vibrations of a Medium for
Given External Forces on Its Boundary," D. I.
Sherman, Moscow Inst of Mech, Acad Sci USSR, 4 pp

"Prikl Mat i Mekh" Vol XIII, No 5

Discusses steady vibrations of an elastic medium
filling a finite simply connected region lying
in the complex plane when effective external
forces are acting upon the curve bounding the
region. Submitted 11 May 46.

149T42
149T42

SHERMAN, D. I.

13900

Šerman, D. I., and Narodeckii, M. Z. On the torsion of some prismatic hollow bodies. Akad. Nauk SSSR. Inženeryi Sbornik 6, 17-46 (1950). (Russian)

The authors consider two prismatic hollow bars with the following cross-sections: 1) an ellipse with a concentric circular hole, and 2) a square with rounded edges with a concentric circular hole. In both cases the bars are twisted by a moment M acting on the outside contour. The problems are solved by conformal mapping, which, due to the fact that the cross-section is a doubly connected region, presents considerable difficulty. The method employed was developed by Šerman [Doklady Akad. Nauk SSSR 63, 499-502 (1948); these Rev. 10, 651]. The obtained stress function, and hence the expressions for the stresses are in a form of complicated series. To make them useful for practical applications the authors tabulate for both cases up to twelve coefficients of the series for several dimension ratios.

T. Leser.

Source: Mathematical Reviews,

Vol 13 No. 9

SMW

SHERMAN, D. I.

158T97

USSR/Physics - Mechanics
Elasticity

Mar/Apr 50

"Problem of Conformal Reflection," M. Z. Narodetskiy, D. I. Sherman, Moscow, 6 pp

"Priklad Matemat i Mekh" Vol XIV, No 2

Gives approximate, but sufficiently effective, solution of problem of conformal reflection, in a doubly connected region S in the complex z-plane against a circular ring. Submitted 31 Dec 49. .

158T97

SHERMAN, D. I.

Serman, D. I. On the stresses in a twisted circular beam weakened by a prismatic cavity. *Izvestiya Akad. Nauk SSSR Otd. Tehn. Nauk* 1951, 969-995 (1951). (Russian) Saint Venant's torsion problem for a long circular beam weakened by a rectangular prismatic cavity with rounded corners is solved by the introduction of an auxiliary function which assumes on the circular boundary the same values as the complex torsion function. The auxiliary function is

shown to satisfy a Fredholm integral equation whose kernel can be replaced, with a known degree of approximation, by a degenerate kernel. Thus the determination of the auxiliary function is reduced to the solution of a system of linear algebraic equations. Once the auxiliary function is known, the torsion function can be computed. Although the convergence of the approximating process is not fully established in the paper, extensive numerical computations testify to the remarkable effectiveness of the proposed method, even when it is applied to a thin-walled section.

I. S. Sokolnikoff (Los Angeles, Calif.).

Source: *Mathematical Reviews*,

Vol 13 No. 3

Serman

Mathematical Reviews
Vol. 15 No. 4
Apr. 1954
Mechanics

✓
Serman, D. I. Torsion of an elliptic cylinder stiffened with
a circular rod. Akad. Nauk SSSR. Inzhenernyi Sbornik
10, 81-108 (1951). (Russian)

A detailed solution of Saint-Venant's torsion problem is
given for a homogeneous and isotropic elliptical beam rein-
forced by a circular rod. The rod is welded onto the cylinder
along the lateral surface, and the axes of the rod and beam
coincide. This is a generalization of the corresponding tor-
sion problem for an elliptical beam weakened by a cylindrical
cavity, solved by D. I. Serman and M. Z. Nafodeckii
[same Sbornik 6, 17-46 (1950); these Rev. 13, 886].

I. S. Sokolnikoff (Los Angeles, Calif.)

② Serman 1951

8-25-54
JPP

SHERMAN, D.I.

Sherman, D. I. On a case of regularization of singular equations. Akad. Nauk SSSR. Prikl. Mat. Meh. 15, 75-82 (1951). (Russian)

In an earlier note [Doklady Akad. Nauk SSSR (N.S.) 59, 647-650 (1948); these Rev. 9, 442] the author indicated a new method of transforming the singular equation (in the sense of principal values)

$$(1) \quad A(t_0)w(t) + \frac{1}{\pi i} B(t_0) \int_L \frac{w(t)}{t-t_0} dt + \int_L w(t)G(t, t_0) dt = f(t_0)$$

into a Fredholm integral equation. Here L is a closed, suitably smooth curve, bounding a bounded, simply connected domain; t, t_0 are points on L ; A, B, G, f are given on L and are of Hölder classes. This transformation was possible when the functions $A \pm B$ vanish at some points of L . In this earlier work it was assumed that $A(t_0), B(t_0), G(t, t_0), f(t_0)$ are analytic in t_0 at those points. In the present work the author studies (1), adapting the method of his preceding note to the regularisation of (1) under less restrictive conditions on the coefficients involved. *W. J. Trjitzinsky.*

Source: Mathematical Reviews,

Vol. 12 No.

USSR/Physics - Stresses in Plates

May/June 51

"Stresses in a Ponderable Half-Plane Weakened by Two Circular Apertures," D. I. Sherman, Moscow, Inst Mech, Acad Sci USSR

"Prik Matemat i Mekh" Vol XV, No 3, pp 297-316

Considers elastic isotropic and homogeneous half-plane possessing 2 openings circular in form which are sufficiently far removed from the margin. Cf. G. V. Kolosov's "Application of Complex Variables to the Theory of Elasticity," 1935, Moscow, and N. I. Muskhelishvili's "Certain Basic Problems in the Mathematical Theory of Elasticity," 1949, Moscow. \Especially interest is in the stress near boundary of apertures. Submitted 16 Mar 51.

1857104

Mathematical Reviews
Vol. 15 No. 4
Apr. 1954
Mechanics

✓
Šerman, D. I. On stresses in a plane heavy medium with two identical symmetrically placed circular openings. Akad. Nauk SSSR. Prikl. Mat. Meh. 15, 751-761 (1951). (Russian)

A homogeneous and isotropic elastic material fills the semi-infinite triply-connected domain S , bounded by the straight line L_0 parallel to the X -axis, and by two non-intersecting circles L_1 and L_2 with equal radii R . The centers of the circles lie on the X -axis at a distance f from L_0 . The material filling S is acted on by a uniform gravitational force in the direction of the Y -axis, and the boundaries L_0, L_1, L_2 are free of external loads. A solution of this two-dimensional elastostatic problem, in the neighborhood of L_1 and L_2 , is obtained under the hypothesis that $R \ll f$. The author utilizes the Muskhelišvili formulations of such problems.

I. S. Sokolnikoff (Los Angeles, Calif.)

СЕРИЯ, I. I.

Mining Engineering

State of pressure in pillars between chambers. Resilient, ponderous mass, weakened by two elliptical holes. Part. 1. Izv. AN SSSR Otd. Tekh. nauk no. 6, 1952.

9. Monthly List of Russian Accessions, Library of Congress, November 195~~8~~₂, Uncl.

3

USSR .

2306 Sherman, D. L. Transverse bending of an elliptic beam weakened by a longitudinal cylindrical hole (in Russian), *Inzh. Sbornik, Akad. Nauk SSSR* 17, 121-150, 1953.

Analysis of cantilever beam of elliptical cross section, with longitudinal circular-cylindrical hole concentric with ellipse, loaded at free end by transverse force in direction of major axis of ellipse. Numerical results are presented in somewhat implicit form for a limited variety of ratios of cross-sectional dimensions, and some of them are compared with elementary solution. Reading of this 30-page paper, as with many other recent Russian articles, is seriously hampered by presence of only one single figure, bibliography consisting of a single item, absence of a list of symbols, and general lack of organization of material.

G. Winter, USA

SHERMAN, D. I.

Mathematical Reviews
Vol. 15 No. 3
March 1954
Analysis

Serman, D. I. On properties of infinite systems of equations in problems of torsion of certain doubly connected profiles. Akad. Nauk SSSR. Prikl. Mat. Meh. 17, 470-476 (1953). (Russian)

The Saint-Venant torsion problems for a circular cylinder weakened by a symmetrically located longitudinal circular cavity or by two longitudinal circular cavities have been reduced by the author [Doklady Akad. Nauk SSSR (N.S.) 63, 499-502 (1948); these Rev. 10, 651] and by R. D. Stepanov and Serman [Akad. Nauk SSSR. Inzhenernyi Sbornik II, 127-150 (1952); these Rev. 14, 430] to the solution of certain Fredholm integral equations. The solution of the integral equations was made to depend on the solution of two systems of linear algebraic equations in infinitely many unknowns. This note gives a demonstration that these systems are completely regular and hence admit of an estimate of the error resulting from truncating the systems.

I. S. Sokolnikoff (Los Angeles, Calif.).

LL

EH
7-21-54

SHERMAN, D. I.

3

Sherman, D. I. On a connection of the fundamental problem of the theory of elasticity with a singular case of a problem of Poincaré. Akad. Nauk SSSR. Prikl. Mat. Meh. 17, 685-692 (1953). (Russian)

The determination of two harmonic functions φ and ψ in a finite multiply connected domain from the boundary conditions

$$\frac{\partial \varphi}{\partial x} + \frac{\partial \psi}{\partial y} = f_1(s), \quad \frac{\partial \varphi}{\partial y} - \frac{\partial \psi}{\partial x} = f_2(s),$$

where the $f(s)$ are prescribed functions of the arc-parameter s on the boundary C of the domain, is reduced to the solution of a Fredholm integral equation. The boundary C consists of the exterior simple closed contour C_0 and several interior simple closed contours C_i not having common points, and the $f(s)$ are assumed to satisfy Hölder's condition on C .

I. S. Sokolnikoff (Los Angeles, Calif.)

SHERMAN, D. I.

Šerman, D. I. On a singular problem from potential theory. Doklady Akad. Nauk SSSR (N.S.) 94, 25-28 (1954). (Russian)

62

Let S be a bounded simply connected domain in the plane of $z = x + iy$, whose frontier L is a suitably smooth closed contour L ; the origin is taken in S . One is to find u, v , harmonic in S , continuous with their first partials on $S + L$, such that on L

$$(1) \quad u_x - v_y + a_{11}u + a_{12}v = f_1, \quad u_y + v_x + a_{21}u + a_{22}v = f_2,$$

where the coefficients are assigned, suitably differentiable functions of arc s ; $0 = a_{11} + a_{22} = a_{21} - a_{12}$ at no point of L . Here u and v are sought in terms of certain potentials, whose real densities satisfy a Fredholm system. The author leaves open the question as to whether two harmonic functions are expressible in the form considered, as well as the question as to the conditions under which problem (1) can be solved.

W. J. Trjitzinsky (Urbana, Ill.).

The rotation of a circular cylinder, surrounded with elliptic rod. "Inzhineriyy
Sbornik" by Academy of Science of the USSR, Department of Technical
Science, Institute of Mechanics. 1955.

SHERMAN, D. I. (Moscow)

Torsion of a circular cylinder reinforced by an elliptical rod.
Inzh.sbor. no.21:79-96 '55. (MIRA 8:11)

1. Institut mekhaniki Akademii Nauk SSSR
(Torsion)

SHERMAN, D. I.
USSR/ Engineering - Theory of elasticity

Card 1/1 Pub. 22 - 10/52

Authors : Sherman, D. I.

Title : ~~On the bending of a circular plate partly rigidly fixed and partly supported along its contour~~
On the bending of a circular plate partly rigidly fixed and partly supported along its contour

Periodical : Dok. AN SSSR 101/4, 623-626, Apr 1, 1955

Abstract : A theoretical analysis is presented of the physical deformations (bending) sustained by a circular plate under a uniformly distributed load, when the plate is partly rigidly fixed and partly supported along its contour. Three USSR references (1936-1952).

Institution : Acad. of Sc., USSR, Institute of Mechanics

Presented by: Academician N. I. Muskhelishvili, January 5, 1955

SHERMAN, D.I.

Bending of a circular plate partially supported and partially free
along its circumference. Dokl.AN SSSR 105 no.6:1180-1183 D '55.
(MLRA 9:4)

I.Institut mekhaniki Akademii nauk SSSR. Predstavleno akademikom
N.I.Muskhelishvili.
(Elastic plates and shells)

SOV/124-57-5-5859

Translation from: Referativnyy zhurnal. Mekhanika, 1957, Nr 5, p 120 (USSR)

AUTHOR: Sherman, D. I.

TITLE: Effective Integral-equation Methods Applied to Some Elasticity Problems (Effektivnyye metody integral'nykh uravneniy v primeneni k nekotorym zadacham teorii uprugosti)

PERIODICAL: Tr. 3-go Vses. Matem. s"yezda. Vol I. Moscow, AN SSSR, 1956, p 216

ABSTRACT: Bibliographic entry

Card 1/1

SHERMAN, D.I. (Moskva)

Solving some problems in torsion, bending and plane theory of elasticity for multiconnected regions [in Ukrainian with summaries in Russian and English]. Prykl. mekh. 3 no.4:363-377 '57.
(MIRA 11:2)

1. Institut mekhaniki AN SRSR.
(Strains and stresses)
(Elastic solids)

AUTHOR: Sherman, D. I.

20-114-4-15/63

TITLE: On a Problem in the Theory of Elasticity With Mixed Homogeneous Conditions (Ob odnoy zadache teorii uprugosti so smeshannymi odnorodnymi usloviyami)

PERIODICAL: Doklady Akademii Nauk SSSR, 1957, Vol. 114, Nr 4, pp. 733-736 (USSR)

ABSTRACT: An elastic, isotropic and homogeneous medium may satisfy a finite and simply connected domain which is located in the plane of the complex variables $z = x + iy$. Further, S is assumed to be surrounded by a sufficiently smooth closed contour L . The coordinate source is assumed to be contained in the domain S . On the boundary L the normal components v_n of the displacement vector and the tangential components of the stress vector T may be assumed. The components of the stress tensor and the displacement vector occurring in the medium are determined. For the purpose of simplification the author confines himself to a simply connected domain S . This problem is here reduced to a new and much simpler system of Fredholm equations and the nuclei are expressed immediately by elementary functions. Such systems of integral equations

Card 1/3

On a Problem in the Theory of Elasticity With Mixed
Homogeneous Conditions

20 114-4-15/63

can be interpreted in a comparatively simple manner by means of modern computation methods. The assumed boundary values of the orders ν_n and T themselves are expressed in the known manner by the two functions $\varphi(z)$ and $\Psi(z)$ of the complex variables z . These two functions are regular in the domain S . The problem is reduced to their determination from two real boundary conditions. The boundary conditions, after some transformations, are explicitly given. After lengthy transformations a system of Fredholm's integral equations is obtained for the unknown densities. The nuclei occurring therein are steady functions of the arguments s and s_0 , where s denotes the length of the arc. In conclusion the author once more proves the solubility of this system of Fredholm's integral equations. This system of equations always has a unique solution. There are 4 references, 4 of which are Soviet.

ASSOCIATION: Institut mekhaniki Akademii nauk SSSR (Mechanical Institute of the AS USSR)

Card 2/3

On a Problem in the Theory of Elasticity With Mixed Homogeneous 20-114-4-15/63
Conditions

PRESENTED: December 26, 1956, by L. I. Sedov, Member, Academy of
Sciences, USSR

SUBMITTED: October 16, 1956

Card 3/3

SHERMAN, D.I. (Moskva)

Stressed state of a twisted square beam with a symmetrical
circular hollow. Prikl.mekh. 4 no.3:250-262 '58. (MIRA 13:8)

1. Institut mekhaniki AN SSSR.
(Girders)

16(1);10(2)

PHASE I BOOK EXPLOITATION

SOV/2659

Akademiya nauk SSSR. Institut mekhaniki

Inzhenernyy sbornik, t. 25 (Engineering Symposium, Vol. 25) Moscow, Izd-vo AN SSSR, 1959. 218 p. Errata slip inserted. 2,200 copies printed.

Ed.: A.A. Il'yushin; Ed. of Publishing House: D.M. Ioffe; Tech. Ed.: Ye. V. Makuni.

PURPOSE: This book is intended for applied mathematicians, physicists and engineers.

COVERAGE: The book is a collection of articles published by the Department of Engineering Sciences of the Institut mekhaniki (Institute of Mechanics) of the Academy of Sciences, USSR. The articles discuss various aspects of the mechanics of materials and of fluid mechanics, such as stress and bending of beams, shells, plates and reels, supersonic gas flows, vibrations, etc. The problems are treated in a highly theoretical, i.e., mathematical, manner. References are given at the end of each article.

TABLE OF CONTENTS:
Card 1/5

Engineering Symposium, Vol. 25

SOV/2659

Sherman, D.I. On the Problem of the Torsion of an Elliptical Beam, Longitudinally Weakened by an Elliptical Cavity	3
Galimov, K.Z. On Large Bendings of a Rectangular Cylindrical Shell	20
Aleksandrov, A.Ya., and N.I. Nazatov. Bending of a Tapered Cantilever Plate	37
Grigor'yev, A.S. On Plates of Equal Resistance to Bending	45
Sheremet'yev, M.P. Bending of an Infinite Plate Which Is Weakened by an Elliptical Hole, and the Edge of Which Is Reinforced by a Thin Ring	51
Alekseyev, S.A. Design of a Circular Elastic Membrane Under Uniform Lateral Load	64
Chuvikovskiy, V.S. Lateral Vibrations of Rods and Plates With Reactive Tensile Forces	81
Stepanov, R.D. On the Flutter of Cylindrical Shells Moving in a Gas	92
Card 2/5	

Engineering Symposium, Vol. 25	SOV/2659	
Livanov, K.K. Stability of a Fastened Shell in a Supersonic Flow		101
Kurdin, N.S. Bending of a Partially Loaded Rectangular Plate With Two Supported and Two Free Edges		104
Karmishin, A.V. A Method of Solving Systems of Five-Termed Algebraic Equations Pertaining to Certain Problems of Engineering Mechanics		111
Nagibin, L.N. On the Stresses of a Weighable Anisotropic Semiplane Weakened by Two Circular Holes		122
Tarabasov, N.D. Stressed State of a Multiply-Connected Semiplane Caused by the Pressing in of Discs		136
Golitsynskaya, Ye.D. Certain Problems of the Stability of Thin Cylindrical Shells		145

Card 3/5

Engineering Symposium, Vol. 25

SOV/2659

Lukash, P.A., and N.N. Leont'yev. Design of a Spherical Shell Supported by a Foundation	154
Vyaz'menskiy, S.P. On the Stability of a Rectilinear Form of the Equilibrium of an Elastic Compressed Twisted Rod	164
Lomakin, V.A. A One-dimensional Problem of the Temperature Stresses in a Plastic-Elastic Medium	174
Marenov, A.Ye. A Study of Heat-Exchange in Supersonic Air Flows in Pipes	179
Yur'yev, I.M. Approximate Solution of the Fundamental Boundary Value Problems of a Supersonic Gas Flow	188
Shekhtman, Yu.M. Flow of Liquid to a Vertical Fissure With Filler	197

Card 4/5

Engineering Symposium, Vol. 25

SOV/2659

Karpychev, V.A. On the Displacement of a Water-Oil Contact
in Formations With Bottom Water

208

AVAILABLE: Library of Congress

Card 5/5

LK/esp
12-29-59

24(6)

PHASE I BOOK EXPLOITATION

SOV/2250

Akademiya nauk SSSR. Institut fiziki zemli

Nekotoryye voprosy mekhaniki deformiruyemykh sred (Some Problems in the Mechanics of Deformable Media). Moscow, Izd-vo AN SSSR, 1959. 219 p. (Series: Its: Trudy, Nr. 2 /169/) Errata slip inserted. 2,000 copies printed.

Ed.: V.A. Magnitskiy, Doctor of Technical Sciences; Ed. of Publishing House: V.A. Kalinin; Tech. Ed.: Yu. V. Rylina.

PURPOSE: This book is intended for engineers and geophysicists concerned with problems of deformations.

COVERAGE: This collection consists of eight articles on the mechanics of deformations in solid plastic media as applied to the solution of geophysical and engineering problems. No personalities are mentioned. References appear at the end of each article.

TABLE OF CONTENTS:

Card 1/5

Some Problems (Cont.)

SOV/2250

Gurevich, G.I., and A.L. Rabinovich. Relation Between Stresses and Displacements in Large Deformations for the Case of a One-dimensional Problem 3

In the analytical study of geometrical and kinetic deformations of elastic and residual nature, which are of significance in attenuation and dispersion of seismic waves, the authors derive general equations of motion. 12

Gurevich, G.I. Relation Between Stresses and Displacements in Large Deformations for the General Case of a Three-dimensional Load 27

The author considers the application of Maxwell's equation to a case of a residually deformed solid-liquid body which can be considered as a "massive" one and to which the usual formulas of the theory of elasticity are applicable.

Gurevich, G.I. Generalized Maxwell Equation for Three Measurements Taking Into Consideration Small Elastic Aftereffect Deformations 60
In the study of rock behavior in cases of static and dynamic

Card 2/5

Some Problems (Cont.)

SOV/2250

loading, the usual Maxwell's equation is not adequate. Taking into account the additional components of deformation, a new equation embodying the relationship between shear deformation and the velocity of full shear deformation is analyzed.

Gurevich, G.I. Initial Considerations in the Approach to Tectonic Modeling 75

The author deals with considerations in the application of the principle of similitude to the modeling of tectonic and hydrodynamic processes in the solution of geodynamic problems. The following names are mentioned: B.L. Shneyerson, Ye. N. Lyustikh, A.A. Ilyushin, M.V. Gzovskiy.

Khaykovich, I.M. Propagation of Vibrations in a Medium With Relaxation of Stresses 145

The theory of propagation of seismic waves in an ideally elastic medium is not adequate for purposes of interpretation. The present article establishes the quantitative corrections for a half-space subjected to axially symmetric loading. Maxwell's three-dimensional equation is used in finding a solution for corrections. The following names are mentioned: G.I.

Card 3/5

Some Problems (Cont.)

SOV/2250

Petrachen', K.I. Ogurtsov.

Khaykovich, I.M. Beam Method of Computing the Wave Intensity in a Relaxing Medium With a Large Relaxation Time 179

The author refers to various scientists offering the solution of nonstationary problems in the theory of elasticity leading to the determination of the intensity and the force of reflected waves. He introduces a so-called beam method for computing the propagation of a wave in nonideal elastic media. The following names are mentioned: G.I. Petrashen', V.M. Babich, G.O. Gurevich.

Sherman, D.I. Problem of the Stressed Condition of a Semiplane Without External Load and With Two Sunken Circular Orifices 187

The article discusses the distribution of stresses caused by gravity in media weakened by holes or openings. The problem is of interest in analyzing the rock pressure in the neighborhood of shaft openings and for the study of seismic conditions.

Card 4/5

SOV/2250

Some Problems (Cont.)

Keylis-Borok, V.I., and V.I. Ul'yanova. Problem of Creep in Hollow
Cylinders Under Normal Pressure 211

The author considers the process of residual deformation in a hollow cylinder and takes into account the time changes of stresses and deformations. This problem is of interest in theoretical studies of seismic behavior and also in studies of the relationship between the creep and interior pressure in pipes. The following names are mentioned: A.F. Golovin, L.I. Kachanov, A.A. Abramov, L.G. Shershen', I.K. Snitko.

AVAILABLE: Library of Congress

IS/ec
10-15-59

Card 5/5

S/124/60/000/008/010/011
A005/A001

Translation from: Referativnyy zhurnal, Mekhanika, 1960, No. 8, p. 120, # 10587

AUTHOR: Sherman, D. I.

TITLE: On the Problem of the Stress State of a Bounded Semi-Infinite Plane With Two Circular Deep-Drawn Apertures

PERIODICAL: Tr. In-ta fiz. Zemli. AN SSSR, 1959, No. 2, (169), pp. 187-210

TEXT: The author investigates the stresses in a plane triply-connected region having the shape of a semi-infinite plane with two circular apertures and affected by the gravity; the straight line connecting the centers of the apertures is perpendicular to the gravity direction. Assuming that the radii of the apertures are small in comparison with the depth of their position, the given problem can be reduced, with a sufficient degree of accuracy, to the problem of stresses in an infinite plane weakened by two circular apertures, at the boundaries of which the following boundary conditions are fulfilled.

$$X_x \cos(n, x) + X_y \cos(n, y) = k_1 \cos(n, x)$$

$$X_y \cos(n, x) + Y_y \cos(n, y) = k_2 \cos(n, y) \quad (*)$$

Card 1/2

S/124/60/000/008/010/011
A005/A001

On the Problem of the Stress State of a Ponderable Semi-Infinite Plane With Two Circular Deep-Drawn Apertures

where k_1, k_2 are certain positive constants. The author solved this problem earlier (Prikl. matem. i mekhan., 1951, Vol. 15, No. 3, No. 6). In the article reviewed, a new simpler solution is given. Using the special formulation of the boundary conditions (*), the author reduces the problem by a particular transformation to the elementary auxiliary problem of the two-dimensional elasticity theory for a plane with one circular aperture and to the determination of a certain infinite sequence of constants, for which an infinite system of linear algebraic equations is obtained. The numerical solution of the system is presented for the case $R_1 = 2 R_2 = 0.5 r$, where R_1, R_2 are the radii of the apertures, r is the distance between their centers. The infinite system is shortened at first to 17 and then to 7 equations; both of finite systems obtained in this way are solved by iterations. The author does not present a detailed substantiation and states that the infinite system mentioned is quasi-regular for arbitrary values of R_1, R_2, r . The case of equal apertures is especially considered.

S. G. Mikhailin

Translator's note: This is the full translation of the original Russian abstract.

Card 2/2

SHERMAN, D.I. (Moskva)

Torsion of an elliptic bar longitudinally weakened by an elliptic cavity. Inzh. sbor. 25:3-19 '59.

(MIRA 13:2)

(Elastic rods and wires) (Torsion)

KEYLIS-BOROK, Vladimir Issakovich; SHERMAN, D.I., otv.red.; YANOVSKAYA,
T.B., red.izd-va; POLYAKOVA, T.V., tekhn.red.

[Interference surface waves] Interferentsionnye poverkhnostnye
volny. Moskva, Izd-vo Akad.nauk SSSR, 1960. 194 p.
(MIRA 13:?)

(Wave motion, Theory of)

PHASE I BOOK EXPLOITATION
 SOVI/4000
 SOVI/12-4-27

Al'manentye nauki SSSR. Institut matematiki
 Inzhenernyy sbornik, t. 27 (Engineering Collection, Vol. 27) Moscow, Izd-vo AN SSSR, 1980. 210 p. 2,000 copies printed.
 Sponsoring Agency: Akademiya nauk SSSR. Otdeleniye tekhnicheskikh nauk.
 Ref. Ed.: A. A. D'yakov; Ed.: V. M. Akhmedov; Ed. of Publishing House: V. M. Akhmedov; Tech. Ed.: A. P. Guseva.
 PURPOSE: This book is intended for engineers, applied physicists, and applied mathematicians.
 COVERAGE: The book consists of 24 articles on such problems as viscosity, superonic flow, theory of shells, stability, plasticity and elasticity, the bending of thin plates and shells, and various aspects of applied mathematics. No personalities are mentioned. References accompany most of the articles.

Pillnov, M. V. On the Problem of Displacing Gas by Water	54
Bobotin, V. V. Application of Statistical Methods for the Evaluation of the Strength of Structures Subjected to Stochastic Forces	58
Morozov, A. A. The Behavior of Complex Eigen Values in the Problem of Buckling Plates	70
Slobodkin, A. M. Stability of an Elastic Beam With Rigid Inclusions in Superionic Flow	77
Shenoy, P. N. S. Vibrations of an Elastic String	81
D'yakov, A. A. Elastoplastic Stability of Structures Containing Rod Elements	87
Pogoy, B. M. Stability of Circular Thin Plates Beyond the Elastic Limit	98
Zubchikov, Y. G. Stability of Structural Rods Beyond the Elastic Limit	101
Tsarkov, I. S. On the Bending of a Closed Cylindrical Shell by a Concentrated Force	114
Shapoval, D. I. Strains in a Physical Medium Weakened by Elliptical and Circular Holes	124
Ugolechikov, A. G. Determination of Stresses Caused by Pressing Several Circular Washers Into a Plate With Variable Interference	137
Mishonov, M. On the Practical Calculation of Bending Moments of Shells Supported by a Rectangular Frame	163
Pheylshteyn, G. I. Statistical Calculation of Latticed Cylindrical Slotted Shells	171
Volkov, A. N. "Contact Method" in the Coupling of a Cylindrical Shell of Open and Closed Profile	179
Burakovskiy, Ye. F. Symmetric Bending of Kirchhoffian and Non-synchronous Orthotropic Shells of Revolution Taking Into Account Large Deflections and an Uneven Temperature Field	185
Kurkin, N. S. Lower Limit of a Kinematically Maximum Admissible Load	200
Levit, D. Ye. Homographic Solution of Equations of the Fourth and Fifth Degree	203
Svaynov, B. D. Application of the Method of Asymptotic Interpolation to the Solution of One Equation of the Natural Vibrations of Shells	207

AVIALENZ: Library of Congress

24.4200
12.8000

36978
S/044/62/000/003/018/092
C111/C222

AUTHOR: Sherman, B. I.

TITLE: On the stresses in a medium with weight which is weakened by an elliptical and a circular hole

PERIODICAL: Referativnyy zhurnal, Matematika, no. 3, 1962, 39, abstract 3B168. ("Inzhenernyy sb.," 1960, 27, 124-156; 28, 151-170)

TEXT: The author examines the stress distribution in a half-plane with two holes sufficiently far away from the boundary; one hole is elliptical and the other circular. The centers of the holes lie on a straight line parallel to the boundary of the half-plane. The complex potentials are determined by a method of the author, which is frequently used to solve elasticity problems for multiply connected domains (cf., e. g., *Rzh. Mat.*, 1960, 11566). The solution is illustrated by a concrete example, in which the calculation scheme is given in great detail. The connection between this problem and that of shocks in mining is considered.

f

[Abstracter's note: Complete translation.]

Card 1/1

27799

S/508/60/028/000/012/022
D251/D305

24,4200 (1103,1327) 1:31

AUTHOR: Sherman, D.I. (Moscow)

TITLE: On the tension in a heavy medium weakened by
elliptical and circular holes. (Part II)PERIODICAL: Akademiya nauk SSSR. Otdeleniye tekhnicheskikh nauk.
Inzhenernyy sbornik, v. 28, 1960, 151 - 170TEXT: This article is a continuation of the author's previous
work with the same title in Inzhenernyy sbornik, v. 27. The limit-
ing conditions are obtained in the form

$$\varphi^*(\tau) + \frac{\left\{ \sigma_1 + A\left(\tau + \frac{1}{\tau}\right) \right\}}{A\left(1 - \frac{\tau^2}{\rho^4}\right)} \overline{\varphi^{*'}(\tau)} + \overline{\psi^*(\tau)} = f^*(\tau) \quad (5.2)$$

where the independent term has the form

Card 1/9

UH

27799

S/508/60/028/000/012/022
D251/D305

On the tension in a heavy ...

$$f^*(\tau) = f_0^*(\tau) + A(2h_1 + 2h_2\rho^{-2})\tau + A(2h_1 + 2h_2\rho^2)\frac{1}{\tau} + C_1. \quad (5.3)$$

It follows easily that

$$\varphi(z) = A(2h_1 + 2h_2\rho^2)\frac{1}{z} + \sum_{k=0}^{\infty} \{\alpha_k^* R_k(\zeta) + \epsilon_k \beta_k^* W_k(\zeta)\}, \quad (5.4)$$

$$\begin{aligned} & \left\{ a_1 + A \left(\frac{\rho^2}{z} + \frac{\zeta}{\rho^2} \right) \right\} \varphi'(z) + \psi(z) = \\ & = A(2h_1\rho^2 + 2h_2)\frac{1}{z} + \sum_{k=0}^{\infty} \{\alpha_k^* \Omega_k(\zeta) + \epsilon_k \beta_k^* G_k(\zeta)\}. \end{aligned} \quad (5.3)$$

These formulae are true outside the contour L_1 and, in particular on L_2 , where

$$\varphi(t) = \sum_{n=0}^{\infty} \varphi_n^* \left(\frac{t-a_2}{R} \right)^n \quad (5.7)$$

Card 2/9

27799

S/508/60/028/000/012/022
D251/D305

On the tension in a heavy ...

with coefficients of the series

$$\varphi_n^* = A(2h_1 + 2h_2\rho^2)\theta_n + \sum_{k=0}^{\infty} \{\alpha_k^* r_{k,n}^* + \varepsilon_k \beta_k^* w_{k,n}^*\}. \quad (5.8)$$

Also, outside L_1

$$\begin{aligned} \bar{z}\varphi'(z) + \psi(z) = R \left\{ \frac{\bar{z}-a_2}{R} - \frac{A}{R} \left[\frac{a_1-a_2}{A} + \left(\frac{\rho^2}{\zeta} + \frac{\zeta}{\rho^2} \right) \right] \right\} \varphi'(z) + \\ + \sum_{k=0}^{\infty} \{\alpha_k^* \Omega_k(\zeta) + \varepsilon_k \beta_k^* G_k(\zeta)\} + A(2h_1\rho^2 + 2h_2) \frac{1}{\zeta}. \end{aligned} \quad (5.9)$$

and hence on L_2

$$-[\bar{t}\varphi'(t) + \psi(t)] = -\varphi_1^* \frac{R}{t-a_2} + \sum_{n=0}^{\infty} \psi_n^* \left(\frac{t-a_2}{R} \right)^n, \quad (5.10)$$

Card 3/9

LH

27799
S/508/60/028/000/012/022
D251/D305

On the tension in a heavy ...

On L_2 (1.7) is obtained in the form

$$\omega(t) = \sum_{n=0}^{\infty} \left\{ \varphi_n \left(\frac{t-a_2}{R} \right)^n + \psi_n \left(\frac{R}{t-a_2} \right)^n \right\} + \beta_0 \frac{t-a_2}{R} + \alpha_{-1} + h_2 R \frac{R}{t-a_2}, \quad (6.1)$$

By means of (1.12) α_k and β_k are expressed by an infinite system of equations, also given. In order to rationalize the calculation further, a transformation is carried out into the form of terms occurring in the formula of Kolosov of the components of stress

$$\begin{aligned} X_x + Y_y &= 4\text{Re}\varphi_1'(z), & (6.10) \\ Y_y - X_x + 2iX_y &= 2[\bar{z}\varphi_1''(z) + \psi_1'(z)]. \end{aligned}$$

The system

LK

Card 4/9

27799

S/508/60/028/000/012/022
D251/D305

On the tension in a heavy ...

$$\alpha_k + \sum_{m=0}^{\infty} \{ \alpha_m^* \rho_{m,k+1}^* + \varepsilon_m \beta_m^* q_{m,k+1}^* \} = -A(2h_1 c_{k+1} + 2h_2 d_{k+1}) \quad (6.5)$$

(k = 0, 1, 2, ...).

$$\beta_k - \sum_{m=0}^{\infty} \{ \alpha_m^* r_{m,k+1}^* + \varepsilon_m \beta_m^* \omega_{m,k+1}^* \} = A(2h_1 + 2h_2 c^2) \theta_{k+1}$$

(k = 1, 2, ...).

is then considered. This system is said to be quasi irregular always for any relative dimensions of the space. On the basis of Cauchy's formula for the function $\zeta^m(\zeta - \lambda_2)^{-m}$ regular inside a closed curve γ , and making use of (3.5) one obtains

$$h_{m,n} = \frac{1}{2\pi i} \int_{\gamma} \frac{\tau^m}{(\tau - \lambda_2)^m} \frac{d\tau}{(\tau - \lambda_1)^{m-n+1}} \quad (n = 1, 2, \dots, m) \quad (7.2)$$

UH

Card 5/9

27799
S/508/60/028/000/012/022
D251/D305

On the tension in a heavy ...

$$|h_{m,n}| \leq \frac{\rho^{m+1}}{(|\lambda_2| - \rho)^m (\rho - |\lambda_1|)^{m-n+1}} \quad (n = 1, 2, \dots, m). \quad (7.3)$$

near $\zeta = \lambda_1$. From the general form of the expansion within $\zeta = \lambda_2$ of a function regular outside γ , and making use of (3.5), one obtains

$$l_{m,n} = \frac{1}{2\pi i} \int_{\gamma} \frac{\tau^m}{(\tau - \lambda_1)^m} \cdot \frac{d\tau}{(\tau - \lambda_2)^{m-n+1}} \quad (n = 1, 2, \dots, m) \quad (7.5)$$

$$l_{m,n} \leq \frac{\rho^{m+1}}{(\rho - |\lambda_1|)^m (|\lambda_2| - \rho)^{m-n+1}} \quad (n = 1, 2, \dots, m). \quad (7.6)$$

The right-hand-sides of (7.3) and (7.6) give the relationship between $h_{m,n}^*$ and $l_{m,n}^*$. The functions

Card 6/9

27799

S/508/60/028/000/012/022
D251/D305

On the tension in a heavy ...

$$h_m(\zeta) = \left(\frac{R}{A}\right)^m \sum_{n=1}^m h_{m,n} \frac{1}{(\zeta - \lambda_1)^n}, \quad (7.7)$$

$$l_m(\zeta) = \left(\frac{R}{A}\right)^m \sum_{n=1}^m l_{m,n} \left(\frac{\zeta}{\rho^2 - \lambda_1 \zeta}\right)^n.$$

are important in the further working. The coefficients of the expansion of $(\lambda_1 - \zeta)^{-1}$ within $z = a_2$ are positive and the expansion has the form

$$\frac{1}{\lambda_1 - \zeta} = \frac{1}{|\lambda_1|} \frac{A}{R} \sum_{k=0}^{\infty} |0_{k+1}| \left(\frac{z - a_2}{R}\right)^k, \quad (7.8)$$

The dominated expansion on L_2 of $(\lambda_1 - \zeta)^{-n}$ for some integer $n > 0$ is of the form,

Card 7/9

27799

S/508/60/028/000/012/022
D251/D305

On the tension in a heavy ...

$$c_0^n \sum_{k=0}^{\infty} (-1)^k C_{-n}^k \theta^k \left(\frac{z-a_2}{R} \right)^k, \quad (7.9)$$

$$c_0 = \frac{A}{R} \frac{1}{|\lambda_1|} a_0 \theta.$$

The Taylor series

$$b_0 \frac{\rho}{c} \frac{1}{1-\rho^2} x^m \sum_{k=0}^{\infty} \Delta^k \left(\frac{z-a_2}{R} \right)^k, \quad \Delta = \max\{\sigma, \theta\}. \quad (7.12)$$

is hence constructed. The expansion of (7.12) is the dominating series of the expansion of the first function of (7.7). The second function of (7.7) (also regular outside γ) gives a nearly-linear expansion

$$(\lambda_2 \sigma_0 - \rho^2 = \lambda_2^2 - \rho^2 > 0)$$

$$\frac{\zeta}{\rho^2 - \lambda_2^2} = \frac{1}{\lambda_2} \frac{\rho^2}{\lambda_2} \frac{1}{(\lambda_2 \sigma_0 - \rho^2) + \lambda_2 \sum_{k=1}^{\infty} \sigma_k \left(\frac{z-a_2}{R} \right)^k}$$

W

Card 8/9

27799
 S/508/60/028/000/012/022
 D251/D305

On the tension in a heavy ...

Hence, by various simplifications, equations of the form

$$\frac{1}{\zeta+1} = \frac{1}{\frac{a_1-a_2}{A}-2} \frac{1+\frac{1}{\zeta}}{1-\frac{R}{a_1-a_2-2A} \frac{z-a_2}{R}}, \quad (7.21)$$

$$\frac{1}{\zeta-1} = \frac{1}{\frac{a_1-a_2}{A}+2} \frac{1-\frac{1}{\zeta}}{1-\frac{R}{a_1-a_2+2A} \frac{z-a_2}{R}},$$

are obtained. The article ends with a review of the theory of the method. There are 2 Soviet-bloc references.

SUBMITTED: May 12, 1959

LH

Card 9/9

LAVRENT'YEV, M.A., otv.red.; MIKHAYLOV, G.K., red.; BITSADZE, A.V.,
red.; VEKUA, I.N., red.; DZHANELIDZE, G.Yu., red.; LUR'YE, A.I.,
red.; MANDZHAVIDZE, G.P., red.; MIKHAYLOV, G.K., red.; SEDOV, L.I.,
red.; SOBOLEV, S.L., red.; SOKOLOVSKIY, V.V., red.; KHRISTIANOVICH,
S.A., red.; SHERMAN, D.L., red.; RYVKIN, A.Z., red.izd-va;
VOLKOVA, V.V., tekhn.red.

[Problems in the mechanics of solids] Problemy mekhaniki sploshnoi
sredy; k semidesiatiletiiu akademika N.I.Muskhelishvili. Moskva,
1961. 577 p. (MIRA 14:3)

1. Akademiya nauk SSSR. (Elastic solids)
(Mechanics, Analytic)

SHERMAN, D. I. (Moskva)

Ponderable medium weakened by periodical circular holes. Part 1.
Inzh.sbor. 31:24-75 '61. (MIRA 14:6)
(Elastic plates and shells)

PHASE I BOOK EXPLOITATION SOV/6201

29

Vsesoyuznyy s"yezd po teoreticheskoy i prikladnoy mekhanike. 1st, Moscow, 1960.

Trudy Vsesoyuznogo s"yezda po teoreticheskoy i prikladnoy mekhanike, 27 yanvarya -- 3 fevralya 1960 g. Obzornyye doklady (Transactions of the All-Union Congress on Theoretical and Applied Mechanics, 27 January to 3 February 1960. Summary Reports). Moscow, Izd-vo AN SSSR, 1962. 467 p. 3000 copies printed.

Sponsoring Agency: Akademiya nauk SSSR. Natsional'nyy komitet SSSR po teoreticheskoy i prikladnoy mekhanike.

Editorial Board: L. I. Sedov, Chairman; V. V. Sokolovskiy, Deputy Chairman; G. S. Shapiro, Scientific Secretary; G. Yu. Dzhanelidze, S. V. Kalinin, L. G. Loytsyanskiy, A. I. Lur'ye, G. K. Mikhaylov, G. I. Petrov, and V. V. Rumyantsev; Resp. Ed.: L. I. Sedov; Ed. of Publishing House: A. G. Chakhirev; Tech. Ed.: R. A. Zamarayeva.

Card 1/6

Transactions of the All-Union Congress (Cont.)

SOV/6201

(25)

PURPOSE: This book is intended for scientific and engineering personnel who are interested in recent work in theoretical and applied mechanics.

COVERAGE: The articles included in these transactions are arranged by general subject matter under the following heads: general and applied mechanics (5 papers), fluid mechanics (10 papers), and the mechanics of rigid bodies (8 papers). Besides the organizational personnel of the congress, no personalities are mentioned. Six of the papers in the present collection have no references; the remaining 17 contain approximately 1400 references in Russian, Ukrainian, English, German, Czechoslovak, Rumanian, French, Italian, and Dutch.

TABLE OF CONTENTS:

SECTION I. GENERAL AND APPLIED MECHANICS

• Artobolevskiy, I. I. Basic Problems of Modern Machine Dynamics	5
• Bogolyubov, N. N., and Yu. A. Mitropol'skiy. Analytic Methods of the Theory of Nonlinear Oscillations	25

Card 2/6

Transactions of the All-Union Congress (Cont.)	SOV/6201	
Kachanov, L. M. On Some Variational Principles and Methods in the Theory of Plasticity		358
Kupradze, V. D. The Singular Integral Equation Method in the Spatial Theory of Elasticity		374
Rabotnov, Yu. N. Creep		384
Florin, V. A. Present State and Future Problems in the Mechanics of Soils		396
Sherman, D. I. Two- and Three-Dimensional Problems in the Static Theory of Elasticity		405

AVAILABLE: Library of Congress

SUBJECT: Physics

Card 6/6

IS/dmp/mas
2-13-62

SHERMAN, D.I.

Reducing certain problems in the theory of steady-state vibrations to Fredholm's integral equation. Izv. AN Arm. SSR. Ser. fiz.-mat. nauk 16 no.4:41-63 '63. (MIRA 16:8)

1. Institut mekhaniki AN SSSR.

ARAMANOVICH, I.G.; SHERMAN, D.I. (Moscow)

"On certain special problems in elasticity".

report presented at the 2nd All-Union Congress on Theoretical and Applied Mechanics, Moscow, 29 January - 5 February 1964

L 32971-66 EWP(k)/EWT(d)/EWT(m)/I-2/EWP(w) IJP(c) EM
ACC NR: AT6016915 (r) SOURCE CODE: UR/0000/65/000/000/0352/0399

37
B+1

AUTHOR: Sherman, D. I.

ORG: Institute of Mechanics, AN SSSR (Institut mekhaniki AN SSSR)

TITLE: A solution of the Dirichlet problem for a circular ring and some applications in potential theory and elasticity theory

SOURCE: International Symposium on Applications of the Theory of Functions in Continuum Mechanics. Tiflis, 1963. Prilozheniya teorii funktsiy v mekhanike sploshnoy sredy. t. 1: Mekhanika tverdogo tela (Applications of the theory of functions in continuum mechanics. v. 1: Mechanics of solids); trudy simpoziuma. Moscow. Izd-vo Nauka, 1965, 352-399

TOPIC TAGS: Dirichlet problem, boundary value problem, elasticity theory, approximate solution

ABSTRACT: The Dirichlet problem is solved for a ring of concentric circles L_1 and L_2 . The desired function is regular in the ring and is defined by the boundary conditions:

$$\varphi(t) + \overline{\varphi(t)} = f_j(t) + 2C_j$$
$$L_j \quad (j = 1, 2), \quad C_1 = 0,$$

where $f_j(t)$ are Holder constants and C_2 is some unknown constant.

L 32971-66

ACC NR: AT6016915

The solution is given in the form

$$\varphi(z) = \sum_{n=0}^{\infty} \left[G(\lambda^n z) + \Omega\left(\frac{z}{\lambda^n}\right) \right] + \frac{1}{4\pi i} \int_{L_1} \frac{f_2(t)}{t} dt.$$

which has the advantage over other solutions (in particular those in power series) that solutions may be determined to any desired degree of accuracy even when the circles L_1 and L_2 are very close to one another. The extension to the case of elliptical rings is sketched. An example for a circular half-ring under twisting stress is given to illustrate the convergence of the solution. The solution is applied to the study of the stress fields of a circular shaft subjected to twisting moments and consisting of two identically armored cylinders which are half-circular in cross section and have the same radii. The extension to other types of cross sections is sketched. The Dirichlet problem and the Dirichlet-Neymann mixed problem are studied for a semi-circle to find the function $\phi_1(z)$ regular in the semi-circular region S . Finally, the case of plane deformation of a semi-circle with the arc free of external forces and the components of displacement given on the diameter is studied. It is shown that the fundamental mixed problem of the theory of elasticity for a semi-circle reduces to the singular integral-differential equation

Card 2/3

L 32971-06

ACC NR: AT6016915

$$(\kappa + A) \varphi_1(t_0) + \overline{\delta(t_0)} = \frac{1}{\pi i} \int_{\gamma} [-\overline{\delta(t)} + (A - \kappa) \varphi_1(t)] \frac{dt}{t - t_0} +$$

$$+ \frac{1}{\pi i} \int_{\gamma} [-A \overline{\delta(t)} + (1 - A\kappa) \varphi_1(t)] \frac{dt}{t - \frac{t_0}{R^2}} + F(t_0), \quad \text{or}$$

$$\overline{\delta(t_0)} + \frac{1}{\pi i} \int_{\gamma} [\overline{\delta(t)} + 2\kappa \varphi_1(t)] \frac{dt}{t - t_0} -$$

$$- \frac{1}{\pi i} \int_{\gamma} [\kappa \overline{\delta(t)} + (\kappa^2 + 1) \varphi_1(t)] \frac{dt}{t - \frac{t_0}{R^2}} = \Lambda(t_0),$$

Orig. art. has: 7 figures, 2 tables, 275 formulas.

SUB CODE: 12,20/

SUBM DATE: 13Aug65/

ORIG REF: 015/

OTH REF: 004

Card 3/3

SHERMAN, D.M. (L'vov)

Effect of the ligation of one carotid artery on the bioelectric
activity of the brain in rabbits. Pat. fiziol. i eksp. terap. 9
no.3:66-67 My=Ja '65. (MIRA 18:9)

L 31850-66

ACC NR: AP6021317

(H)

SOURCE CODE: UR/0390/65/028/005/0550/0550

AUTHOR: Shorman, D. M.

29

B

ORG: none

TITLE: Effect of dithilin and diplotsin on the outcome of traumatic shock

SOURCE: Farmakologiya i toksikologiya, v. 28, no. 5, 1965, 550

TOPIC TAGS: injury, rabbit, drug effect, blood pressure, biologic respiration, reflex activity, bioelectric phenomenon

ABSTRACT: Experiments were conducted on 92 rabbits which were struck 300-400 blows on the hip with an iron bar. During the experiment the change in arterial pressure, pressor sinocartid reflexes, respiration, temperature in the right intestines and muscles, bioelectric activity of the brain and heart, reaction of the animal to pain and temperature irritations, etc., were studied. The basic indicators were the survival and length of life of the experimental animals.

In 15 control experiments trauma causes a reaction with 4 phases: the agitation phase, parabiotic, transitional phase, torpid shock phase, and collapse phase. After this, terminal state and death

Card 1/2

UDC: 617.001.36-092.9-085.785.3+615.785.3

L 31850-66

ACC NR: AP6021317

occured. In 13 control experiments the trauma reaction was limited to only the agitation phase, resulting in collapse and death of the animals.

In the experimental series trauma was applied after a single venous administration of dithilin or diplatsin in a dose of 1 mg/kg. Artificial respiration was applied. Of 10 rabbits given dithilin, 5 survived; of 10 rabbits given diplatsin, 4 survived.

Administration of dithilin or diplatsin 15-30 minutes after trauma in the torpid shock phase also increase the survivability of the animals. The protective action of dithilin lasted 30 minutes, of diplatsin, for 3 hours. Artificial respiration, without the myo-relaxants, did not affect the course of traumatic shock and survivability.

The data indicate these drugs alter the dynamics of the course of traumatic shock. [JPRS]

SUB CODE: 06 / SUBM DATE: 09Jul64

Card 2/2 JS

SHERMAN, D. M.

Thermoregulation disorders in traumatic shock. Eksper. khir. i
anest. no.2:91-93 '62. (MIRA 15:6)

1. Iz eksperimental'noy laboratorii L'vovskogo okruzhnogo voyennogo
gospitalya.

(SHOCK) (BODY TEMPERATURE--REGULATION)

SHERMAN, F.M. (Lvov)

Effect of heating on the results of treating experimental shock with
N.G. Belen'kii's serum. Trudy Kiev. nauch.-issl. inst. pereb. krov. i
neotlozh. khir. 3:116-152 '61. (MIRA 17:10)

С. 111, . . . и т. д., А. В.

25515

Исторический очерк о деятельности и деятельности в области. Журнал "История" - Ленин.
Исторический очерк о деятельности, деятельности, деятельности, деятельности и деятельности,
VII. 23, 1919, с. 21 - 28

С. 111, . . . и т. д.

SHERMAN, D.S., inzhener; SHAMAROVA, T.A., redaktor; KUZ'MIN, G.M., tekhnicheskii redaktor.

[Aid for workers in linear measurements] Posobie dlia rabochikh na liniinykh izmereniakh. Izd. 3 Moskva, izd-vo geodezicheskoi lit-r7, 1955.
[Microfilm] (MIRA 8:5)
(Base measuring)

SUNDAKOV, Yakov Arnol'dovich; ~~SHERMAN, D.S.~~, redaktor; KHROMCHENKO, P.I.,
redaktor izdatel'stva; KUZ'MIN, G.M., tekhnicheskiy redaktor

[Geodetic work for the construction of multistory buildings]
Geodezicheskie raboty pri stroitel'stve mnogoetazhnykh zdaniy.
Moskva, Izd-vo geodezicheskoi lit-ry, 1956. 222 p. (MLRA 9:8)
(Building) (Geodesy)

SHERMAN, Daniil Savel'yevich; LARIN, B.A., kand. tekhn. nauk, red.; ZUBAKOV, A.G.,
red. izd-va; PREYS, E M., tekhn. red.

[Manual for processing base-line measurements in first-order,
second-order, and third-order triangulation] Rukovodstvo po ka-
meral'noi obrabotke bazisov 1 2 i 3 klassov. 3. izd., perer. i dop.
Moskva, Izd-vo geodez. lit-ry 1961. 179 p. (MIRA 14:10)
(Surveying—Tables, etc.)

SHERMAN, D.S., red.; KOMAR'KOVA, L.M., red.izd-va; ROMANOVA, V.V.,
tekhn. red.

[Safety rules for topographic and geodesic work. In effect from
October 1, 1962]Pravila po tekhnike bezopasnosti na topografo-
geodezicheskikh rabotakh. Vvoditsia v deistvie s 1 oktiabria
1962. g. Moskva, Izd-vo geodez.lit-ry, 1962. 73 p.

(MIRA 16:2)

1. Russia (1923- U.S.S.R.)Glavnoye upravleniye geodezii i karto-
grafii.

(Topography--Safety measures) (Geodesy--Safety measures)

KRIVOY, TS.P.; TREYSTER, Yu.Ya.; SHERMAN, E.M.

Automatic control of the blast distribution to blast furnace
tuyeres. Metallurg 7 no.2:6-8 F '62. (MIRA 15:3)

1. Tsentral'naya laboratoriya avtomatiki.
(Blast furnaces) (Automatic control)