

SOURCE: Svarochnoye proizvodstvo, no. 6, 1965, 15-17

TOPIC TACS: build-up sequence, press tool, press bushing, press die, carbon  
bushings, metal bushings, metal parts, metal parts.

"APPROVED FOR RELEASE: 07/16/2001

CIA-RDP86-00513R001756120013-5

NO REF ID#:

DATE: 00/00/00

APPROVED FOR RELEASE: 07/16/2001

CIA-RDP86-00513R001756120013-5"

TOLSTOV, K. D.

"Mass and Trajectory of Fission Products of Plutonium," *Uspekhi Fiz. Nauk*, 37, No.3, 1949.

"APPROVED FOR RELEASE: 07/16/2001

CIA-RDP86-00513R001756120013-5

TOLSTOV, K. D.

"Uranium-235 in 'Tukholite'", Uspekhi Fizicheskikh Nauk, Vol. XL, No. 3, 1950

TRANSLATION AVAILABLE In W-12587, 4 Aug 1950.

APPROVED FOR RELEASE: 07/16/2001

CIA-RDP86-00513R001756120013-5"

TOLSTOV, K. P.

600  
RMC

✓4076 AEC-tr-2435((Pt. 1) (p.85-109))  
AVERAGE NEUTRON VELOCITIES IN VARIOUS MEDIA.  
K. D. Tolstov, F. L. Shapiro, and I. V. Shtrankh. p.85-  
108 of CONFERENCE OF THE ACADEMY OF SCIENCES  
OF THE USSR ON THE PEACEFUL USES OF ATOMIC  
ENERGY, JULY 1-6, 1955. SESSION OF THE DIVISION OF  
PHYSICAL AND MATHEMATICAL SCIENCES. (Transla-

tion). 14p.  
This paper was originally abstracted from the Russian  
and appeared in Nuclear Science Abstracts as NSA 9-7892.

PMI sent

*Tolstoy, K. D.*

*9 F  
RM*

*Investigation of the parameters of uranium-graphite heterogeneous systems by the prism method.* — L. V. Grod, I. I. Reznik, G. I. Kozmets, Z. E. Lazareva, K. D. Tolstoy, E. E. Leibnitz, I. M. Frank, F. L. Shapiro, and I. V. Shtrankh.

*Soviet Akad. Nauk SSSR po Miruomu Izdatelstvenyyu Atomnoi Energii, Zaschitnye Otdel. Fiz.-Mat. Nauk* 1955, 21-50 (English summary, 51-2). — The buckling,  $x^2$ , the multiplication factor  $k$ , and the thermal utilization factor  $\theta$  in a U-graphite heterogeneous system with const. slug diam. were investigated as a function of the U concn.,  $C$ , in a Cd-shielded prism of  $180 \times 180 \times 420$  cm. The equidistant channels in the graphite could be filled either with U slugs or graphite rods.  $C$  in the prism was varied by removing metal from the channels. An at. concn. of  $N/N_e = 0.013$ , corresponding to a spacing of about 20 cm, is the arbitrary unit. About half of the channels were of larger diam. so that an air gap existed around the slugs; by moving the slugs from one channel into a wider one, the gap effect could be measured. The values  $x^2$  and  $k$  were determined according to the exponential method; both had a max. at  $C = 0.8$ , and were 0 at  $C = 0.3$ , and 1 at  $C = 1.0$ ;  $\theta$  was obtained from Cd-ratio measurements; a formula was derived

$(1 - \theta)/\theta = (g/C) - k$ , where  $g$  and  $k$  are consts. This agreed with the results of the diffusion theory,  $k/\theta$  is a linear function of  $C$ ; by extrapolating to  $C = 0$   $k$  can be split into two empirical factors  $\nu_1$  and  $\varphi$ .  $\nu_1$  is the no. of neutrons generated per thermal neutron captured; it is  $1.34 \pm 0.02$ .  $\varphi$  is the probability of fission neutrons reaching the thermal energy region. Thus  $1 - \varphi$  is the radiation capture in U, partially compensated by epicedium  $U^{235}$  fission. It is a linear function of  $C$ . Both air gaps and  $H_2O$  gaps were tried for cooling. At  $C < 1.5$   $H_2O$  has a neg. effect on  $k$  and at  $> 1.5$  a pos. one; the air gap influences  $\theta$  mostly, but both gap effects of air or  $H_2O$  were smaller than anticipated from the elementary diffusion theory; this is simply due to the one-velocity-group treatment, which is mathematically insufficient.

Werner Jacobson

(2)

TOLSTOV, K. D.

1100  
-RML

14091 AEC-tr-2435((Pt. 1)(p.39-58))

INVESTIGATION OF THE PARAMETERS OF HETERO-

GENEOUS URANIUM-GRAPHITE SYSTEMS BY THE

PRISM METHOD. L. V. Groshev, O. I. Kozinets, I. E.

Lazareva, K. D. Tolstov, E. L. Feinberg, I. M. Frank,

F. L. Shapiro, and F. V. Shtanikh. p.39-58 of CONFER-

ENCE OF THE ACADEMY OF SCIENCES OF THE USSR ON

THE PEACEFUL USES OF ATOMIC ENERGY, JULY 1-6,

1955. SESSION OF THE DIVISION OF PHYSICAL AND

MATHEMATICAL SCIENCES. (Translation). 20p.

This paper was originally abstracted from the Russian and  
appeared in Nuclear Science Abstracts as NSA 9-7925.

TOLSTOV, K.D.

1946

Average neutron velocities in various media. K. D.

Tolstov, V. L. Shapiro, and I. V. Shfranikh. Sessiya Akad. Nauk S.S.R. po Miriamu Tsplotoriumu Atomnoi Energetiki, Zasedaniyu Otdel. Fiz.-Mat. Nauk 1955, 103-29 (English summary, 120-31).—The process of slowing down neutrons in the vicinity of the thermal equil. region ( $v = 2200 \text{ m./sec.}$ ) is influenced by many factors. The av. energy and spectrum of the neutrons depend on the mean energy transferred owing to inelastic collisions, on the capture cross section, the structure and vol. of the moderator, and the temp. of the medium. The influence of these factors on the av. velocity  $\bar{v}$  and on the spectrum of thermal neutrons was studied. The  $\bar{v}$  was detd. by the ratio flux of neutrons to their d. The d. was measured with a  $\text{BF}_3$  ionization chamber, the flux with a Geiger counter, which counted the  $\gamma$ -quanta owing to neutron capture in Cd. The  $\bar{v}$  was obtained by measuring the transmission of neutrons by a  $1/\nu$  absorber. The temp. function of the diffusion length was detd. by the exponential method. The changes with temp. of the diffusion coeff. of the neutrons was found by measuring the variation of the neutron density when the medium was heated; the mean free transport path changes were found by measuring the changes of the albedo. Expts. were done with prisms of  $60 \times 60 \times 100 \text{ cm.}$  or  $20 \times 100 \times 120 \text{ cm.}$  of graphite (I), paraffin (II),  $\text{H}_2\text{O}$ , and  $\text{H}_2\text{O-B}$ , with and without increments having  $1/\bar{v}$  absorption. Also, heterogeneous systems with I and  $\text{H}_2\text{O}$  as moderators, like the U-I

system, were investigated, and it was found that in I, II, and  $\text{H}_2\text{O}$  the neutrons attain thermal equil. and their velocity spectrum is Maxwellian. If the lifetime of the neutrons within the prism is decreased to  $1/20$  by introducing neutron absorbers, or to  $1/50$  by diminishing the dimensions of the prism,  $\bar{v}$  is increased up to 40%. Neutrons in the vicinity of the thermal equil. suffer many collisions before they lose their energy; the results agree with the theory of slowing down of neutrons in a cryst. medium. If I is heated up to  $300^\circ$ , the increase of  $\bar{v}$  is  $6 \pm 2\%$  larger than is that of the square of the diffusion length, thus the mean free transport path  $\lambda_{fr}$  decreases  $5 \pm 1\%$ . The  $\bar{v}$  was measured at  $20$  and  $300^\circ$  in the U-I system with various U contents, and it was found that for any concn.  $\bar{v}$  in a homogeneous system is higher than in such a heterogeneous one. It was found by calcs. and measurement that the neutron spectrum emitted by the moderator surface is not only detd. by the neutron spectrum in the moderator, but also by the energy function of  $\lambda_{fr}$ . Therefore, the spectrum of the

neutrons in the medium is different from that of the neutrons that have passed through the prism. W. J.

(2)

RMS DM

*Tol'st'oy K. D.*

✓ Measurement of temperature effects in uranium-graphite over a layer of I adjacent to the slug. At identical U concns.,  
 Subcritical systems. B. P. Ad'yasevich, O. T. Koznits, the  $\theta$  of a heterogeneous system can be increased over that  
 K. D. Tolstoy, I. M. Frank, F. L. Shapiro, and I. V. of a homogeneous one by inhomogeneously heating the  
 Smirnovich, Sessiya Akad. Nauk S.S.R. po Mirovym moderator. In order to evaluate correctly the temp. effect  
*Ispol'zovaniyu Atomnoj Energii, Zasedaniyu Otdel. Fiz.* on the  $\nu$  escape probability,  $\nu$ , cold H<sub>2</sub>O and H<sub>2</sub>O at 80° were  
*Mat. Nauk 1955, 132-54* (English summary, 154-5). The circulated alternately through the slugs, the temp. of the I  
 temp. effects on the multiplication const.  $k_{\infty} = \nu_{\infty} \theta$  and being kept const. The variation of  $\varphi$  was obtained from  
 also on all the factors thereof were investigated by two exponential measurements of the buckling  $x^2$  and from mea-  
 120 X 120 X 200 cm. prisms, each in its own thermostat, surements of the influence of heating on the ep-Cl neutron  
 heated to 300-450°. U-graphite (1) lattices were used with d. in the vicinity of the source. The temp. effect of the  
 and a Po-Be neutron source were employed. The thermal integral is found to be  $\delta\rho/\rho_0 = (1.95 \pm 0.1) \times 10^{-4}$  per de-  
 utilization factor  $\theta$  was measured by the Cl ratio method; gree. The temp. effect on  $k_{\infty}$  was detd. by measuring  $x^2$   
 a pos. temp. effect was found which increased with decreas- by the exponential method. The temp. changes of  $\nu_{\infty}$  (no.  
 ing U concn. and was greater in the presence of cooling H<sub>2</sub>O. of neutrons generated, no. of neutrons captured) were calc.  
 This effect was already predicted by the elementary diffu- from the changes of  $\varphi$  and  $\theta$  and from the temp. effect on  
 sion theory, but the theory furnished too high values in the  $k_{\infty}$ . It was found that  $\nu_{\infty}$  has a neg. temp. effect, which is  
 absence and too low values in the presence of H<sub>2</sub>O. This roughly proportional to the variation of the mean energy of  
 discrepancy was due to the fact that  $\theta$  increased when the thermal neutrons, caused by the heating of the system;  
 neutrons were cooled in H<sub>2</sub>O upon entering the slug. The DOS. effect on  $\theta$  increased also, if the cooling was extended  
 Werner Jacobson

"APPROVED FOR RELEASE: 07/16/2001

CIA-RDP86-00513R001756120013-5

FOR STOVA K.D.

APPROVED FOR RELEASE: 07/16/2001

CIA-RDP86-00513R001756120013-5"

USSR/Nuclear Physics - Elementary Particles

C-3

Abst Journal : Referat Zhur - Fizika, No 12, 1956, 33920

Author : Dul'kova, L. S., Romanova, T. A., Sokolova, I. B., Sukhov, L. V.,  
Tolstov, K. D., Shafranova, M. G.

Institution : None

Title : Interaction of 300-Mev  $\pi^-$ -Mesons with Protons, Deuterons, and  
Nuclei of a Photographic Emulsion

Original  
Periodical : Dokl. AN SSSR, 1956, 107, No 1, 43-46

Abstract : AIKFI plates of the "p" type, enriched with H or loaded with  
D by impregnating in a 30% water solution of lithium acetate,  
were radiated in the phasotron of the Institute for Nuclear  
Problems, Academy of Sciences USSR by  $\pi^-$ -mesons of  $225 \pm 8$  Mev.  
The H content reached  $6 \cdot 10^{22}$ , and the D content reached  
 $3 \cdot 10^{22}$  per  $\text{cm}^3$ . The presence of Li made it possible to  
control the evenness of the loading. The increased value of

Card 1/2

USSR/Nuclear Physics-Elementary Particles

C-3

**Abst** Journal : Referat Zhur-Fizika, No. 12, 1956, 33920

pH of the lithium acetate contributed to a reduction in regression. The radiation was carried out up to a density of  $10^4$  to  $10^5$  tracks per  $\text{cm}^2$ . The examination was made by areas end along the track. The average free path for all the processes, including scattering by an angle greater than  $20^\circ$  was  $88 \pm 5\%$  of the geometric. The principal contributions are made by processes of inelastic scattering and absorption with star formation.

The area inspection method was used to trace 1,240 stars. A distribution was made by the number of rays. Fifty cases of scattering by H and 11 cases of scattering by D were found; the elastic-scattering sections were respectively  $H = 14 \pm 3.6$  millibarn and  $D = 15 \pm 5.5$  millibarns. The scattering by D is strongly anisotropic. A histogram is given for the differential scattering of  $\pi^-$ -mesons by H in a center of gravity system. A discrepancy is noticed from the theoretical curve for small scattering angles.

Card 2/2

"APPROVED FOR RELEASE: 07/16/2001

CIA-RDP86-00513R001756120013-5

APPROVED FOR RELEASE: 07/16/2001

CIA-RDP86-00513R001756120013-5"

TOLSTOV, K.D.

BELYAKOV, V.A.; IVANOVA, L.N.; KOZLOVA, L.G.; TOLSTOV, K.D.

Experiments with 600 micron layers from the "R" Emulsion of the  
Motion Picture and Photography Scientific Research Institute.

Zhur. nauch. i prikl. fot. i kin. 2 no.5:325-329 S-0 '57.

(MIRA 10:11)

1. Ob'yedinenyy institut yadernykh issledovaniy.  
(Photographic emulsions)

120-6-30/36

AUTHORS: Otroshchenko, V.A., Sviridov, V.A., Tolstov, K.D.,  
and Shal'nikov, A.I.

TITLE: Solid Hydrogen Targets on the Surface of Photographic  
Emulsions (Tverdyye vodorodnyye misheni na poverkhnosti  
fotoemul'sii)

PERIODICAL: Pribory i Tekhnika Eksperimenta, 1957, No.6,  
pp. 110 - 111 (USSR).

ABSTRACT: It is difficult to study interactions between elementary particles and protons and deuterons which are included in nuclear emulsions because their number is small compared with the total number of nucleons bound in the nuclei of the emulsion. This is still true even when the emulsion is specially loaded with deuterium and hydrogen. To remove this difficulty, it is convenient to have a target of solid hydrogen or deuterium deposited directly on the surface of the emulsion. In this method of preparation of targets the temperature of the emulsion cannot be greater than 12 to 15 °K. Because of this, the temperature dependence of the sensitivity of NIKFI-R emulsions was investigated (Ref.1). Already at 20 °K, the sensitivity of emulsion is down by a factor of 2 and therefore it is difficult to use this emulsion with mini-Card1/2  $\mu$ m ionisation particles. However, different types of

120-6-30/36

Solid Hydrogen Targets on the Surface of Photographic Emulsions.

emulsion have been described (Refs. 2 and 3) which have the property that their sensitivity falls much lower with temperature. To obtain sufficiently thick solid hydrogen targets on top of emulsions, a special device shown in Fig. 1 was used. The photoemulsion, 2, kept at the bottom of the plate-holder, 1, was surrounded by liquid hydrogen. Through the tube, 3, deuterium gas was introduced into the plate-holder and slowly solidified on top of the emulsion due to the cooling effect of the surrounding liquid hydrogen. After this, the Dewar containing the hydrogen was evacuated and the plate-holder was lifted into position for irradiation by a beam from an accelerator. In the case of the solid hydrogen target, liquid helium was used as the cooling agent. V.I. Veksler and I.B. Danilov collaborated. There are 2 figures, 2 diagrams and 3 references, 2 of which are Slavic.

ASSOCIATION: United Institute for Nuclear Studies  
(Ob'yedinennyj Institut yadernykh issledovaniy)  
Department of Low Temperature Physics of MGU  
(Kafedra Fiziki nizkikh temperatur MGU)

SUBMITTED: May 20, 1957.

AVAILABLE: Library of Congress.  
Card2/2

TOLSTOV K.D.

AUTHORS: Bogachev, N. P., Van Shu-Fen', Gramenitskiy, I. M., Kirillova, L. F., Lebedev, R. M., Lyubimov, V. B., Markov, P. K., Merekov, Yu. P., Podgoretskiy, M. I., Sidorov, V. M., Tolstov, K. D., Shaframova, M. G.

TITLE: The Interaction of 9 Bev Protons With Nuclei in Photo-Emulsion  
(Vzaimodeystviye protonov s energiyey 9 Bev s yadrami foto-emul'sii)

PERIODICAL: Atomnaya Energiya, 1958, Vol. 4, Nr 3, pp. 281 - 284 (USSR)

ABSTRACT: The photoemulsion H  $\mu$  KФИ-P with a layer of about 450  $\mu$  was irradiated with proton within and out of the vacuum chamber of the 9 Bev synchrophasotron. The mean range of 9 Bev protons for an interaction is  $34,7 \pm 1,5$  cm. (The scattering for angles below  $5^\circ$  was not taken into account). 258 cases of a nuclear interaction were observed. The mean number of fast particles n generated a process of interaction to  $3,4 \pm 0,7$ . The angular distribution of these particles shows a clearly preferred forward motion. The mean number of black and grey traces  $N_n$  - the recoil nuclei not being considered - is  $8,3 \pm 0,5$ . From 249 found stars 18 can be considered to constitute in interaction of the initial protons with "free" or "quasifree" protons. 13 stars can be considered to represent

an interaction between protons and "quasifree" neutrons. All of them have an odd number of traces, and in the point of formation of the stars  $\beta$ -traces can be observed. The mean number of fast particles in these 13 star traces is 3,1  $\pm$  0,3. There are 5 figures, 1 table, and 7 references, 1 of which is Slavic.

SOV/89-5-4-7/24

AUTHORS: Barashenkov, V. S., Van Shu-fen<sup>1</sup>, Tolstov, K. D.

TITLE: Collisions of Protons With an Energy of 9 BeV With Nucleons  
("stolknoveniya protonov s energiyey 9 Bev s nuklonami")

JOURNAL: Atomnaya energiya, 1958, Vol 5, Nr 4, pp 453-454 (USSR)

ABSTRACT: The nuclear emulsion **NIKFI-R**, which has a thickness of layer of  $\sim 450\mu$ , was bombarded with protons of  $\approx 9$  BeV (synchro-phasotron of the Ob'yedinenyyi institut yadernykh issledovaniy (United Nuclear Research Institute)). From 372 stars 39 were counted as p,p-collisions and from 264 stars 11 were counted as p,n-collisions. Three of the 39 stars could be fixed as being elastic p,p-collisions. The following further values were obtained by counting (average number of the particles taking part in the p,p- and p,n-collisions respectively):

$$\bar{n}_p = 3,8 \pm 0,3$$

$$\bar{n}_n = 2,8 \pm 0,6$$

Card 1/2  $\bar{n}_p^{(s)} = 3,1 \pm 0,3$  (s) = fast particles

SOV/89-5-4-7/24

Collisions of Protons With an Energy of 9 BeV With Nuclei

$$\bar{n}_n^{(s)} = 2,5 \pm 0,6 \quad (s) = \text{fast particles}$$

The angular distribution for  $n_p^{(s)}(\Theta)$  is recorded graphically. The average loss of energy for the forming of ions at p,p-collision amounts to about 50% of the energy of the primary protons.

E. G. Bubelev, V. M. Mal'tsev, and Ten-Gyn assisted in the theoretical calculations. L. F. Kirillova and V. A. Belyakov assisted in experiments. There are 1 figure, 1 table, and 3 references, 2 of which are Soviet.

SUBMITTED: August 1, 1958

Card 2-2

BARASHENKOV, V. S., BELYAKOV, V. A., BUBELEV, E. G., MALTSEV, V. M., TOLSTOV, K. D.,  
WANG SHOU FENG, and TEN GYN.

"Multiple Production of Particles in Collisions between 9 GeV Protons and  
Nucleons." Nuclear Physics, vol. 9, No. 1, Nov 1958.

Joint Inst. Nuclear Research, Lab Theoretical Physics and High Energy Lab., Dubna.

Abstract: Some theoretical calculations pertaining to multiple production of particles in nucleon-nucleon collisions at 7-10 GeV were presented in ref. 1. Some preliminary experimental results obtained by irradiating photographic emulsions with proton beam from the synchrocyclotron of the Joint Inst. Nuclear Research were given in ref. 2. In the present paper we compare the theoretical results of ref. 1. with the results of some new experiments. 372 stars of which 50 were classified as proton-nucleon collisions, were recorded in NIKFI-R photographic emulsions along the tracks of  $\sim$  9 GeV protons accelerated in the JINR proton synchrocyclotron. The mean number of charged particles created in these collisions was  $3.6 \pm 0.5$ . The angular distribution of fast charged particles is obtained. As a whole the experimental results agree with the statistical theory of multiple particle production within the limits of the experimental  $\pm 2\%$  errors. Some discrepancy is evident in the small angle range and may be due to the contribution of non-central impacts and to asymmetry of the angular distribution in the c.m.s.

(23.3000)

66838

SOV/77-4-6-5/16

AUTHOR: Belyakov, V.A., Kozlova, L.G., Sviridov, V.A. Tolstov,  
K.D.

TITLE: Dependence of the Sensitivity of Nuclear Emulsions on  
Temperature Within the Range of 2-300° K

PERIODICAL: Zhurnal nauchnoy i prikladnoy fotografii i kinematografii  
1959, Vol 4, Nr 6, pp 427-429 (USSR)

ABSTRACT: The author reports on recent Soviet study of the dependence of the recording properties of various nuclear emulsions on temperature within the range of 2-300° K. The results of the first experiments were published in the paper of N.A. Dolina, V.A. Sviridov, K.D. Tolstov and E.N. Tsyganov [Ref 1]. Subsequently, an attempt was made to improve the recording properties of the emulsion NIIFI R 400μ by a change in the processing conditions. Curve 1 of the graph (taken from the paper of V.A. Belyakov, L.G. Kozlova, V.A. Sviridov, K.D. Tolstov and E.N. Tsyganov [Ref 2]) corresponds to the normal processing conditions of emulsions, which with

Card 1/3

66838

SOV/77-4-6-5/16

Dependence of the Sensitivity of Nuclear Emulsions on Temperature  
Within the Range of 2-300° K

regard to the correlation trace density of fog are most suitable for exposure at room temperature. Curve 2 corresponds to intensified development conditions, the fog increasing in this case by 50%. NIKFI low-temperature emulsions without silver iodide gave better results. Under normal processing conditions, the relative sensitivity at 20° K for the best series of emulsions was equal to  $45 \pm 3\%$ , and the absolute density of the tracks of the relativistic particles amounted to 17 grains for  $100\mu$ . The microphotograph shows the tracks of  $\pi$ -mesons with an energy of 340 Mev and nuclear fission at an exposure of the emulsion at 20° K. Fine-grained emulsions developed by N.A. Perfilov, N.R. Novikova and Ye.T. Prokof'yeva [Ref 3] showed at 75° K a relative sensitivity of 75%. The density of the grains on the tracks of the relativistic particles at 300° K amounted to 46 grains per  $100\mu$ . Experiments with Ilford ("Il'ford") G-5 600 $\mu$  layers [Ref 4] were also carried out (see

Card 2/3

X

66838

SOV/77-4-6-5/16

Dependence of the Sensitivity of Nuclear Emulsions on Temperature  
Within the Range of 2-300° K

Table). The grain density at exposure within the range of 2-215° K averages 15-17 grains per  $100\mu$  of particle track. The fog is approximately constant. The layers were processed under conditions recommended by the firm of Ilford. Comparative data on NIKFI and Ilford emulsions are given in the graph. There are 1 graph, 1 microphotograph, 1 table and 4 references, 3 of which are Soviet and 1 English.

ASSOCIATION: Ob"yedinenyyi institut yadernykh issledovaniy (Joint Institute of Nuclear Research) ✓  
SUBMITTED: September 23, 1957  
Card 3/3

To Ls Toy, K.D.

2) (8)  
ARTICLES:  
Barashenkov, V. S., Belyakov, V. A., Sov. 59, 74-12/26  
Yan Shun-sen, Gligorov, V. V., Dolzhikov, M. P., Evtushenko, I. P.,  
Lebedev, R. N., Matveev, V. M., Savchenko, T. N., Solntsev, R. D.,  
Tseytlin, B. M., Shekhter, M. G., Iao Ching-shih  
TITLE:  
The Interaction of Fast Nucleons With Protons of the Photoemulsion KLEPTA.

PUBLICATIONS:  
Izmeriya sverchastoty, 1959, Vol. 7, No. 4, pp. 516-517 (USSR)

ABSTRACT:  
The present paper deals with the interaction between 9 Bev-protons, which were accelerated in the beam of the synchrotron-cyclotron of the Obninskayu Institute Yadernyy Isledovaniyu (Joint Institute of Nuclear Research), and the nuclei of a polyacrylate of the KLEPTA type. The results of these measurements are shown by a table. On the basis of the data thus found it is possible to draw several conclusions as to the mechanism of the interaction between a fast proton and a nucleus. If the primary nucleon-nucleus collision is an inelastic interaction between nucleon and channel, the velocity of the center of mass in an interaction of silver and bromine with the channel will be considerably less than in an interaction with light nuclei. Therefore, also the number of  $\alpha$ -particles

must be considerably greater. In the experiment, the numbers of  $\alpha$ -particles for light and heavy nuclei are, however, nearly the same. This is explainable on the basis of the cascade mechanism of interaction, in which the energy of the  $\alpha$ -particles decreases rapidly in cascade collisions. The multiplicity of the particles produced decreases simultaneously. In the case of the greater number of  $\alpha$ -particles nucleons are concerned, which may be explained by the cascade mechanism of nucleon-nucleus interaction. Also the agreement between the traversals  $\sigma_{\text{KLEPTA}}$  for  $\alpha$ -protons originating from interactions with light and heavy nuclei points in the direction of the interaction cascade mechanism. Besides, a search was made for strange particles by employing the method of investigating according to areas. The cross section of the production of  $\Lambda$ -particles with an energy of  $E < 10$  Bev in a medium-weight nucleus of the photoemulsion amounts to  $(5 \pm 2) \times 10^{-27}$  cm<sup>2</sup>. Besides, the amount of the production cross section, the wide angular distribution of the  $\Lambda$ -mesons, as well as other facts indicate that a noticeable fraction of

slow strange particles is produced in an intranuclear cascade process. Furthermore, the medium-weight energy losses of a fast nucleon are evaluated in the case of a single nucleon-nucleus collision. A 9-bev-proton gives up an average of  $(5.1 \pm 0.8)$  per to a medium-weight nucleus of the photoemulsion, which amounts to  $(60 \pm 10)\%$  of its initial energy. 4.05 Bev are used for the production of pions, and 1.05 Bev are transferred to the nucleus of the nucleon. As a proton in an medium-weight nucleus undergoes approximately 2 collisions in the photoemulsion, a single nucleon-nucleus collision, losses  $\Delta E = 5 \pm 10\%$  of its initial energy. By means of other measurements of the pion energy spectrum carried out independently of the present paper in a nucleon-nucleus collision,  $\Delta E = 10 \pm 10\%$  is obtained. The statistical theory of multiple-particle production furnishes  $\Delta E = (40 - 50)\%$ . The authors thank G. Basov and V. Kuznetsov and Yu. M. Krikunov for their help in the measurements, and L. Popov for his assistance in analyzing measuring results. There are 1 table and 1 reference.

Card 1/3

Card 2/3

Card 3/3

85363

S/120/60/000/005/037/051  
E032/E314*21.5.200*  
AUTHORS: Zavaritskiy, N.V., Sviridov, V.A. and Tolstov, K.D.TITLE: Sensitivity and Thermal Conductivity of Nuclear  
Emulsions at Low TemperaturesPERIODICAL: Pribory i tekhnika eksperimenta, 1960, No. 5,  
pp. 131 - 132TEXT: НИКФИ-Р (NIKFI-R) and Ilford G-5 nuclear emulsions were  
investigated. The thermal conductivity of the emulsions was  
measured by the method described in an earlier paper (Ref. 1). In  
the temperature interval 4 - 1.5 K the coefficient of thermal  
conductivity of the NIKFI-R emulsions can be expressed by the  
formula:

$$K \sim 2.2 \cdot T^{2.8} \cdot 10^{-5} \text{ W/cm. } ^\circ\text{K} \quad (1) .$$

A description is given of a low-temperature device which was  
used to cool the emulsions below 1 K. The emulsions are  
cooled by connecting them through a heat-conducting rod to a  
block of an adiabatically demagnetised material. The sensitivity  
was measured after irradiating the emulsions with  $\text{Co}^{60}$   $\gamma$ -rays

Card 1/3

85363

S/120/60/000/005/037/051  
E032/E314

Sensitivity and Thermal Conductivity of Nuclear Emulsions at Low Temperatures

at 0.1, 0.3, 1.6 and 300 °K. The results obtained are summarised in the following table:

Emulsion	Temperature, °K	300	1.6	0.3	0.1	Absolute sensitivity at 300 °K (blobs/100μ)
NIKFI-R		100%	$(36^{+15}_{-10})\%$	$(31^{+15}_{-10})\%$	$(21^{+15}_{-10})\%$	~ 60
Ilford G-5		100%	$(69^{+15})\%$	-	$(70^{+15})\%$	~ 25

The sensitivity at 300 °K was taken at 100%. Acknowledgments are expressed to P. L. Kapitsa for collaboration in this work.

Card 2/3

85363

S/120/60/000/005/037/051  
E032/E314

Sensitivity and Thermal Conductivity of Nuclear Emulsions at  
Low Temperatures

There are 2 figures, 1 table and 1 Soviet reference.

ASSOCIATION: Ob'yedinenyy institut yadernykh issledovaniy  
(Joint Institute for Nuclear Studies)

SUBMITTED: August 13, 1959

X

Card 3/3

PEREYGIN, V.P.; TOLSTOV, K.D.

Cross section of the reaction  $\text{Li}^6(n,\alpha)\text{H}^3$  for 2.15 Mev neutrons.  
Atom. energ. 9 no.6:488-489 D '60. (MIRA 13:12)  
(Lithium--Isotopes) (Tritium)

TOLSTOV, K. D.

Card 5/1

SUMMITED: May 11, 1960

ASSOCIATION: Obshodimnyy Institut Yadernyykh Issledovaniy (OII)

Institute of Nuclear Research

Card 2/3

maximum value of  $4 \cdot 10^{-14}$  cm. The authors summarize the results as follows:

Average momenta of protons ( $0.870 \pm 0.001$ ) MeV/c, average transverse momenta ( $0.170 \pm 0.001$ ) MeV/c, asymmetry of angular distribution of all pions ( $-1.520 \pm 0.10$ ) pions with  $p > 0.5$  have are emitted in the forward direction, their average momentum equalling  $(0.810 \pm 0.06)$  MeV/c and agrees, with that of the protons. The authors thank Dr. J. Blümlein, Dr. L. Kesten, Dr. H. Kesten (for discussion and advice, there are 9 figures, 3 tables, and 23 references; 9 Soviet, 8 US, 1 British, 1 German, 1 Italian, 1 Japanese, and 1 Polish.

X

TITLE: *Inelastic interaction of 7 Bev  $\pi$ -mesons and Nucleons*  
 PERIODICAL: *Zhurnal eksperimental'noi i teoretičeskoy fiziki*, 1960, Vol. 39, No. 4(10), pp. 937-947

TEXT: The inelastic interaction of 7-Bev  $\pi$ -mesons with nucleons is studied in this paper. The preliminary results were communicated to the audience of the International Conference on High-Energy Physics (Kiev, Ukraine). The emulsion chamber consisted of 210 Mylar- $\rho$  (UVECH) layers with a thickness of 400  $\mu$ , 5100 interactions with the nuclei of pionization were observed. Of these, 515 inelastic interactions were analyzed (Table 1). The theoretical distribution of the charged particles was calculated by V. S. Karatachyan. Scattering scattering was eliminated by special measurements (Table 2). 439 pions and 134 protons

Card 3/3

were identified. The angular distribution of pions and the total distribution of all stars (in c.m.s.) are shown in Fig. 1. For smaller number of charged particles, the asymmetry increases strongly. This is principally due to pions with large momenta (Fig. 2). Therefore, the angular distributions are very different for pions and for neutral pions with momenta  $< 0.5$  MeV/c than in almost isotropic distribution. From the proton conserves their initial direction. From the momenta distributions of pions and nucleons, the authors conclude that the average momentum of the nucleons and of the charged pions does not depend on the increase of the number of charged particles. The same result follows from the data on the average transverse momenta  $p_T$  of protons and pions given in Table 2. Fig. 7 shows the number of neutral stars as a function of the number of charged particles. The results can be interpreted only partly by the statistical theory. The asymmetry of the angular distribution of the secondary pions can only be explained by a peripheral collision of the pion with a pion of the nucleus shell (Fig. 8 and 9). An estimate of the radius of the nuclear core gave the

24-604

X

RELENTOV, V. A., VAN SHU-SHUN, GLEBOVSKAYA, MALKHATZ,

SHUBIKOV, S. M., MELNIKOV, N. N., MITIN, V. A.,

PARKHOMENKO, V. A., SOKOLOV, K. I.

8/05/60 07/01/1982/005/048

B60/2070

8436

X

7

*TOLSTOV, K.D.*32991  
S/641/61/000/000/018/033  
B108/B102

246500

AUTHORS: Mikhaylina, K. M., Nomofilov, A. A., Romanova, T. A.,  
Sviridov, V. A., Tikhomirov, F. A., Tolstov, K. D.

TITLE: Interaction of 14.1-Mev neutrons with Li<sup>6</sup> and Li<sup>7</sup>

SOURCE: Krupchitskiy, P. A., ed. Neytronnaya fizika; sbornik statej.  
Moscow, 1961, 249 - 257

TEXT: Interaction of 14.1-Mev neutrons with Li<sup>6</sup> and Li<sup>7</sup> nuclei was studied both with targets prepared from Ilford E<sub>1</sub> photoemulsions bearing the lithium and with targets of metallic lithium isotopes. The latter method was used for small angles of the departing particles. The mean number of Li nuclei in the photoemulsion was  $2.3 \cdot 10^{19} \text{ cm}^{-2}$ . The integral neutron flux striking the emulsion at right angles was about  $10^8 \text{ cm}^{-2}$ . Altogether, 412 events were recorded on a  $2.5 \text{ cm}^2$  area. 96 events were from the reaction Li<sup>6</sup>(n,t)<sup>3</sup>H with a cross section  $\sigma = 27 \pm 6 \text{ mb}$ . Seven Li<sup>6</sup>(n,p)He<sup>6</sup> reactions with a cross section of about 5 mb were found, moreover

Card 1/2

32991  
S/641/61/000/000/018/033  
B108/B102

Interaction of 14.1-Mev...

$\text{Li}^6(n,d)\text{He}^5$  reactions with a differential cross section of  $2.15 \pm 2.5$  mb/sterad in the range between  $70^\circ$  and  $142^\circ$ . The cross section of the  $\text{Li}^6(n,n')\text{Li}^{6*} \rightarrow d + \alpha$  events was  $70 \pm 12$  mb, that of the reaction  $\text{Li}^6(n,2n)\text{Li}^{5*} \rightarrow n + p$  was equal to  $50 \pm 10$  mb. Interaction with  $\text{Li}^7$  yielded the reactions  $\text{Li}^7(n,t)\text{He}^5$ ,  $\text{Li}^7(n,n')\text{Li}^{7*}$ , and seven  $\text{Li}^7(n,d)\text{He}^5$  events. In the experiments with pure lithium targets the reactions observed were  $\text{Li}^6(n,d)\text{He}^5$  ( $\sigma = 58 \pm 10$  mb),  $\text{Li}^7(n,t)\text{He}^5$  ( $58 \pm 12$  mb),  $\text{Li}^7(n,n')\text{Li}^{7*} \rightarrow t + \alpha$ ,  $\text{Li}^7(n,d)\text{He}^6$ . The overall cross section of  $(n,r')$  and  $(n,2n)$  processes for  $\text{Li}^6$  was  $179 \pm 20$  mb. The results obtained are consistent with those of other publications. I. M. Frank, O. I. Kozinets, L. N. Katsurov, and D. I. Ivanov are thanked for help. There are 6 figures, 1 table, and 7 references: 2 Soviet and 5 non-Soviet. The four most recent references to English-language publications read as follows: Frye, G. M. Phys. Rev., 93, 1086 (1954); Battat, M. E., Ribe, F. L. Phys. Rev., 92, 80 (1953); Frye, G. M., Rosen, L. Phys. Rev. 90, 381 (1953); Moak, C. E. Phys. Rev., 92, 383 (1953).

Card 2/2

3.2100 (also 4303)

37199  
S/560/61/000/011/005/012  
E032/E514

AUTHORS: Veprik, Ya.M., Kurnosova, L.V., Razorenov, L.A.,  
Tolstov, K.D. Fradkin, M.I. and Chukin, V.S.

TITLE: Experiment on the development of photographic  
emulsions on board the second cosmic spaceship

SOURCE: Akademiya nauk SSSR. Iskusstvennye sputniki Zemli.  
no.11. Moscow, 1961. Rezul'taty nauchnykh  
issledovaniy, provedennykh vo vremya poletov vtorogo  
i tret'yeogo kosmicheskikh korabley-sputnikov, 35-41

TEXT: The second Soviet cosmic spaceship carried stacks  
of thick nuclear emulsions. Owing to the fact that the spaceship  
remained in orbit for a considerable time, the number of particles  
recorded in the emulsions was very large, which could complicate  
subsequent scanning and identification of particle tracks. It was,  
therefore, necessary to develop the emulsions before too many  
particles had been recorded. An account is given in the present  
paper of how the emulsions were in fact developed on board the  
spaceship. The operation was carried out in four stages, namely:  
1) exposure of the emulsions to the radiations for a given time,

Card 1/2

2

Experiment on the development ...

S/560/61/000/011/005/012  
EO32/E514

2) development, 3) storage of the emulsions (latent-image centres produced during this period could not be developed), 4) subsequent laboratory analysis on the Earth's surface. The whole operation was carried out in a hermetically sealed container. The emulsion stack (20 unbacked emulsions 300  $\mu$  thick each) had to be so arranged that after the exposure the emulsions could be separated from each other and the developer let in. This was done by a piston device (a schematic drawing of the latter is reproduced). After this operation the developer was removed and a stopping solution was introduced. The emulsions remained in this solution until they were returned to the laboratory for final treatment. It was found that relativistic tracks were easily visible in these emulsions, although the sensitivity to the latter turned out to be somewhat lower than usual. Two particle-track microphotographs are reproduced to illustrate the possibilities of the method. There are 3 figures.

SUBMITTED: July 7, 1961

Card 2/2

29612  
S/120/61/000/004/023/034  
E032/E514

24,6830

AUTHORS: Perelygin, V. P., Myachkova, S.A. and Tolstov, K.D.

TITLE: Introduction of beryllium grains into photographic emulsions

PERIODICAL: Pribory i tekhnika eksperimenta, 1961, No. 4,  
pp. 145-147

TEXT: Zh.S. Takibayev (Ref. 3; Zh. eksperim. i teor. fiz.,  
1953, 24, 229) is said to have been the first to introduce  
spherical metal grains into photographic emulsions. Quantitative  
experiments concerned with the determination of cross-sections  
using non-spherical beryllium grains were described by  
S.S. Vasil'yev, V. V. Komarov, A.M. Popova (Ref. 4; PTE, 1959,  
No. 1, 48). The dimensions of the grains depend on the minimum  
range of charged particles which can be recorded in an ordinary  
emulsion (3  $\mu$  approximately). However, the grains cannot be too  
small since otherwise there may be confusion as to whether the  
event takes place in the grain or the emulsion. The present  
authors have used the spark discharge method of evaporation of  
metals described by B. R. Lazarenko, N. I. Lazarenko (Ref. 5).

✓

Card 1/8 3

29612  
Introduction of beryllium grains ... S/120/61/000/004/023/034  
E032/E514

Elektroiskrovaya obrabotka metallov (Electric Spark Treatment of Metals), 1950, Gcsenergoizdat). In the case of a spark discharge between two electrodes located in a dielectric, it is found that in most cases the metal grains produced during the process are spherical in form. It is stated that the "usual circuit" was employed with  $R = 115$  Ohm,  $C = 2.8 \mu F$ ,  $V = 110$  V. The average beryllium grain diameter was about  $1.5 \mu$ . The volume of the dielectric was 50 to 100 cc and the evaporation process was continued for 60 to 90 min. At first the dielectric employed was absolute alcohol. However, the spark discharge in alcohol leads to the formation of BeO and  $Be(OH)_2$  and complex insoluble compounds. Tests were therefore made to determine whether the grains could be obtained with a spark discharge in liquefied argon. The evaporation was carried out in a dewar having a volume of about 200 cc. The argon was then driven off and the volume was filled with alcohol. In this way it was possible to obtain isolated beryllium grains and the suspension could be kept for long periods of time. In order to introduce the grains into the emulsion, the photographic plates were placed horizontally and the

*✓*

Card 2/4 3

Introduction of beryllium grains ... <sup>296L</sup>  
S/120/61/000/004/023/034  
E032/E514

suspension was poured on to it. The particles then sedimented down onto the surface and the alcohol was evaporated. The photographic plate was then covered by a wet, unbacked emulsion and the composite emulsion was placed in a 5% solution of glycerine at 15°C for 45 min. The emulsion was then removed from the glass backing and dried with filter paper. The two-layer photo plates were then placed into a water bath at 45-48°C for 3 to 5 min. In this bath the upper layer fused into the lower one and the separation boundary could not be seen through a microscope. The procedure has been successfully used with Ilford E-1, C-2, NIKFI Ya-2, (NIKFI Ya-2), T-1 and T-3 emulsions. Fig.2 shows the diameter (I) and mass (II) distributions. N in this figure is the number of grains, M is the weight of the grains in units of  $10^{-9}$  g/cm<sup>2</sup>, and d is the diameter in microns (horizontal axis). The method has been used in nuclear reaction studies with  $^{14}$  MeV neutrons. Acknowledgments are expressed to G.Ye.Belovitskiy for advice. There are 2 figures and 5 references: all Soviet.

ASSOCIATION: Fizicheskiy institut AN SSSR (Physics Institute AS USSR)

Card 3/13

X

TOLSTOV, K. D., VEPRIK (fhu), KURNOSOVA, L. V., RAZORENVO, L. A., FRADKIN, M.I., CHUKIN

"Controlled exposition of nuclear emulsions on sputniks"

Fourth International Colloquium on Photography (Corpuscular) - Munich, West  
Germany, 3-8 Sep 62

TOLSTOV, K.D.

[Determination of the effectiveness of observation and the  
number of events] Opredelenie effektivnosti nabliudeni i  
chisla sobytii. Dubna, Ob"edinennyi in-t iadernykh issle-  
dovanii, 1962. 9 p. (MIRA 15:2)  
(Photography, Particle track)

SOKOLOV, S.N.; TOLSTOV, K.D.; SARANTSEVA, V.R., tekhn. red.

[Verification of counting efficiency and estimating the true  
number of events] Kontrol' effektivnosti nabliudenii i otsenka  
istinnogo chisla sobytii. Dubna, Ob"edinennyi in-t iadernykh  
issledovaniii, 1962. 10 p. (MIRA 15:12)  
(Nuclear counters) (Mathematical statistics)

TOLSTOV, V.N.; TOLSTOV, K.N.

Orientation method for the determination of blood sugar. Zdrav.  
Kazakh. 22 no.10:75-76 '62. (MIRA 17:5)

1. Iz Ural'skoy oblastnoy bol'nitsy.

KOROLEV, P.A.; TOLSTOV, K.S.

Smoke generator of a multiple effect. Zashch. rast. ot vred. i bol.  
6 no.12:9-11 D '61. (MIRA 16:5)

LUCHINSKIY, G.P., doktor khimicheskikh nauk; TOLSTOV, K.S., kand.tehnicheskikh nauk; KOROLEV, P.A.

Use of hexachloran in insecticidal fumigants. Khim.prom. no.4:235-  
237 Ap '61. (MIRA 14:4)

(Cyclohexane) (Insecticides)

TOLSTOV, L.A.

Effect of temperature on the oil yield in water flooding.  
Neft. khoz. 43 no.6:38-42 Je '65. (MIRA 18:7)

NOVIKOV, V.N.; TOLSTOV, L.K.; SEREBRYAKOVA, Ye.K.; SOKOLOV, B.M.; Prinimal uchastiye: Melent'yev, Yu.I.; KAPGER, V.S.; ZORCHENKO, I.P.; KARPOV, K.F.; Kushnarenko, V.S.; SHEVCHENKO, L.I.; TRIFONOVA, N.I.; PODZHUNAS, V.A.; MASLITSKAYA, M.P.

Obtaining industrial naphthalene from the centrifugal naphthalene of the Gubakha Coke and Coal Chemicals Plant. Koks i khim. no.8: 35-38 '62.  
(MIRA 17:2)

1. Vostochnyy uglekhimicheskiy institut (for Novikov, Tolstov, Serebryakova). 2. Gubakhinskiy koksokhimicheskiy zavod (for Sokolov).

ZOLOTAVIN, V.L.; BUKREYEV, Yu.F.; TOLSTOV, L.K.; BEZRUKOV, I.Ya.

Photometric determination of sodium in pure vanadium pentoxide.  
Zhur. prikl. spektr. 2 no.5:461-462 My '65. (MIRA 18:7)

TOLSTOV L. M.

238T17

USSR/Chemistry - Viscosimeters Aug 52

"The Sliding of Mercury on Glass," L. M. Tolstov,  
Moscow Mach Tool and Tool Inst imeni I. V. Stalin

"DAN SSSR" Vol 85, No 6, pp 1329-1332

The sliding effect of Hg can be studied in a capillary having a diam about the size of a micron. Since this excludes the use of ordinary viscosimeters, a specially constructed instrument was made from a long-stem thermometer. Negative results obtained by Brillouin agree with those obtained by using this instrument. Presented by Acad P. A. Rebinder  
14 Jun 52

238T17

SIL'CHENKO, Serafim Semenovich; TOLSTOV, M.A., inzh., retsenzent;  
LIOZNYANSKIY, M.I., inzh., red.; YERMAKOV, N.P., tekhn.red.

[Hydraulic equipment of metal cutting machines; manual for  
repairman] Gidravlicheskoе oborudovanie metallorezhushchikh  
stankov; posobie dlia slesarei-remontnikov. Izd.2., dop.  
Moskva, Gos.nauchno-tekhn.izd-vo mashinostroit.lit-ry, 1958.  
169 p.

(MIRA 12:5)

(Machine tools--Hydraulic driving)

TOLSTOV, MIKHAIL ALEKSEYEVICH

N/5  
741.43  
.T6

PNEUMATICHESKIYE I PNEVMOIDRAVLICHESKIYE PRISPOSOBLENIYA / PNEUMATIC  
AND PNEUMOHYDRAULIC DEVICES/ MOSKVA, MASHGIZ, 1953-

V. DIAGRS. INCLUDES BIBLIOGRAPHIES. LIB.HAS: 1953 (1st ED.)  
1956 (2D ED.)

TOLSTOV, M.A.; SHNEYDERMAN, K.A., red.

[Footprint on the earth] Sled na Zemle. Rostov-na-Donu,  
Rostovskoe knizhnoe izd-vo, 1965. 138 p. (MIRA 18:8)

MEN'SHAKOV, Vladimir Mikhailovich; STUDENOK, G.A., inzhener, redaktor;  
KUVSHINSKIY, V.V., kandidat tekhnicheskikh nauk, retsenzent;  
~~TOLSTOV, M.A.~~, inzhener, retsenzent; DUGINA, N.A., tekhnicheskiy  
redaktor

[Planing machines] Strogal'nye stanki. Pod red. G.A. Studenok.  
Moskva, Gos. nauchno-tekhn. izd-vo mashinostroit. lit-ry, 1956.  
54 p. (Nauchno-populiarnaia biblioteka rabochego stanochnika,  
no. 22)

(Planing machines)

TOLSTOV, Mikhail Alekseyevich; KERNER, D.B., inzhener, retsenzent; KUVSHINSKIY, V.V., kandidat tekhnicheskikh nauk, redaktor; DUGINA, N.A., tekhnicheskiy redaktor  
[Pneumatic and pneumatic-hydraulic attachments] Pnevmaticheskie i pnevmogidravlicheeskie prispособleniya. Izd. 2-oe, dop. i perer. Moskva, Gos. nauchno-tekh. izd-vo mashinostroit. lit-ry, 1956. 202 p.  
(Machine tools--Attachments) (MIRA 10:4)

DOROFEEV, Vasiliy Ivanovich, zootehnik; TOLSTOV, M.A., red.

[We raise ducks without bodies of water] Vyrashchivaem  
utok bez vodoemov. Rostov-na-Donu, Rostovskoe knizhnoe  
izd-vo, 1963. 14 p. (MIRA 17:9)

1. Zernogradskaya gosudarstvennaya selektsionnaya stantsiya  
(for Dorofeyev).

SHKURIN, Gennadiy Alekseyevich; TOLSTOV, M. S., red.

[State dairy farm for Rostov workers] Molochnyi sovkhоз -  
trudashchimsia Rostova. Rostov-na-Donu, Rostovskoe knizh-  
noe izd-vo, 1963. 20 p. (MIRA 17:10)

1. Direktor molochnogo sovkhosa "Aksayskiy" Azovskogo pro-  
izvodstvennogo upravleniya Rostovskoy oblasti (for Shkuri).

OGANESYAN, Yefrem Avetisovi , deputat Verkhovnogo Soveta SSSR;  
TOLSTOV,M.A.,red.

[Creators of golden fleece] Tsvetny zolotogo runa. Rostov-na-Donu, Rostovskoe knizhnoe izd-vo, 1963. 35 p.

(MIRA 17:10)

1. Predsedatel' kolkhoza imeni 17 partkonferentsii,  
Remontnenskogo proizvodstvennogo upravleniya, Rostovskoy  
oblasti (for Oganesyan).

CHEPEL', Yakov Grigor'yevich; TOLSTOV, M.A., red.

[Let's discover, let's utilize the resources] Nakhodim,  
ispol'zuem rezervy. Rostov-na-Donu, Rostovskoe knizhnoe  
izd-vo, 1963. 28 p. (MIRA 17:10)

1. Direktor ordena Lenina sovkhoza "TSelinskiy" Yegorlykskogo  
proizvodstvennogo upravleniya, Rostovskoy oblasti (for Chepel').

GENIKA, Lyubov' Vasil'yevna, kand. veter. nauk; TCLSTOV, M.A.,  
red.

[Trichomoniasis in cattle] Trichomonoz krupnogo rogatogo  
skota. Rostov-na-Donu, Rostovskoe knizhnoe izd-vo, 1963.  
26 p. (MIRA 17:6)

FEDOROV, Boris Fedorovich; SLOBODYANNIKOV, S.S., kand.tekhn.nauk,  
retsenzent; TOLSTOV, M.A., inzh., retsenzent; BOGOSLAVETS,  
N.P., tekhn.red.

[Mechanization and automation of fitting and assembling  
operations] Mekhanizatsiia i avtomatizatsiia slesarno-  
sborochnykh rabot. Moskva, Mashgiz, 1962. 310 p.

(Assembly-line methods) (Machine-shop practice)  
(Automation) (MIRA 15:5)

ROZIN, Aleksandr Iosifovich; REVENKO, Vyacheslav Andreyevich;  
TOLSTOV, M.A., inzh., red.; KUTENKOVA, G.M., tekhn.red.

[Using straight and reverse tooth generation in the manufacture  
of cutting tools] Metod priamoi i obratnoi obkstki v proiz-  
vodstve instrumenta. Sverdlovsk, Tsentr.biuro tekhn.informatsii,  
1959. 40 p.

(Metal cutting)

(MIRA 14:4)

TOLSTOV, M. A.

LOSKUTOV, V.V.; BREYEV, B.G., kand.tekhn.nauk, retsenzent; KITAYEV,  
V.I., inzh., retsenzent; TOLSTOV, M.A., inzh., red.; MODEL',  
B.I., tekhn.red.

[Automatic and semiautomatic grinding machines] Shlifoval'nye  
avtomaty i poluvavtomaty. Moskva, Gos.sachno-tekhn.izd-vo  
mashinostroit.lit-ry, 1959. 292 p.  
(Grinding machines) (MIRA 13:3)

TOLSTOV, M.A.

Pnevmaticheskie i pnevmogidravliche-  
skie prisposobleniia (Pneumatic and pneumohydraulic  
devices). Sverdlovsk, Mashgiz, 1953. 164 p.

SO: Monthly List of Russian Accessions, Vol. 7, No. 5, August 1954

TOLSTOV M.A.

ALEKSEYEV, B.A.; ROZIN, A.I.; KLIMOV, V.I., inzhener, retsenzent; TOLSTOV, M.A.,  
inzhener, retsenzent; SOMOVA, T.M., inzhener, vedushchiy redaktor, re-  
daktor literatury po kholodnoy obrabotke metallov.

[Metal cutting tools; design and manufacture] Instrumental'noe delo.  
Moskva, Gos.nauchno-tekhn.izd-vo mashinostroit.lit-ry, 1952. 319 p.  
[Microfilm] (MLRA 7:10)

1. Uralo-Sibirskoye otdeleniye Mashgiza (for Somova).  
(Cutting tools)

TOLSTOV, M. A.

Pnevmaticheskiye i pnevmogidravlicheskiye prispособleniya (Pneumatic and pneumo-hydraulic devices) Moskva, Mashgiz, 1953. 162 p. diagrs. "Literatura i istochniki": p. 161.

SO: N/5  
741.43  
•T6

TEL. 51809 102-821

KUVSHINSKIY, Vladimir Vladimirovich; TOLSTOV, M.A., retaenzent; DUGINA,  
N.A. tekhnicheskiy redakter.

[Milling machinery] Frezernye stanki. Moskva, Gos.nauchno-tekhn.  
izd-vo mashinostroitel'noi lit-ry, 1955. 62 p. (Nauchno-popular-  
naia biblioteka rabochego stanochnika, no.24) (MLRA 9:1)  
(Milling machines)

SHUMAKOV, B.A., akademik, red.; TOLSTOV, M.A., red.

[Work mechanization in irrigation farming] Mekhaniza-  
tsiya rabot v oroshaemom zemledelii. Rostov-na-Donu,  
Rostovskoe knizhnoe izd-vo, 1965. 152 p.

(MIRA 19:1)

1. Vsesoyuznaya akademiya sel'skokhozyaystvennykh nauk  
imeni V.I.Lenina (for Shumakov).

TOLSTOV, Mikhail Alekseyevich; DUGINA, N.A., tekhn. red.

[Pneumatic and hydraulic-pneumatic devices] Pnevmaticheskie i  
pnevmogidravlicheskie prispособlenia. Izd.3., dop. i perer.  
Moskva, Gos. nauchno-tekhn. izd-vo mashinostroit. lit-ry,  
1961. 270 p. (MIRA 14:9)  
(Hydraulic machinery) (Pneumatic machinery)

TOlstov, M.A.; Murav'ev, K.N., inzhener, retsenzent; Kuvshinskiy, V.V.,  
kandidat tekhnicheskikh nauk, redaktor.

[Pneumatic and pneumohydraulic devices] Pnevmaticheskie i pnevmogidrav-  
licheskie prispособleniya. Sverdlovsk, Gos. nauchno-tekhn. izd-vo mashino-  
stroit. i sudostroit. lit-ry [Uralo-Sibirske otd-nie] 1953. 162 p.  
(MLRA 7:6)

(Pneumatic tools)

TOLSTOY, Mikhail Ivanovich; MIRONETS, Ye.M., red.

[Some problems in the evaluation of the metal potential  
in geological formations and revealing of endogenetic  
geochemical halos] Nekotorye voprosy otsenki metallo-  
nosnosti geologicheskikh obrazovanii i obnaruzheniya  
endogennykh geokhimicheskikh oreolov. Kiev, Izd-vo  
Kievskogo univ., 1964. 150 p. (MIRA 18:7,

KARISS, Ya.E.; TOlstoy, M.N.; FEDOFLOV, P.P.

Luminescence and absorption of trivalent neodymium in fluorite  
type crystals. Opt. i spektr. 18 no.3:440-445 Mr '65.  
(MIRA 18:5)

SEYFUL'-MULYUKOV, R.B., kand.geol.-mineral.nauk; TOLSTOY, M.P., prof.,  
doktor geol.-mineral.nauk, red.

[Genetic types of Quaternary sediments and basic data on geo-  
morphology] Geneticheskie tipy chetvertichnykh otlozhenii s  
osnovnymi svodeniiami po geomorfologii; kratkoe uchebnoe posobie.  
Pod red. M.P.Tolstogo. Moskva, Mosk.sel'khoz.akad.im. K.A.  
Timiriazeva, 1959. 79 p. (MIRA 13:9)  
(Sediments (Geology)) (Geology, Structural)

TOLSTOV, N.

Close contact with the economic council. Prof.-tekhn. obr. 15 no.4:  
3-4 Ap '58.  
(MIRA 11:5)

1. Zamestritel' nachal'nika Luganskogo oblastnogo upravleniya trudovykh  
rezervov.

(Lugansk Province--Technical education)

L 446-66 EWT(1)/EWT(m)/EWP(t)/EWP(b) IJP(c) JD  
ACCESSION NR: AP5017897 UR/0051/65/019/001/0097/0101  
535.377 48  
B

AUTHORS: Sokolov, V. A. Tolstoy, N. A. 44,55

TITLE: Thermal afterglow and thermostimulated current in TlCl single crystals 21,44,55

SOURCE: Optika i spektroskopiya, v. 19, no. 1, 1965, 97-101

TOPIC TAGS: luminescence, thallium compound, thermoluminescence, crystal lattice defect, optic transition

ABSTRACT: This is a continuation of earlier work (Sb. 'Fizika shchelochno galoidnykh kristallov. (II Vses. soveshch.) [Collection Physics of alkali halide crystals, Second All-Union Conference] p. 411, Riga, 1962), where it was shown that luminescence of thallium chloride undergoes strong temperature quenching when heated above 160C. To determine the level spectrum in the forbidden band and its variations in single crystals of thallium chlorides from different sources, which have different luminescence spectra, the authors in-

Card 1/4

L 4446-66

ACCESSION NR: AP5017897

O

vestigated the thermoluminescence and at the same time also the thermostimulated current (TSC) in the same single crystal specimens of TlCl. A block diagram of the experimental setup for the simultaneous measurement is shown in Fig. 1 of the Enclosure. The crystals were excited at -190°C with the 365-nm mercury line for five minutes, which was adequate for an equilibrium population of the capture levels. All the investigated single-crystal TlCl specimens displayed thermal afterglow (TA) and TSC. Specimens from different lots had similar shapes of TA and TSC curves, but different luminescence spectra. Different specimens cut from the same crystal (with similar luminescence spectra) may have different ratios of the peak maxima of the TA and TSC curves. The peaks of the TA and TSC of all the TlCl specimens lie within narrow temperature ranges. The net result is that the capture levels of the carriers in single TlCl crystals are connected with the intrinsic defects of the crystal lattice, and that the luminescence mechanism of this sort, when excited by the band-band transition, is similar to the Schoen-Klasens mechanism. Orig. art. has: 5 figures and 1 table.

Card 2/4

L 446-66

ACCESSION NR: AP5017897

0

ASSOCIATION: None

SUBMITTED: 28Apr64

ENCL: 01

SUB CODE: OP, SS

NR REF SOV: 013

OTHER: 010

Card 3/4

L 4446-66

ACCESSION NR: AP5017897

ENCLOSURE: 01




Fig. 1. Block diagram of setup for simultaneous measurement of thermoluminescence and thermostimulated current.

T - Thermocouple, O - direction of exciting light, II - oven of sample holder,  
 R - 15 Meg resistor, 0.6μF - sample, Φ3V - photomultiplier, M-95 - millivoltmeter,  
 M-95 - microammeter

*Deb*  
 Card 4/4

ACCESSION NR: AP4032872

S/0051/64/016/004/0677/0683

AUTHOR: Tolstoy, N.A.; Yepifanov, M.V.

TITLE: Shape of light pulses from a flash tube

SOURCE: Optika i spektroskopiya, v.16, no.4, 1964, 677-683

TOPIC TAGS: flash lamp, flash tube pulse, light impulse, flash tube pulse cutoff, taumeter, taumeter circuitry

ABSTRACT: Flash tubes are now successfully employed in conjunction with pulse taumeters; the use of the flash-pulse technique has made it feasible to advance from measurement of integral luminescence lifetimes to measurement of the duration of individual relaxation processes. However, the pulse taumeter technique, as compared with the ordinary taumeter procedure, has two related shortcomings which stem from the fact that the exciting pulses are not square and the fact that the trailing edge of the light pulse is longer than the leading edge. The last reduces the time-sensitivity of the pulse taumeter measurements. Accordingly, the authors propose a method for enhancing the time sensitivity by shortening the trailing edge of the light pulses from the flash tube. This is based on the assumption that the rise and fall

Card 1/2

ACCESSION: NR:AP4032872

are both exponential, and is realized in practice by modification of the flash tube supply (discharge) circuit to provide for sharp cutoff of the current to the tube. The circuit used by the authors with an IFK-120 infrared flash tube is diagramed. The shapes of the light pulses yielded by the tube under different supply conditions are shown in figures. An analytic expression for the pulse shape is also adduced. Orig.art.has: 7 formulas, 4 figures and 1 table.

ASSOCIATION: none

SUBMITTED: 25Jan63

DATE ACQ: 07May64

ENCL: 00

SUB CODE: OP, EE

NR REF Sov: 004

OTHER: 000

Card 2/2

RYSKIN, A.I.; TKACHUK, A.M.; TOLSTOY, N.A.

Properties of ethanolamine compounds of divalent platinum.  
Opt. i spektr. 18 no.3:422-431 Mr '65.  
(MIRA 18:5)

RYSKIN, A.I.; TKACHUK, A.M.; TOLSTOY, N.A.

Optical properties of complex compounds of bivalent platinum.

Izv. AN SSSR. Ser.fiz. 29 no.3:512-515 Mr '65.

(MIRA 18:4)

"APPROVED FOR RELEASE: 07/16/2001

CIA-RDP86-00513R001756120013-5

APPROVED FOR RELEASE: 07/16/2001

CIA-RDP86-00513R001756120013-5"

BABUSHKIN, L.N., doktor geogr. nauk, prof., glav. red.; AKRAMOV,  
Z.M., doktor geogr. nauk, red.; SULTANOV, G.S., kand.  
biol. nauk, red.; PETROSYANTS, M.A., kand. fiz.-matem.  
nauk, red.; ZARIFOV, Kh.T., kand. filolog. nauk, red.;  
TOLSTOV, N.N., red.; BAUDINA, S.B., red.; VOLKOVA-VOLK,  
V.M., red.

[Atlas of the Uzbek Soviet Socialist Republic] Atlas  
Uzbekskoi Sovetskoi Sotsialisticheskoi Respublikи.  
Tashkent, 1963. 53 p. (MIRA 18:2)

1. Russia (1923- U.S.S.R.) Glavnaya upravleniye geodezii  
i kartografii. 2. Tashkentskiy Gosudarstvennyy universitet  
(for Babushkin).

LEBEDEV, N.A.; TOLSTOY, N.S.; KHALKIN, V.A.

Microchromatographic column with remote control. Radiokhimiya 7  
no.1:115-117 '65. (MirA 18:6)

SAVENKO, Yuriy Filippovich; TOLSTOV, Nikolay Timofeyevich; VOSTORV, I.D.,  
otv.red.; SMIRENSKIY, M.M., red. izd-va; LOMILINA, L.N., tekhn. red.

[Stoper] Gornorabochii ochistnogo zaboia. Moskva, Gos.nauchno-  
tekhn.izd-vo lit-ry po gornomu delu, 1961. 183 p. (MIRA 14:12)  
(Stoping (Mining))

TOLSTOV, R.

Moon is made on earth. Znan.-sila 37 no.9:21 S '62.  
(MIRA 15:12)  
(Moon--Surface)

MASHKARENKO, A., inzhener-podpolkovnik; TOLSTOV, S., inzhener-podpolkovnik;  
KUNIN, V., inzhener-polkovnik; NETYKSA, V., podpolkovnik

Evacuation of tracklaying vehicles. Tekh. i vooruzh. no.6:  
(MIRA 17:7)  
46-49 Je'64

I. Tolstov, Sergei Pavlovich, ed.

Transactions of the Khorazmsk Archaeological and Ethnographic Expedition, 1937-  
DK861.K45

1. Khiva--Antiq. 2. Ethnology--Asia, Central. I. Tolstov, Sergei Pavlovich, ed.

TOLSTOV, S. P., Prof.

Prof. of the Moscow State University (1949)

"Ancient Khorezm," 1948

Stalin 1st Prize, 1948, publ.

Current Digest of the Soviet Press, Vol. 1, No. 15, 1949, page 17. (In █ Library)

TOLSTOV, S. P.

Ethnology

Crisis in bourgeois ethnography. Trudy Inst. etn. AM SSSR., No. 12, 1951

9. Monthly List of Russian Accessions, Library of Congress, March 1953, Unclassified.

TOLSTOV S. P.

Turkmenistan - Antiquities

Archaeological surveys along the route of the Main Turkmen Canal (Uzboy river bed).  
Krat. soob. Inst. etn. AN SSSR 14, 1952.

Monthly List of Russian Accessions, Library of Congress, September 1952. UNCLASSIFIED

TOLSTOV, PROF. S. F.

Antiquities-Turkmenistan

Archaeological remains on the route of the Main Turkmen Canal. Prof. S. P. Tolstov.  
Vest. AN SSSR 22, No. 4, 1952.

9. Monthly List of Russian Accessions, Library of Congress, September 1953<sup>2</sup> Uncl.

TOLSTOV, S.P.

N.N. Miklukho-Maklai; 65th anniversary of his death. Sov. etn. no.2:3-9  
153. (MIRA 6:6)  
(Miklukho-Maklai, Nikolai Nikolaevich, 1846-1888)

TULSTOV, S.P., doktor istoricheskikh nauk.

N.N. Miklukha-Maklai (on the occasion of the 65th anniversary of his death).  
Nauka i zhizn' 20 no.4:33-35 Ap '53. (MLRA 6:5)  
(Miklukha-Maklai, Nikolai Nikolaevich, 1846-1888)

TOLSTOV, S.P.

Sredneaziatskii ethnograficheskii sbornik  
(Collection of articles on Central Asian ethnography)  
Moskva, Izd. Ak. nauk SSSR, 1954. 412 p.

SO: Monthly List of Russian Accessions, Vol. 7, No. 5, August 1954

**USSR/ Geography - Geology**

**Card 1/1 Pub. 45 - 5/16**

**Authors** : Tolstov, S. P.; Kes', A. S.; and Zhdanko, T. A.

**Title** : The history of the Sarikamish Lake in the Middle Ages

**Periodical** : Izv. AN SSSR. ser. geog. 1, 41-50, Jan-Feb 1954

**Abstract** : The origin of Lake Sarikamish is traced to the Pliocene Epoch at which time it was full of water and formed a large basin. During the first half of the Quaternary Period it became dry and in the second half of the same period it again filled with water due to the change in the course of the Amu-Darya River. In the 16th Century the level of the water began to sink, the water became salty and it finally dried out altogether. Fifteen Russian and USSR references (1879-1953). Maps; drawings.

**Institution** : Ethnographical and Geographical Institute of the Soviet Academy of Science

**Submitted** : ...

TOlstov, S.

Mikhail Mikhailovich D' iakovov; obituary. Sov. etn. no.3:122-124  
'54. (MLRA 7:11)  
(D' iakovov, Mikhail Mikhailovich, d. 1954?)

GULYAMOV, Yakh'ya Gulyamovich, prof.; TOLSTOV, S.P., otvetstvennyy red.;  
IMVINA, A.A., red.izd-va; SHAPEL'KOV, A.T., tekhn.red.

[History of irrigation in Khorezm from ancient times to our day]  
Istoriia oroshenija Khorezma s drevneishikh vremen do nashikh  
dnej. Tashkent, Izd-vo Akad.nauk Uzbekskoi SSR, 1957. 313 p.

1. Pochetnyy akademik AN UzSSR, chlen-korrespondent AN SSSR (for  
Tolstov)  
(Khorezm Province--Irrigation)

70/5104, Ser.  
AL-BIRUNI; ABDULLAYEV, Kh.M., akademik, red.; AZIMDZHANOVA, S.A., kand. istor.nauk, red.; BELENITSKIY, A.M., kand. istorich.nauk, red.; BELYAYEV, V.I., kand.filologicheskikh nauk, red.; GULYAMOV, Ya.G., red.; KARY-NIYAZOV, T.N., akademik, red.; LEMMLEYH, G.G., prof., red.; SAL'YE, M.A., kand.filolog.nauk, red.; SEMENOV, A.A., red.; TOlstov, S.P., pochetnyy akademik, red.; UKLONSKIY, A.S., akademik, red.; LIUBECHANSKAYA, N.I., red.; GOR'KOVAYA, Z.P., tekhn.red.

[Selected works] Izbrannye proizvedeniia. Tashkent, Izd-vo Akad.nauk Uzbekskoi SSR. Vol.1. 1957. 485 p. (MIRA 11:1)

1. AN UzSSR (for Abdullayev, Kary-Niyazov, Tolstov, Uklonskiy).
2. Chlen-korrespondent AN UzSSR (for Gulyamov, Semenov).  
(Science, Medieval)

SHCHERBAKOV, D.I., akademik; FRUMKIN, A.N., akademik; KHACHATUROV, T.S.;  
VINOGRADOV, A.P., akademik; SOBOLEV, S.L., akademik; KOSTENKO, M.P.,  
akademik; TOLSTOV, S.P.; SAZHIN, N.P.; KAZARNOVSKIY, I.A.; VUL, B.M.;  
TROFIMUK, A.A., akademik

Discussion of the annual report. Vest. AN SSSR 33 no.3:25-34  
Mr '63. (MIRA 16:3)

1. Chleny-korrespondenty AN SSSR (for Khachaturov, Tolstov, Sazhin,  
Kazarnovskiy, Vul). (Academy of Sciences of the U.S.S.R.)

TOLSTOV, S.P.

Ancient irrigation system and outlook for modern irrigation; based  
on investigations in the ancient Syr-Darya Delta. Vest. AN SSSR  
31 no.11:59-65 N '61. (MIRA 14:11)

1. Chlen-korrespondent AN SSSR.  
(Aral Sea region--Irrigation)

TOLSTOV, S.P.; KES', A.S., kand.geograf.nauk; ITINA, M.A., kand.istor.  
nauk; ANDRIANOV, B.V., kand.istor.nauk; ZHDANKO, T.A., kand.  
istor.nauk; VISHNEVSKAYA, O.A., nauchnyy sotrudnik; VAKTURSKAYA,  
N.N., kand.istor.nauk. Prinimeli uchastiye LEVINA, L.M.,  
aspirantka; TRUDNOVSKAYA, S.A.; DAVIDOVICH, Ye.A., kand.istor.  
nauk; ANDRIANOV, B.V., red.izd-va; LEBEDEVA, L.A., tekhn.red.

[The lower reaches of the Amu Darya, the Sarykamysh and the Uzboy;  
history of their formation and settlement] Nizov'ia Amu-Dar'i,  
Sarykamysh, Uzboi; istoriia formirovaniia i zасeleniia. Pod  
obshchei red. S.P.Tolstova. Moskva, 1960. 346 p. (Materialy  
Khorezmskoi ekspeditsii, no.3). (MIRA 14:2)

1. Akademiya nauk SSSR. Institut etnografii. 2. Chlen-korrespon-  
dent AN SSSR (for Tolstov). 3. Institut etnografii AN SSSR (for  
Levina). 4. Akademiya nauk Tadzhikskoy SSR (for Davidovich).  
(Amu Darya Valley)