

VAGNER, V.V.

Restrictive semigroups. *Izv.vys.ucheb.zav.*; mat. no.6:19-27 '62.
(MIRA 15:12)

1. Saratovskiy gosudarstvennyy universitet imeni N.G.
Chernyshevskogo.

(Groups, Theory of)

VAGNER, V. V.

"Generalized groups of particle transformations"

report submitted at the Intl Conf of Mathematics, Stockholm, Sweden,
15-22 Aug 62

VAGNER, V.V.; LOPSHITS, A.M.

Iakov Semenovich Dubnov; obituary. Trudy Sem.po vekt.i tenz.anal.
no.11:3-17 '61. (MIRA 15:3)
(Dubnov, Iakov Semenovich, 1887-1957)

S/044/62/000/004/007/099
C111/C444

AUTHOR: Vagner, V. V.
 TITLE: Variation theory as field theory of the central semi-cones
 PERIODICAL: Referativnyy zhurnal, Matematika, no. 4, 1962, 69,
 abstract 4A422. ("Nauchn. yezhegodnik. Saratovsk. un-t.
 Mekhan.-matem. fak., 1955". Saratov, 1959, 27-34)
 TEXT: Let in X_n a differential m -pseudometric be given, i. e.
 let a unique binary relation $\rho \subset E_n(X_n) \times R$ between the points of the
 fibred space $E_n(X_n)$ and the real numbers be given to which there
 corresponds in the fibred space $E_n \times R(X_n)$ which is in one-to-one corres-
 pondence to the space $E_n(X_n) \times R$ in a natural way, the central semi-conic
 m -secant surface $K_{(n+1)+m}$ i. e. an m -secant surface for which the inter-
 sections $K_{(n+1)+m} \cap E_n \times R(p)$ for all points $p \in \rho(K_{(n+1)+m})$ are the
 central m -semicones K_m , where ρ is the canonical mapping of $E_n \times R(X_n)$
 on X_n . Let
 Card 1/4

Variation theory as field theory ...

S/044/62/000/004/007/099
C111/C444

$$x^\lambda = L^\lambda(\xi^\lambda, \eta^a), \quad x = L(\xi^\lambda, \eta^a)$$

be the coordinate equations of $K_{(n+1)+m}$. The first n equations define the central semiconic m -secant surface $pr_1 \mathcal{S}$ in $E_n(X_n)$, while the function L gives an expression for the pseudomodulus of the measurable vector $x \in pr_1 K_m$ in the tangential $E_n(p(\xi^\lambda))$ in its homogeneous curvilinear coordinates η^a .

An oriented curve C in X_n is called measurable with respect to the given m -pseudometric \mathcal{S} , if its positive tangential semi straight lines belong to $pr_1 \mathcal{S}$. If the measurable curve C is defined by the equations

$$\xi^\lambda = \xi^\lambda(t) \quad (t_1 \leq t \leq t_2) \quad (1)$$

and if $\eta^a = \eta^a(t)$ is the field of its contact vectors, then the scalar

Card 2/4

S/044/62/000/004/007/099
C111/C444

Variation theory as field theory ...

$$\int_{t_1}^{t_2} L(\dot{x}(t), x(t)) dt$$

is an integral invariant which is called a pseudoarc-length. To every measurable curve C in X_n with the equations (1) in the $(n+1)$ space $X_n + R$ (Cartesian product of X_n and the numerical line R) one makes correspond a curve \hat{C} , having the equations


$$\hat{C} = \left\{ \begin{aligned} \dot{x} &= \dot{x}(t), \\ \dot{t} &= \sqrt{L(\dot{x}(t), x(t))} \end{aligned} \right. dt$$

If \hat{C} is anormal, then C is geodetic (in this article the notion of an anormal curve is defined by aid of the there introduced binary relation of the variation of a curve).

Card 3/4

Variation theory as field theory ...

S/044/62/000/004/007/099
C111/C444

It is proved: In order a measurable curve to possess a weakly extremal pseudoarc-length, it is necessary that it be geodetic. It is stated that the variation problem of Lagrange on the conditional extremum may be also considered as the problem of the determination of curves with extremal pseudolengths in X_m in a differential m -pseudometric. 

[Abstracter's note: Complete translation.]

Card 4/4

VAGNER, V.V.

Transformative semigroups. Izv. vys. ucheb. zav.; mat. no.4:36-48
'60. (MIRA 13:10)

1. Saratovskiy gosudarstvennyy universitet im. N.G. Chernyshevskogo.
(Groups, Theory of)

85197

14.2000

S/140/60/000/004/010/023 XX
C111/C222AUTHOR: Vagner, V.V.TITLE: Transformative Semigroups ¹⁶PERIODICAL: Izvestiya vysshikh uchebnykh zavedeniy. Matematika, 1960,
No. 4, pp. 36 - 48

TEXT: The author uses notations and notions of (Ref. 1).
A semigroup is called transformative if in it, beside of the fundamental binary operation \circ which here is understood as a ternary relation between the elements of the semigroup, two binary relations are given: quasiorder χ and quasiequivalence ξ satisfying the condition that there exists an eigenrepresentation of the semigroup with the aid of partial transformations of a set, for which χ can be represented as an enclosure relation of the first projections of the partial transformations and ξ as a union relation of the partial transformations.

In the present paper the author proposes an axiomatic foundation of the theory of transformative semigroups, where these are understood as ordered systems (G, \circ, χ, ξ) where the set of the elements of the semigroup and \circ, χ, ξ are the mentioned relations. Eight theorems are given, the
Card 1/2

85497

Transformative Semigroups

S/140/60/000/004/010/023 XX
C111/C222

assertions of which are partially already contained in (Ref. 1) (theorems 2 and 5) or are known from the general theory (theorem 6). As an exceptional case the author considers the representation of semigroups with the aid of set transformations. It is pointed out that the axiomatic theory of representable ordered semigroups can be obtained immediately from the theory of transformative semigroups.

There is 1 Soviet reference.

[Abstracter's note : (Ref. 1) is a paper of V.V. Vagner in Matematicheskii sbornik, 1956, Vol. 38, pp. 203 - 240]

ASSOCIATION: Saratovskiy gosudarstvennyy universitet imeni N.G. Chernyshevskogo (Saratov State University imeni N.G. Chernyshevskiy)

SUBMITTED: January 28, 1960

Card 2/2

85943

S/039/60/052/001/005/009 XX
C111/C222

16.2000

AUTHOR: Vagner, V.V. (Saratov)

TITLE: Semigroups Associated With a Generalized Heap

PERIODICAL: Matematicheskiy sbornik, 1960, Vol. 52, No. 1, pp.597-628

TEXT: The author uses notations and notions of (Ref. 4,5).
 The author considers the binary operative (G, o) , i.e. the set G together with the ternary relation $o \subset G \times G \times G$ which determines a binary algebraic operation defined everywhere. Let in brief

$$(1) \quad o(g_1, g_2) = g_1 g_2 \quad \checkmark$$

The following dual relations are defined:

1) right and left ideal relations τ_d and τ_s :

$$(3) \quad \tau_d = \bigcup_{(g_1, g_2)} \bigvee_g (g_1 g = g_2) \quad , \quad \tau_s = \bigcup_{(g_1, g_2)} \bigvee_g (g g_1 = g_2)$$

2) right and left canonical equivalence relations ϵ_d, ϵ_s :

Card 1/8

85943

Semigroups Associated With a Generalized Heap S/039/60/052/001/005/009 XX
C111/C222

$$(5) \quad \epsilon_d = \bigcap_{(g_1, g_2)} \bigwedge_g (gg_1 = gg_2) \quad , \quad \epsilon_s = \bigcap_{(g_1, g_2)} \bigwedge_g (g_1g = g_2g)$$

3) right and left absorption relations α_d, α_s :

$$(6) \quad \alpha_d = \bigcap_{(g_1, g_2)} (g_1g_2 = g_1) \quad , \quad \alpha_s = \bigcap_{g_1, g_2} (g_2g_1 = g_1)$$

The absorption relation is denoted with

$$(7) \quad \alpha = \alpha_d \cap \alpha_s = \bigcap_{(g_1, g_2)} (g_1g_2 = g_2g_1 = g_1)$$

The invariant, symmetric, reflective, binary relation being dual to itself

$$(8) \quad \zeta = \bigcap_{(g_1, g_2)} (g_1g_2 = g_2g_1) = \text{pr}_{12} (o \cap c^*)$$

is called the commutativity relation, where o^* is defined by

Card 2/8

85943

S/039/60/052/001/005/009 XX
C111/C222

Semigroups Associated With a Generalized
Heap

$$(4) \quad o^*(g_1, g_2) = o(g_2, g_1)$$

and $pr_{12} o = G \times G$.

Let the considered operative (G, o) be a semigroup, i.e. let o be associative. The factor semigroups of (G, o) with respect to stable ϵ_d (resp. ϵ_s) are called right (resp. left) canonical factor semigroups. | b

Theorem 1 : For every semigroup, α_d , α_s and α are transitive binary relations.

A semigroup is called idempotent if all elements are idempotent.

Theorem 2 : For an idempotent semigroup, α_d and α_s are quasiordering relations and α is an ordering relation.

Theorem 3 : For idempotent semigroups it holds

$$(15) \quad \tau_d = \bar{\alpha}_s, \quad \tau_s = \bar{\alpha}_d.$$

Card 3/8

X

85943

Semigroups Associated With a Generalized
Heap

S/039/60/052/001/005/009 XX
C111/C222

Theorem 4 : If \mathcal{E} is a stable equivalence relation between the elements of an idempotent semigroup, then the binary relation $\alpha_d \circ \mathcal{E}$ (resp. $\alpha_s \circ \mathcal{E}$) is the maximal canonical inverse image of the right (resp. left) relation of absorption of the factor semigroup of this semigroup with respect to \mathcal{E} . X

Theorem 5 , Conclusion :

A semigroup is called pseudocommutative from the left (right) if its left (right) canonical factor semigroup is commutative.

Theorem 6 : For an idempotent semigroup (G, \circ) pseudocommutative from the right (left) it holds

$$(18) \quad \varepsilon_d = \alpha_d \cap \alpha_d^{-1} \quad (\varepsilon_s = \alpha_s \cap \alpha_s^{-1})$$

Theorem 7 : For an idempotent semigroup pseudocommutative from the right (left) it holds

$$(21) \quad \alpha_d = \alpha_s \circ (\alpha_d \cap \alpha_d^{-1}) \quad (\alpha_s = \alpha_d \circ (\alpha_s \cap \alpha_s^{-1})) .$$

Theorems 8 and 9 are conclusions from the theorems 6 and 7 .

Card 4/ 8

85943

Semigroups Associated With a Generalized
Heap

S/039/60/052/001/005/009 XX
C111/C222

The fact that for idempotent semigroups the absorption relation is an ordering relation, permits to use the theory of ordered sets in the theory of idempotent semigroups. Let (G, ω) be an ordered set, i.e. a set G with the ordering relation ω . Let g_1 be a minorant of g_2 and g_2 be a majorant of g_1 if

$$(25) \quad g_1 \{ g_2 \leftrightarrow (g_1, g_2) \in \omega.$$

The subset \mathcal{M} is called minorized (majorized) if it has a minorant (majorant) i.e. if $\bar{\omega}^{-1}(\mathcal{M}) \neq \emptyset$ ($\bar{\omega}(\mathcal{M}) \neq \emptyset$). The relation $\omega \circ \bar{\omega}^{-1}$ ($\bar{\omega} \circ \omega$) is called minorizing (majorizing) relation. The subset \mathcal{M} is called a minorant (majorant) raster if each of its pairs of elements has a minorant (majorant) belonging to this subset.

Theorem 10 : In order that all maximal connected subsets of an ordered set (G, ω) are minorant (majorant) rasters it is necessary and sufficient that

$$(31) \quad \bar{\omega}^{-1} \circ \omega \subset \omega \circ \bar{\omega}^{-1} \quad (\text{resp.} \quad \bar{\omega} \circ \omega \subset \bar{\omega}^{-1} \circ \omega).$$

Card 5/8

X

85943

Semigroups Associated With a Generalized
Heap

S/039/60/052/001/005/009 XX
C111/C222

Now every idempotent semigroup (G, o) is considered as a set ordered with the aid of the absorption relation α . It is stated that the theory of commutative idempotent semigroups in essential is equivalent to the theory of minorant semilattices.

Theorem 12 : For every idempotent semigroup it holds

(38) $\xi \subset \alpha \circ \alpha^{-1}$.

Theorem 13 : For an idempotent semigroup pseudocommutative from the right (left) it holds

(39) $\alpha^{-1} \circ \alpha \subset \xi$.

Theorem 14 : For an idempotent semigroup pseudocommutative from the right (left) which is ordered with the aid of the absorption relation, all maximal connected subsets are minimal raster.

The author considers the ternary operative (K, o) , i.e. a set K with the quaternary relation $o \subset K \times K \times K \times K$ which explains an everywhere defined ternary algebraic operation. Let in brief

(41) $o(k_1, k_2, k_3) = [k_1 k_2 k_3]$

Card 6/8

UX

85943

Semigroups Associated With a Generalized Heap

5/039/60/052/001/005/009 XX
C111/C222

Three binary algebraic operations o_1, o_2, o_3 are associated with the ternary algebraic operation o and they are denoted with $\triangleright, \triangleleft, \nabla$:

$$(44) \quad o_1(k_1, k_2) = k_1 \triangleright k_2 = [k_1 k_1 k_2], \quad o_2(k_1, k_2) = k_1 \triangleleft k_2 = [k_1 k_2 k_2];$$
$$o_3(k_1, k_2) = k_1 \nabla k_2 = [k_1 k_2 k_1]$$

The three binary operatives $(K, o_1), (K, o_2), (K, o_3)$ are denoted as the first, second and third associative binary operative of the ternary operative (K, o) . The ternary operative (K, o) is called a semiheap if for arbitrary k_1, \dots, k_s it holds

$$(45) \quad \left[[k_1 k_2 k_3] k_4 k_5 \right] = \left[k_1 [k_4 k_3 k_2] k_5 \right] = \left[k_1 k_2 [k_3 k_4 k_5] \right].$$

The semiheap (K, o) is called a generalized heap if the operation o is idempotent and bicommutative.

Theorem 15 : For the generalized heap (K, o) the first (second) associated binary operative is the idempotent semigroup pseudocommutative from the Card 7/8

X

85943

Semigroups Associated With a Generalized
Heap

S/039/60/052/001/005/009 XX
C111/C222

left (right).

Theorem 16 : The absorption relations of the first and second associated semigroup of a generalized heap are identical.

After introducing numerous further definitions, the author proves further 12 theorems on the connection between a generalized heap and the semigroups associated with it. The proofs result in essential from the theorems 1 - 14. The theorems 25 and 26 are already contained in (Ref. 6).

There are 6 references : 3 Soviet, 2 French and 1 German.

[Abstracter's note : (Ref. 4) concerns a paper of V.V. Vagner in Matematicheskiy sbornik, 1953, Vol. 32, pp. 545-632, (Ref. 5) is a paper of V.V. Vagner in Ukrainskiy matematicheskiy zhurnal, 1956, Vol. 8, pp. 235-252; (Ref.6) is a paper of V.V. Vagner in Ukrainskiy matematicheskiy zhurnal, 1959, Vol. 11, pp. 231-242] .

SUBMITTED: January 14, 1959

Card 8/8

TARASLENKOVA, Ye.M.; ABALENKOVA, Ye.A.; MANKIN, P.A.; VAGNER, V.V.

Catalytic transformations of n-heptane over manganese oxides.
Trudy LIEI no. 46:96-102 '63. (MIRA 17:6)

BURMISTROV, Ye.F., dots., red.; VAGNER, V.V., prof., red.; LIBER,
A.Ye., prof., red.; FAL'KOVICH, S.V., prof., red.;
PERSHIN, A.I., st. преподаvatel', red.; PERSOVA, V.M., red.

[Work of young scientists; mathematics issue] Trudy molodykh
uchenykh; vypusk matematicheskii. Saratov, 1964. 121 p.

(MIRA 18:8)

1. Saratov. Universitet. 2. Kafedra matematiki i statistiki
Saratovskogo ekonomicheskogo instituta (for Pershin).

VAGNER, V.V.; GLUSKIN, L.M.; AYZENSHTAT, A.Ya.

Evgeni Sergeevich Liapin, 1914-; on his 50th birthday. Usp. mat.
nauk 20 no.1:244-245 Ja-F '65. (MIRA 18:4)

VAGNER, V.V. (Saratov)

Shifts in a groupoid. Izv.vys.ucheb.zav.; mat. no.6:
37-47 '65. (MIRA 19:1)

1. Submitted May 27, 1965.

VAGNER, V.V. (Saratov)

Groupoid theory. Izv.vys.ucheb.zav.; mat. no.5:31-42 '65.

(MIRA 18:16)

VAGNER, Ye.A.

Diagnosis and treatment of injuries of the pericardium and the
heart. Khirurgiia, no.9:70-71 S '55. (MLBA 9:2)

1. Iz khirurgicheskogo otdeleniya Beresnikovskoy gorodskoy bol'nitsy
(glavnyy khirurg goroda--kandidat meditsinskikh nauk A.P. Nõskov)

(HEART, wounds and inj.
diag. & ther.)

(PERICARDIUM, wounds and inj.
same)

(WOUNDS AND INJURIES
heart & pericardium, diag. & ther)

VAGNER, YE. A.

VAGNER, YE. A.: "Surgical tactics in penetrating wounds of the chest under peacetime conditions." Molotov State Medical Inst. Molotov, 1956. (Dissertation for the Degree of Candidate in Medical Science.)

So: Knizhnaya letopis', No. 37, 1956. Moscow.

VAGNER, Ye.A., kand.med.nauk (Berezniki, Permskoy oblasti, ul. Demeneva,
d.11, kv.17)

Surgical treatment of penetrating wounds of the chest in peacetime.
Vest.khir. 83 no.9:89-95 S '59. (MIRA 13:2)

1. Iz khirurgicheskogo otdeleniya Bereznikovskoy gorodskoy bol'nitsy
(glavnyy khirurg - kand.med.nauk A.P. Noskov) i kafedry obshchey
khirurgii (zaveduyushchiy - prof. N.M. Stepanov) Permskogo meditsin-
skogo instituta.
(THORAX, wds. & inj.)

VAGNER, Ye.A.; TSUKANOV, V.I.

Our experience in the treatment of concussions of the brain.
Khirurgiia 36 no.11:102-105 N '60. (MIRA 13:12)

1. Iz otdeleniya neotlozhnoy khirurgii i travmatologii (zav. -
kand.med.nauk Ye.A. Vagner) oblastnoy bol'nitsy g. Berezniki
Permskoy oblasti.

(BRAIN--CONCUSSION)

VAGNER, Yevgeniy Antonovich; DEKHTYAR', Ye.G., red.

[Surgical treatment of penetrating chest wounds in peace
time] Khirurgicheskoe lechenie pronikaiushchikh ranenii
grudi v mirnoe vremia. Moskva, Meditsina, 1964. 191 p.
(MIRA 18:2)

VAGNER, Ya.A., dotsent (g. berezniki); SPUNBERG, Ya.A. (g. berezniki)

Review of V.P. Smol'nikov's book "Problems of emergency anesthesia."
Vest. khir. 93 no.9-138-142 S. 194. (MIRA 19-6)

GOLOVIN, Igor' Filippovich; MIKHAYEV, A.V., redsentsent; ~~VAGNER-MEMCHINOV,~~
A.A., spetsredaktor; MOROZOVA, I.I., redaktor; GOLLIB, E.M.,
tehnicheskiyredaktor

[Whaling techniques] Tekhnika kitoboinogo promysla. [n.p.]
Rishnepromisdat [1956] 110 p. (MIRA 10:1)
(Whaling)

ANDREYEV, Nikolay Nikiforovich; ~~VAGNER-NEMCHINOV, A.A.~~, spets. red.;
KCSSOVA, O.N., red.; SCKOLOVA, I.A., tekhn. red.

[Manual on fishing implements, nets and equipment] Spravochnik
po orudiam lova, setesnastnym materialam i promyslovomu sna-
riazheniiu. Moskva, Pishchepromizdat, 1962. 503 p.
(MIRA 16:1)

(Fishing--Equipment and supplies)

PETROSYAN, Ye.A.; VAGNER-SAKHAROVA, M.P.; STARCHEUS, A.V.; VARGINA, A.K.

Production of sorbed diphtheria and tetanus anatoxin. Nauch. ozn.
proizv. bakt. prep. 10:107-117 '61. (MIRA 18:7)

1. Moskovskiy institut vaktsin i syvorotok im. Mechnikova.

PRAT, V.; HATAIA, M.; BEBESOVA, D.; Technicka asistence: DIVIS, Z.; DVORAKOVA,
J.; VAGNEROVA, E.; VIDMAROVA, H.

Experimental contribution to the problem of "nephropathogenity"
of some strains of E. coli. Cas. lek. cesk. 103 no.40:1097-1102
2 0 '64.

1. Ustav pro choroby obehu krevniho v Praze (reditel prof. dr.
J. Brod, DrSc.); Ustav klinicke a experimentalni chirurgie v
Praze (reditel prof. dr. B. Spacek, DrSc.) a Katedra patolo-
gicke anatomie a mikrobiologie fakulty detskeho lekarstvi Kar-
lovy University v Praze (vedouci prof. dr. V. Kubelka).

1. *Effect of chlorotetracycline on the growth of the bacterium
ex: rimental with antibiotic (chlorotetracycline) in a medium.*
no. 42:313-319 (1959) G. B. Johnson.

*Effect of chlorotetracycline on the growth of the bacterium
ex: rimental with antibiotic (chlorotetracycline) in a medium.*
no. 42:313-319 (1959) G. B. Johnson.

1. *Enteropneusts of the genus* (1959) G. B. Johnson.
and, D. G. Johnson; *Enteropneusts of the genus*
(1959) G. B. Johnson, D. G. Johnson, and G. B. Johnson.
in a medium with antibiotic (chlorotetracycline) in a medium.
G. B. Johnson, D. G. Johnson, and G. B. Johnson.

PRAT, V.; HEJL, Z.; DEJDAR, R. Technicka spoluprace: VAGNEROVA, E.; TROUSIL, V.

Our experiences with transfemoral aortography by the Seldinger
technic. Rozhl. chir. 43 no.12:812-817 D '64.

1. Ustav pro choroby obehu krevniho v Fraze (reditel prof. dr.
J. Brod, DrSc.)

HATAIA, H.; HAVL, H.; KLIK, O. Technická spolupráce: KNEŽOVÁ, K.;
VAGNEROVÁ, E.

The use of TTC - a test for determining the significance of
bacteriuria in pregnancy. Cas. lek. cesl. 104 no.8:221
26 F'65.

1. Ústav klinické a experimentální chirurgie v Praze (ředitel:
prof. dr. R. Spáček, DrSc.); Ústav pro choroby oběhu krve v
Praze (ředitel: prof. dr. J. Bredl, DrSc.); a Ženské oddělení Ústavu
narozeního zvlášť ONV v Praze 4, (ředitel: UNZ MUDr. Z. Vaclavík).

PRAT, V., (Praha-Krc, Budejovicka 800); HATALA, M.; BEER, O.; SRAMEK, L.;
VAGNEROVA, E.; Laboratorni spoluprace: KREJCOVA, K.; CHRPOVA, V.

Bacteriuria in pregnancy and its relation to the maturity of
infants. Cas. lek. Cesk. 104 no.48:1327-1328 3 D '65.

1. Ustav pro choroby obehu krevniho v Praze (reditel prof. dr.
J. Brod, DrSc), Ustav klinicke a experimentalni chirurgie v
Praze (reditel prof. dr. B. Spacek, DrSc) a Zenske oddeleni
Ustavu narodniho zdravi Okresniho narodniho vyboru v Praze 4,
(reditel MUDr. Z. Vaclavik).

HATALA, M.; PRAT, V.; ROSSMANN, P. Technicka asistence: DIVIS, Z.;
DVORAKOVA, J.; VAGNEROVA, E.; VIDMAROVA, H.

Pathogenicity of different species of various serotypes of
Escherichia coli in the kidneys and urinary tract of rabbits.
Rozhl. chir. 44 no.5:326-333 My'65.

1. Ustav klinicke a experimentalni chirurgie v Praze (reditel:
prof. dr. B. Spacek, DrSc.); a Ustav pro choroby obehu krevniho
v Praze (reditel: prof. dr. J. Brod. DrSc.).

VAGNEROVA, Kamila; MACURA, J.; CATSKA, Vlasta

Rhizosphere microflora of wheat. I. Composition and properties of bacterial flora during the first stages of wheat growth. II. Composition and properties of bacterial flora during the vegetation period of wheat. III. Fungal flora of wheat rizosphere. Folia microbiol 5 no.5:298-330 '60. (EEAI 10:4)

1. Department of Microbiology, Institute of Biology, Czechoslovak Academy of Sciences, Prague.

(Wheat) (Roots (Botany)) (Bacteria)
(Rhizosphere) (Fungi)

VAGNEROVA, Kamila; MANCURA, V.

Production and utilization of aminoacids by various species of rhizosphere bacteria. Folia microbiol. 7 no.1:55-60 '62.

1. Department of Soil Microbiology, Institute of Microbiology Czechoslovak Academy of Sciences, Prague 6.
(AMINO ACIDS metab) (BACTERIA metab)

VAGNEROVA, Kamila; VANCURA, Vlastimil; LASIK, Jaromir

Rhizosphere microflora of wheat. Pt. 4. Rost vyroba 9 no. 7/2:637-692
Jl-Ag '63.

1. Mikrobiologicky ustav, Ceskoslovenska akademie ved, oddeleni
pudni mikrobiologie, Praha.

VAGO, A.

Complications and reaction in blood transfusion. Orv. hetil., Budap.
93 no.3:74-79 20 Jan 52. (CIHL 21:5)

1. Doctor.

--- VA00, F.

45. The determination of the quantity of mullite in baked ceramic products -- *Mullit mennyiségének meghatározása égetett kerámiái termékekben* -- J. Grofcsik and E. Vágó, (Building Materials -- *Építőanyag* -- Vol. 4, 1952, No. 11-12, pp. 211-215, 5 figs., 6 tabs)

Finding the optimal conditions for the formation of mullite, such as baking time, composition of the raw material, catalysts, etc. is very important in the manufacture of both refractories and ceramics. A rapid and accurate method for determining the mullite content of the finished products is therefore necessary. If the material to be examined is finely pulverized with 15% ammonium chloride it can be moulded with Canada balsam into 0.4-0.5 mm dia sticks for Debye-Scherrer X ray test. X-ray photos are subsequently made of the mixture with $\text{CuK}\alpha$ rays, Ni filter, 32 kv and 15 ma, after which the intensity of the 1.50 Å and 0.935 Å lines can be determined photometrically. The intensity of the lines is proportional to the quantity of mullite present. This method yields very accurate results which are not influenced by the possible presence of sillimanite, β -corundum, or zirconium.

Hungarian Technical Abst.
Vol. 5 No. 4 1953

VAGC, E.

VAGC, E.; TARDY, P.

"Use of Dolomite in the production of fireproof materials", p. 116,
(MATEM. FIS, Vol. 6, No. 6, June 1954, Budapest, Hungary)

SC: Monthly list of East European Accessions (EMAI), 16, Vol. 4, No. 3,
March 1955, Uncl.

HUNGARY/Physical Chemistry. Crystals.

B

Abs Jour: Ref Zhur-Khimiya, No 22, 1958, 73044.

Author : J. Grofcsik, E. Vago.
Inst : Academy of Sciences of Hungary.
Title : Study of Polymorphous Conversions of Quartz
Taking Tridymite Formation Into Consideration.
(Preliminary Report.)

Orig Pub: Acta chim. Acad. sci. hung., 1957, 11, No 3-4,
357-358.

Abstract: It is shown that the addition of various metal oxides to quartz, when it is calcined, results in the formation of various modifications of quartz of various specific gravity (2.65 g per cub. cm in the case of low-temperature quartz and 2.3 g per cub.cm in the case of high-temperature modi-

Card : 1/2

HUNGARY/Physical Chemistry. Crystals.

B

Abs Jour: Ref Zhur-Khimiya, No 22, 1958, 73044.

fications). Besides the specific gravity determination, the process of polymorphous conversions of quartz was studied also roentgenographically, the pictures indicating the presence of cristobalite and unaltered quartz but not tridymite, although the specific gravity of the obtained calcined quartz glass was about 2.3 g per cub.cm. The x-ray data refute the generally accepted opinion that tridymite is the most important component of Dinas and prove that the methods recommended for the determination of the mineralogical composition of refractory Dinas are not applicable.

Card : 2/2

COUNTRY : Hungary
CATEGORY :

8-13

ACQ. JOUR. : Bulletin, No. 20, 1959, No. 27027

AUTHOR :
EDIT. :
TITLE :

ORIG. PUB. :

ABSTRACT : No. 2027 (15-20, 948). Documentation of the
Bulletin, 1959, No. 27, 1959. -- G. Schmidt.

CARD: 2/2

HUNGARY/Microbiology. General Microbiology. Physiology F
and Biochemistry.

Abs Jour: Ref Zhur-Biol., No 2, 1959, 5480.

Author : Gal, D.; Vago, E.

Inst : Not given.

Title : Use of the Kinetic Isotopic Method for the Study
of Transport of Substances by Bacteria.

Orig Pub: Agrokom. es talaj, 1957, 6, No 3, 223-232.

Abstract: The existence of the process of "active trans-
port" of substance into a cell is demonstrated.

Card 1/1

16

FOLDI, H.; RUSZNYAK, I.; SZABO, G.; VAGO, E.

Antihyaluronidase titer of plasma in renal disease and in
cardiac edema. *Magy. belorv. arch.* 4 no.2:66-69 1951.
(CML 20:11)

1. Doctors. 2. First Internal Clinic (Director -- Dr. Istvan
Rusznayk), Budapest Medical University.

MAGYAR, I.; VAGO, E.; DUBSKY, M.

4-5 Years follow-up of diseases condition after acute hepatitis. Orv.
hetil. 94 no.18:488-490 3 May 1953. (CIML 24:5)

1. Doctors. 2. First Internal Clinic (Director -- Prof. Dr. Istvan
Rusznayak), Budapest University.

V1150) E.

MAGYAR, I.; STEKKER, K.; VAGO, E.

Unusual course of hepatitis with intrahepatic obstruction. Orv. hetil.
94 no.46:1261-1267 15 Nov 1953. (CML 25:5)

1. Doctors. 2. First Internal Clinic (Director — Prof. Dr. Istvan
Rusznayk), Budapest Medical University.

MAGYAR, I.; RONA, Gy.; VAGO, E.

Hyperglycemia and arteriosclerosis. Acta med. hung. Suppl. 6 no.1:
64-66 1954.

1. I Klinik für innere Medizin der Medizinischen Universität,
Budapest.

(HYPERGLYCEMIA, exper.
in etiol. of arteriosclerosis in rabbits)
(ARTERIOSCLEROSIS, exper.
prod. by hyperglycemia in rabbits)

MAGYAR, Imre.; VAGO, Ersebet.; MATH, Zoltan.

Carbohydrate and kalium metabolism. 3. Effect of glycogen contents of the liver and the muscles on the kalium metabolism. Kisorletes orvostud. 7 no.1:66-72 Jan 55.

1. Budapesti Orvostudományi Egyetem I. sz. Belklinikája.
(POTASSIUM, metabolism, eff. of glycogen contents in liver & musc.)
(GLYCOGEN
liver & musc., eff. on potassium metab.)
(LIVER, metabolism
potassium, eff. of glycogen contents)
(MUSCLES,
glycogen in, eff. on potassium metab.)

MAGYAR, Imre.; VAGO, Erzsebet.; MATE, Zoltan.

Carbohydrate and kalium metabolism. 4. Levulose and kalium.
Kiserletes orvostud. 7 no.1:72-77 Jan 55.

1. Budapesti Orvostudományi Egyetem I. sz. Belklinikája.
(FRUCTOSE, effects
on liver metab., relation to potassium)
(POTASSIUM, effects
on liver metab., relation to fructose)

MAGYAR, I.; ROMA, Gy.; VAGO, M.

Experimental studies on the pathogenesis of diabetic angiopathy.
Acta med. hung. 8 no.1:37-59 1955.

1. 1-st department of medicine and 1-st Institute of Pathological
Anatomy University Medical School, Budapest.
(DIABETES MELLITUS, experimental,
causing arteriosclerosis)
(ARTERIOSCLEROSIS, experimental,
caused by diabetes mellitus)

VAGO, E.

HUNGARY/Pharmacology and Toxicology. Toxicology.

V

Abs Jour: Ref Zhur-Biol., No 10, 1958, 90027.

Author : Vago, E.; Macyar, I.

Inst : Hungarian Academy of Sciences.

Title : Relationship Between Carbohydrate and Potassium Metabolism in the Liver of Dogs in Carbon Tetrachloride Poisoning.

Orig Pub: Acta physiol. Acad. Sci. hung., 1956, 9, Suppl. 37-38.

Abstract: It was demonstrated previously that a change in serum K values, following administration of sugars, is very instructive as to the status of the glyco-genic and glycogenolytic functions of the liver and muscles. Experiments were carried out on dogs poisoned with carbon tetrachloride (CCl₄), and also . . .

Card : 1/3

HUNGARY/Pharmacology and Toxicology. Toxicology.

V

Abs Jour: Ref Zhur-Biol., No. 19, 1958, 90027.

levulose, which is transformed into glycogen only in the liver, but not in the muscles. Following introduction of dextrose into the intestines of dogs, no accumulation of glycogen was observed in the poisoned liver, and the K level was higher in the blood of the hepatic veins than in the portal veins. In the muscles, glycogen was synthesized under these conditions and the K value was lower in the peripheral veins than in the arteries. Following intravenous administration of dextrose, when it first reaches the muscles and then the liver, there is so much K retained in the muscles, that its level decreases in the peripheral blood although to a lesser extent than in normal animals under identical conditions (CCl₄ depresses glycogen metabolism

Card : 2/3

V-51

HUNGARY/Pharmacology and Toxicology. Toxicology.

V

Abs Jour: Ref Zhur-Biol., No 19, 1958, 90027.

in the muscles). After administration of levulose, the poisoned liver synthesizes glycogen better than following administration of dextrose. K is also retained in the moderately damaged liver, although hypokalemia is compensated by passage of K from the muscles. No glycogen synthesis or K retention occurs in the liver or muscles in severe liver damage. Administration of galactose, as well as of glucose, causes hypokalemia. The glycogenic function of the liver can be estimated when dextrose is administered orally (and not intravenously) by the serum K level; the lowering of the K level means that the liver synthesizes glycogen, absence of changes in the K values signifies suppression of this synthesis. -- A.G. Brusilovskaya.

Card : 3/3

MAGYAR, Imre, dr.; VAGO, Erzsébet, dr.; JELLINEK, Harry, dr.

Carbohydrate metabolism and potassium. V. Carbohydrate metabolism and potassium in liver disease. Magy. belorv. arch. 9 no.4:119-123 Aug 56.

1. A Budapesti Orvostudományi Egyetem I. sz. Belklinikájának (igazgató: Ruzsnyák, István, dr.; egyetemi tanár) közleménye.

(LIVER, metab.

potassium, eff. of sugars in exper. lesions induced by carbontetrachloride in dogs (Hun))

(CARBOHYDRATES, eff.

sugars, on liver potassium content in exper. liver lesions induced by carbontetrachloride in dogs (Hun))

(POTASSIUM, metab.

liver, eff. of sugars in exper. liver lesions induced by carbontetrachloride in dogs (Hun))

VAGO, E.

HUNGARY/General Problems of Pathology - Tumors.

T-5

Abs Jour : Ref Zhur - Biol., No 1, 1958, 3230

Author : Horanyi, M., Vago, E.

Inst : -

Title : Positive Thorn Test in Gastric Carcinoma Accompanied by Eosinophilic Leukemoid Reaction.

Orig Pub : Orv. Hetilap, 1956, 97, No 22, 611-612.

Abstract : An eosinophilic leukemoid reaction (22,000 WBD/1 cu mm; 54% eosinophils) was observed in a 76 year-old patient with a gastric carcinoma. In the bone marrow as well, eosinophils predominated (50-60%); there were 2-3% myeloblasts. Following administration of 25 mg of ACTH the number of eosinophils was decreased by 61%.

Card 1/1

SZOLNOKI, J.; VAGO, E.T.

Decomposition and humification in the soil of straw marked with isotope C^{14} . Acta agronom Hung 9 no.3/4:407-414 '59. (EEAI 9:7)

1. Bodenbiologische Gruppe des Forschungsinstituts für Bodenkunde and Agrikulturchemie der Ungarischen Akademie der Wissenschaften, Budapest.

(Hungary--Straw) (Radiosotopes) (Carbon)

1-20-57

V 3421, V 686, E. Electric spark machining. *Electronica*,
G/p 7, 1, 15-24, 1957

HUNG.

AN

Author assumes that electric spark material cutting can be regarded as a thermal process, the amount of material separated by the unit of energy depending on the quantity of heat needed for melting or evaporizing the material and on the value of thermal conductivity. The amount separated from a certain material depends—other data of the circuit being unaltered—on the charging coefficient of the condenser.

At first, solenoids, and later, electrodynamic control gear have been used by the author; the latter one worked well.

The spark cutting machine at the Laboratory of the K. Gottwald, former "Ganz" Electric Co., has been constructed with electrodynamic control, and every grade of machining can be switched on either directly or through a magnet switch by hand control. The accuracy of the tools produced with the spark cutting machine is 0.01 mm. Form fidelity is obtained by small output, surface finish by the use of small-capacity condensers.

Machining of tungsten carbide steel can be accomplished—the separation of material being 0.5-1.0 mm³/min²—with a surface finish of $R_{a, max} = 1$ micron.

Author claims that this procedure is also used with excellent results in a special large-scale production.

From author's summary by A. Lenkel, Hungary

df

VAGÓ, F.


HUNG •


621.9.018.5

1995. Theory of electric spark machining and the equipment used. F. VAGÓ. *Elektrotechnika*, 48, No. 1-2, 21-35 (~~Jan.-Feb. 1995~~) in Hungarian.


Basic problems of electro-erosion machining are explained and examples of its application are shown in tool-making. Details of control systems are given and the application of the new techniques discussed.

L. CSUEOS





VACO, Gy.



VASO, 6Y

[Faint, mostly illegible text, possibly bleed-through from the reverse side of the page]

[Handwritten signature or initials]

YAGG G.

V/10-0 G .

HUNGARY / Chemical Technology. Chemical Products and H-35
Their Application. Leather. Fur. Gelatin.
Tanning Agents. Industrial Proteins.

Abs Jour: Ref Zhur-Khimiya, No 1, 1959, 3363.

Author : Vago, G.

Inst : Not given.

Title : Concerning the Article "New Liming Methods".

Orig Pub: Bor es cipotechnika, 1956, 6, No 3, 68-69.

Abstract: See R. Zh. Khim., 1957, 3173.

Card 1/1

110

VAGO, Gy.

Problems of storing and lixivating sumac; changeability of the tannin of sumac.
p. 1. (Bor-Es Cipotechnika, Vol. 7, No. 1, Mar 1957, Budapest, Hungary)

SO: Monthly List of East European Accessions (EEAI) LC, Vol. 6, No. 2, Aug 1957. Uncl.

VAGO, György

H-35

HUNGARY/Chemical Technology, Chemical Products and Their
Application, Part 4. - Leather, Furs, Gelatin,
Tanning Agents, Industrial Proteins.

Abs Jour: Referat. Zhurnal Khimiya, No 10, 1958, 34833.

Author : László Radnóti, György Vago, Kálmán Fekete.

Inst : Not given.

Title : Retanning of Vegetably Tanned Leather Fibers in
Leather Board Manufacture.

Orig Pub: Bőr- és cipőtechnika, 1957, 7, No 2, 47-50.

Abstract: Waste leather was retanned with Al and Cr salts after
the tannides had been removed. For example, a 10 to
15% aal $Al_2(SO_4)_3$ solution was alkalized with soda
to the alkalinity of 25 to 30% at 35 to 40° before the
tannage; in order to avoid the formation of Al soap,
leather should be washed thoroughly before oiling.

Card : 1/2

16

LIUNGATY/Chemical Technology, Chemical Products and Their
Application, Part 4. - Leather, Furs, Gelatin,
Tanning Agents, Industrial Proteins.

K-35

Abs Jour: Referat. Zhurnal Khimiya, No 10, 1958, 34333.

An excess of $Al_2(SO_4)_3$ decreases the tensile strength of fibers. Similar results were obtained at the retannage with 1% aq Cr_2O_3 . Considering that the retanned fibers are not strong enough, it is expedient to use them mixed with chrome leather shavings for manufacturing leather board. A treatment of retanned fibers with CH_2O does not alter their mechanical properties, but rises their sweat resistance.

Card : 2/2

VAGO, Györy

HUNGARY/Chemical Technology, Chemical Products and Their
Application, Part 4. - Leather, Furs, Gelatin,
Tanning Agents, Industrial Proteins.

H-35

Abs Jour: Referat. Zhurnal Khimiya, No 10, 1958, 34831.

Author : Györy Vago, Laszlo Wiedner.

Inst : Not given.

Title : Tanning with Dicyandiamide Resins.

Orig Pub: Bőr-és cipőtechn., 1957, 7, No 3, 53-57.

Abstract: A synthetic tanning agent "Plastan", condensation
product of dicyandiamide with formaldehyde, was pre-
pared. It is soluble in water and contains about
44% of dry substances and about 1.6% of ashes. In
the presence of an alkaline catalyst (C), the condensation
product congeals in storage; in the case of the reaction
with an equivalent amount of hexamethylenetetramine in-

Card : 1/2


HUNGARY/Chemical Technology, Chemical Products and Their
Application, Part 4. - Leather, Furs, Gelatin,
Tanning Agents, Industrial Proteins.

H-35

Abs Jour: Referat. Zhurnal Khimiya, No 10, 1958, 34831.

stead of $\text{CH}_2\text{O} + \text{NH}_3$, sol is preserved 1 year. An increase
of CH_2O concentration accelerates the condensation at
first, a great excess of CH_2O stabilizes the resin.
"Plastan" is similar to the syntan Retingan R6.

Card : 2/2



COUNTRY : CZECHOSLOVAKIA
 CATEGORY : Chemical Technology, Chemical Products and Their Applications, Leather, Fur, Gelatine*
 ABS. JOUR. : PRAHA, NO. 23 1959, NO. 24520
 AUTHOR : Novak, O. & Bell, J. V.
 TITLE : Auto-Oxidation of Vegetable Oils in the Leather Industry
 ORIG. PUB. : Kozmetika, 1959, W, No 2, 11-14
 ABSTRACT : The auto-oxidation of vegetable oils and its dependence on the quality of top shoe leather were investigated. The spoilage occurs mainly due to leaching of the dried, unfinished and highly greased top shoe leather. Suitability of an oil for leather greasing may be controlled by determining its iron content, oxidation number and iodine number. It is recommended that in the greasing of leather, oxidation retardants/inhibitors be employed.
 INDEXING :
 Manning Materials, Industrial Proteins.
 1/2

COUNTRY :
CATEGORY :

ABST. JOUR. : RUSKOP, No. 93 1959, No. 84520

AUTHOR :
TITLE :
FIELD :

ORIG. PUB. :

ABSTRACT
Cont'd

: together with the exclusion of sunflower oil.
If this is not possible, the mild sulfonation
of the latter is recommended. Tanning proper-
ties of the epoxy-compounds were also studied.
-- M. Luksemburg.

CARD: 2/2

11 - 7

VAGO, GY.; HALASZ, I.

Waterproofing and its agents in the leather industry. (To be contd.) p. 55.

BOR-ES CIPOTECHNIKA. (Boripari Tudományos Egyesület mint a Magyar Tudományos Egyesületek Szövetsége Tagegyesülete) Budapest, Hungary, Vol. 9, No. 2, April 1959.

Monthly list of East European Accessions (EEAI) LC, Vol. 8, No. 7, July 1959. Uncla.

VAGG, SY.

Utilization of sulfite liquors in Hungary. p.146

BCR- ES CIPOTECHNIKA (Borigipari Tudományos Egyesület mint a Magyar Tudományos
Egyesületek Szövetsége Taggyesülete) Budapest, Hungary
Vol. 9, no.5, Oct.1959

Monthly List of East European Accessions (BEAI) IC., Vol.8, no.12, Dec. 1959
Uncl.

VAGO, GY., RETI, J., VARGA, P.

Autoxidation of vegetable oils in the leather industry. In German, p. 368.

ACTA Chimica. Budapest, Hungary, Vol. 20, No. 4, 1959

Monthly List of East European Accessions (EEAI) LC, Vol. 9, No. 2, Feb 1960
Uncl.

VAGO, Gyorgy, Dr.; LENART, Anna

A new method for measuring the sludginess of vegetable tanning
extracts. Bor cipo 10 no.5:129-134 S '60.

1. Boripari Kutato Intezet.

VAGO, Gyorgy, dr.

Modified protein binding materials. Bor cipo 12 no.5/6:151-155
0 '62.

1. Boripari Kutato Intezet.

GEHERNE GLUCKLICH, Judit; VAGO, Gyorgy

Polarographic determination of the monomer ethyl-acrylate contents of ethyl-acrylate- casein mixtures. *Magy kem folyoir* 68 no.4:181-184 Ap '62

1. Bor- Cipo- es Szormeipari Kutato Intezet, Budapest.

VAGO, Gyorgy, dr.; ERDI, Pal, dr.

Newer method for leather technology in the light of the application of auxiliary materials. Bor cipo 14 no.4:113-117 51 1964.

1. Research Institute of the Leather Industry (for Vago). 2. Leather Industry Enterprise; Editorial Board member, "Hor- or Cipotechnika" (for Erdi).

VAGO, Gyorgy, dr.

Stuffing of leathers dried on plates (pressed and vacuum-dried).
Bor cipo 1/4 no.6:165-169 N '64.

L 31456-66 EWP(t)/ETI IJP(c) JD/JG/AT

ACC NR: AP6023098

SOURCE CODE: HU/0031/66/000/002/0041/0043

AUTHOR: Foti, Erno (Staff scientist); Szucs, Tibor--Syuch, T. (Staff scientist); Vago, Gyorgy (Staff scientist) 61
B

ORG: [Foti] Central Research Institute for Physics (Kozponti Fizikai Kutato Intezet); [Szucs, Vago] Research Institute for Communications Technological Industry (Hiradastechnikai Ipari Kutato Intezet)

TITLE: Suspended-drop metal evaporation by electron bombardment

SOURCE: Finommechanika, no. 2, 1966, 41-43

TOPIC TAGS: electron bombardment, niobium, generator, evaporation

ABSTRACT: The suspended-drop technique for metal evaporation by electron bombardment was described. The instruments and operations involved were discussed on the basis of an example involving the evaporation of niobium. The current generator and output stabilization were described in some detail and some special considerations pertaining to the process were outlined. [JPRS]

SUB CODE: 11, 18, 20 / SUEM DATE: none

Card 1/1 ST

0915

1303

L 11352-67 EWP(k)/EWP(v)/EWP(t)/ETI IM/JD
ACC NR: AP6032798 SOURCE CODE: HU/0031/66/000/009/0265/0273

AUTHOR: Szucs, T. (Scientific assistant); Vago, G. (Scientific assistant)

ORG: Research Institute for the Communications Technological Industry
(Hiradastechnikai Ipari Kutato Intezet)

TITLE: Electron beam welding of thin metal sheets [Paper presented at a Session sponsored jointly by the Technology Department of the Association for Measuring Techniques and Automation, the Association for Acoustics and Film Technology, and the Scientific Association for Communications Technology held on 23 May 1966]

SOURCE: Finommechanika, no. 9, 1966, 265-273

TOPIC TAGS: electron beam welding, metal welding, welding technology, beryllium compound, heat sink

ABSTRACT: The authors review the literature dealing with the electron beam welding of < 1mm thick metal sheets and describe their experiences gained in the welding of beryllium bronze thin films at the Mechanical Measuring Instrument Works (Mechanikai Meromuszerek Gyara). This film is 0.12 mm thick. Satisfactory results were obtained by using the following setup. (Fig. 1) The frame is made of Bz 2 tin-bronze; the configuration illustrated provides for adequate heat sink, a lack of which function had prevented electron-ray welding of this membrane since the large amounts of heat required for the melting of the Bz 2 in the configurations used previously damaged the

Card 1/2

L 11352-67

ACC NR: AP6032798




Fig. 1. Welding technology developed for Type C heads.

film edge. The parameters of the welding operations are: acceleration voltage, 25 kV; beam current (in the operating chamber), 6 mA; welding time, 3 sec (one revolution of the jig); amount of heat supplied, 450 Wsec (i.e., 107 cal). Operating hints and means of preventing common errors are presented. The authors thank their associates at the Third Main Department for Physics at the Central Research Institute for Physics (Kozponti Fizikai Kutató Intézet III. Fizikai Főosztály), primarily Erno Foti, the project leader, for the theoretical and practical assistance given in the experimental work; Katalin Marosvolgyi for her conscientious metallographic and other analytical efforts; and Gyorgy Ispanki for his assistance in the welding and evaluation of the numerous test specimens. Orig. art. has: 2 tables and 18 figures.

SUB CODE: 11/ SUBM DATE: none

Card 2/2

VAGO, I. (Budapest XI., Muegyetem rakpart 3); UZSOKY, M. (Budapest XI., Muegyetem rakpart 3)

New method for calculating admittance of disc triodes at small signals. Periodica polytechn electr 5 no.2:107-127 '61.

1. Lehrstuhl fur Theoretische Electrizaritslehre an der Technischen Universitat. Vorgelegt von Prof. Dr. K. Simonyi.

VAGO, I. (Budapest, XI., Muegyetem rakpart 3)

A new high-frequency measuring bridge. Periodica polytechn electr
5 no.4:389-393 '61.

1. Lehrstuhl fur Theoretische Elektrizitatslehre, Technische
Universitat, Budapest Vorgelegt von Prof.Dr.K.Simonyi.

VAGO, Istvan

A theory of transmission lines consisting of cylindrical
conductors based on electromagnetic fields. *High Voltage*,
no. 6 (1966), p. 104.

1. Chair of Theoretical Electricity, Budapest Technical
University.

VAGO, Istvan (Budapest, XI., Muegyetem rakpart 3)

Calculation of the characteristics of radiating traveling waves with losses in overhead horizontal wire antennas. Periodica polytechn electr 8 no.1:47-63 '64.

1. Kafedra teoreticheskikh osnov elektrotekhniki Budapestskogo politekhnicheskogo instituta. Predstavleno zav. kafedroy prof. K. Simonyi.

VAGO, Istvan (Budapest, XI., Muogyetem rakpart 3)

The thoery of transmission lines consisting of cylindrical leads
on the basis of the electromagnetic field. Periodica polytehn
electr 8 no.3:251-264 '64.

1. Department of Theoretical Electricity of Polytechnical University
Budapest. Received Februrary 17, 1964.

VAGO, Janos

Incentive awards and plans for labor productivity increase in the
Czechoslovak Socialist Republic. Pt.1. Munka szemle 5 no.3:28-31
Mr '61.

VAGO, Janos

Incentive awards and plans for labor productivity increase in the Czechoslovak Socialist Republic. Pt.3. Munka szemle 5 no.4:28-32 Ap '61.

VAGO, Janos

State of the average wage economy in the iron and metallurgical industry during the year 1963. Munka szemle 8 no. 6: 4-8 Je '64.

KOVALCCZY, Gyorgy; VAGO, Jeno

The Szentes television transmitter. Hir techn ll no.3:91-96 Ja '60.

1. Magyar Posta.