

YARMOLENKO, V.Ye., kand.tekhn.nauk (Gomel'); GRUNTOV, P.S., inzh. (Gomel');  
SHUL'ZHENKO, P.A., kand.tekhn.nauk (Gomel')

New conditions of the work of section stations and improvement of their  
techniques. Zhel.dor.transp. 44 no.12:41-44 D '62. (MIRA 15:12)  
(Railroads--Stations)

YARMOLIK, I.F., nauchnyy sotrudnik

Role of open wells in rural areas in the epidemiology of  
Salmonella infections. Gig. i san. 26 no.9:35-40 S '61.

(MIRA 15:3)

1. Iz Saratovskogo instituta sel'skoy gigiyeny.  
(WATER SUPPLY, RURAL--HYGIENIC ASPECTS)  
(SALMONELLA)

YARMOLIK, I.F.

Comparative characteristics of some methods of studying water for the presence of microbes of the paratyphoid group. Lab. delo 8 no.4:42-45 Ap '62. (MIRA 15:5)

1. Otdel obshchey gigiyeny (zav. - prof. L.I.Los') Saratovskogo nauchno-issledovatel'skogo instituta sel'skoy gigiyeny Ministerstva zdravookhraneniya RSFSR.

(WATER--MICROBIOLOGY) (SALMONELLA)

YARMOLINSKIY, M.; YARMOLINSKIY, A.

Our reply to professor I.F. Zelikman. Sakh. prom. 32 no.4:71-72  
Ap '58.

(MIRA 11:6)

(Sugar industry)

YAPASKURT, V.V.; YEPISHIN, A.S.; SHAKIN, A.N.; SILIN, P.M.; ZHIDKOV, A.A.;  
KHELEMSKIY, M.Z.; SHEMYAKIN, P.N.; BOVIKOV, V.A.; POPOV, V.D.; BENIN,  
G.S.; HAYDENOV, A.K.; KURBATOVA, V.S.; KARTASHOV, A.K.; YARMOLINSKIY,  
A.K.; ZIBOROV, D.K.; VAYSMAN, M.L.; ZAMIROVSKIY, V.A.; SVYATENKO, M.M.

IULii Markovich Zhvirblianski; obituary. Sakh.prom.29 no.6:48 '55.  
(Zhvирblianski, IULii Markovich, 1894-1955) (MIRA 9:1)

SEMENOV, Andrey Petrovich; YARMOLINSKIY, A.S., nauchnyy red.;  
GOLOVANIVSKAYA, E.N., red.; BARANOVA, N.N., tekhn.red.

[Mechanization and automation of woodworking] Mekhaniza-  
tsiia i avtomatizatsiia derevoobrabotki; metodicheskoe po-  
sobie. Moskva, Proftekhizdat, 1963. 54 p. (MIRA 16:5)  
(Woodworking machinery) (Automatic control)

KALASHNIKOV, Petr Leont'yevich; BAKHTIYAROV, V.D., inzh.,  
retsenzent; YARMOLINSKIY, A.S., inzh., retsenzent;  
AKINDINOV, M.V., red.; KITTEL', L.S., red.izd-va;  
AKOPOVA, V.M., tekhn. red.

[Commercial study of wood and forest products] Drevesino-  
vedenie i lesnoe tovarovedenie. Moskva, Goslesbumizdat,  
1963. 253 p. (MIRA 16:12)  
(Forest products)

YARPOLINSKIY, I.B.; DROZDOV, G.M.

Over-all mechanization of the production of shoe parts and  
cardboard for shoes. Kozh.-obuv.prom. no.7:14-17 J1 '59.  
(MIRA 12:11)

(Shoe manufacture)



BERENSHTEYN, S.A.; VAYSLEYB, V.P.; VARENIK, I.F.; DOBRYNCHENKO, M.V.;  
YEGOROV, B.P.; KLISENKO, Yu.F.; MOGILEVSKIY, I.I. [deceased];  
PEREYASLAVTSEV, N.A.; PILIPENKO, V.I.; SAPOZENIKOV, F.V., inzh.;  
SHEPELEV, V.M.; SIMULEVICH, M.L.; YARMOLINSKIY, I.M.; SHAGALOV,  
Ye.S., red.; KORIKOVSKIY, I.K., red.; LARIONOV, G.Ye., tekhn. red.

[Construction of the V.I.Lenin State Regional Electric Power  
Plant in Simferopol] Opyt stroitel'stva Simferopol'skoi GRES  
im. V.I.Lenina [By] S.A.Berenshtein i dr. Moskva, Gosenergoizdat,  
1962. 151 p. (MIRA 15:6)

(Simferopol--Electric power plants)

YARMOLINSKIY, I.M., inzh.; LERNER, A.Ye,

Experience in the installation of ~~HK~~-120-100-QM boilers in a  
semienclosed type boiler system. Energ.stroi, no.30:12-20 '62.  
(MIRA 16:2)

1. Odesskiy ob'yedinenny montazhnyy uchastok tresta "Yuzhteplo-  
energomontazh."

(Boilers)

YARMOLINSKIY, I.S.

Malignant tumors of the carotid gland. Vrach. delo no.12:126-127  
D '60. (MIRA 14:1)

1. Gosptal'naya khirurgicheskaya klinika Pervogo moskovogo  
meditsinskogo instituta.  
(CAROTID BODY--CANCER)

YARMOLINSKIY, I.S.

Immediate and late results of therapy for perforated gastric ulcer  
from data of hospitals of the Don area Tula Province for the past  
10 years. Khirurgia 36 no.4:39-43 Ap '60. (MIRA 13:12)  
(PEPTIC ULCER)

KRYLOV, V.S.; YARMOLINSKIY, I.S.

Device for the introduction of vascular protheses in surgical formation of a permanent collateral shunt from the femoral into the popliteal artery. Eksper. khir. i anest. 7 no.5:49-50 S-O '62. (MIRA 17:10)

1. Iz gospital'noy khirurgicheskoy kliniki I Moskovskogo ordena Lenina meditsinskogo instituta imeni Sechenova.

YARMOLINSKIY, I.S. (Moskva, Novo-Ostapc'vskaya ul., d.6, kv.114)

Significance of oscillography in atherosclerotic occlusions of the femoral and popliteal arteries. Vest. khir. 89 no.10:25-29  
O '62. (MIRA 17:10)

1. Iz gosital'noy khirurgicheskoy kliniki 1-go Moskovskogo ordena Lenina meditsinskogo instituta imeni Sechenova (zav. - prof. B.V. Petrovskiy).

YARMOLINSKIY, I.S.

Clinical aspects, diagnosis, and surgical treatment in atherosclerotic occlusions of the femoral and popliteal arteries.  
Med. sestra 22, no.1:31.-33 Ja '63. (MIRA 16:7)

1. Iz Gospital'noy khirurgicheskoy kliniki I Moskovskogo ordena Lenina meditsinskogo instituta imeni I.M.Sechenova, Moskva.  
(ARTERIOSCLEROSIS) (ARTERIES—SURGERY)

YARMOLINSKIY, I.S.

Operation of permanent shunting with vascular prostheses in arteriosclerotic occlusions of the femoral and popliteal arteries. Sovet. med. 27 no.6:9-13 Ja'63 (MIRA 17:2)

1. Iz gosspital'noy khirurgicheskoy kliniki (direktor - day-stvitel'nyy chlen AMN SSSR prof. B.V.Petrovskiy) i Moskovskogo ordena Lenina meditsinskogo instituta imeni I.M.Sechenova.



YARMOLINSKIY, M.

Quality of white sugar. Sakh.prom. JB no.1:69-70 Ja '64.

(MIRA 17:2)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut sakharnoy promyshlennosti.

YARIGOLENITSKY, H. S.

PROCESSES AND PROPERTIES

CA 20

A new combined process for the industrial utilization of extracted beet slices. *Med. Khim.* 1934, II, 382.—A process is given for obtaining pectin, yeast-cake adhesives, and pressed cake for cattle feed from extd. beet slices. The slices are boiled 15-30 min., sepd. from the ext. (A), and by heating the residue (contg. 7.5% dry substance) at 1-1.5 atm. for 15-20 min. and pressing, thick sirup (B) and pressed cake are obtained. A is evapd. until the dry substance amounts to 14-16%, cooled to 25°, 3% of the dry wt. of yeast added and the material allowed to ferment. During fermentation sepn. into 2 layers occurs. From the upper, sol. pectin is obtained; the org. acids and odor-producing materials are completely, and the bitter taste is partially, removed. The lower—yeast and insol. pectin—serves for cattle feed. B is treated like A; after fermentation the upper layer yields glucan; the lower, insol. pectin.

W. A. Moore

APR-51A METALLURGICAL LITERATURE CLASSIFICATION

Common Transmittal Unit

YARMOLINSKIY, M. B.

CA

Рациональный метод в кристаллизации низкосортной маассекулты. М. Б. Яромлинский, *Сакхурнаы Пром.* 23, No. 5, 31-2 (1949). -- A device is described for spraying hot water on low-grade masscutes discharged from the vacuum pan into crystallizers. The purpose of this spraying is to redissolve false grains. The device is simple in design, is installed in a trough, and activated automatically by the discharged masscuite. V. E. B.

ASS-55A METALLURGICAL LITERATURE CLASSIFICATION

1ST AND 2ND LETTERS

3RD AND 4TH LETTERS

5TH LETTER

6TH LETTER

7TH LETTER

8TH LETTER

9TH LETTER

10TH LETTER

11TH LETTER

12TH LETTER

13TH LETTER

14TH LETTER

15TH LETTER

16TH LETTER

17TH LETTER

18TH LETTER

19TH LETTER

20TH LETTER

21ST LETTER

22ND LETTER

23RD LETTER

24TH LETTER

25TH LETTER

26TH LETTER

27TH LETTER

28TH LETTER

29TH LETTER

30TH LETTER

31ST LETTER

32ND LETTER

33RD LETTER

34TH LETTER

35TH LETTER

36TH LETTER

37TH LETTER

38TH LETTER

39TH LETTER

40TH LETTER

41ST LETTER

42ND LETTER

43RD LETTER

44TH LETTER

45TH LETTER

46TH LETTER

47TH LETTER

48TH LETTER

49TH LETTER

50TH LETTER

51ST LETTER

52ND LETTER

53RD LETTER

54TH LETTER

55TH LETTER

56TH LETTER

57TH LETTER

58TH LETTER

59TH LETTER

60TH LETTER

61ST LETTER

62ND LETTER

63RD LETTER

64TH LETTER

65TH LETTER

66TH LETTER

67TH LETTER

68TH LETTER

69TH LETTER

70TH LETTER

71ST LETTER

72ND LETTER

73RD LETTER

74TH LETTER

75TH LETTER

76TH LETTER

77TH LETTER

78TH LETTER

79TH LETTER

80TH LETTER

81ST LETTER

82ND LETTER

83RD LETTER

84TH LETTER

85TH LETTER

86TH LETTER

87TH LETTER

88TH LETTER

89TH LETTER

90TH LETTER

91ST LETTER

92ND LETTER

93RD LETTER

94TH LETTER

95TH LETTER

96TH LETTER

97TH LETTER


98TH LETTER

99TH LETTER

100TH LETTER

YARMOLINSKI, M.

Crystallization of sugar from its aqueous solutions. M. Yarmolinski. *Trudy Vsesoyuz. Nauch.-Issledovatel. Inst. Sakhar. Prom.* 1953, No. 2, 104-15; *Referat. Zhur. Khim.* 1955, No. 1397.—The rate of crystal formation of sugar from its aq. solns. and the amt. of crystals increase with the supersatn coeff. In supersatd. sugar solns. there is a metastable zone outside of which spontaneous crystn. occurs. Stirring of a supersatd. sugar soln. promotes the formation of seed crystals. M. Harech



"Investigation of the Formation of Sugar Crystals." Cand Tech Sci, Moscow Technological Inst of the Food Industry, Min Higher Education USSR, Moscow, 1955. (KL, No 11, Apr 55)

SO: Sum. No. 704, 2 Nov 55 - Survey of Scientific and Technical Dissertations Defended at USSR Higher Educational Institutions (16).

YAROMOLINSKIY, M.B.; TERESHIN, B.N.

Nomogram for determining sugar crystals in masscuite. Sakh.prom.  
30 no.3164 Mr '56. (MLRA 9:7)

1. TSentral'nyy nauchno-issledovatel'skiy institut sakharnoy promy-  
shlennosti.

(Sugar--Analysis and tesing)

YARMOLINSKIY, M.B., kandidat tekhnicheskikh nauk; ZDANOVICH, I.L.,  
ispolnyayushchiy obyazannosti starshego nauchnogo sotrudnika;  
OZOL, M.Ya., khimik-analitik.

Use of corrugated cardboard boxes for packing pressed sugar and  
its transportation by sea. Trudy TSINS no.4:148-162 '56.  
(MLRA 10:5)

1. Rafinadnaya laboratoriya.  
(Boxes) (Sugar--Transportation)



YARMOLINSKIY, M.

Sugar packing machine. Sakh.prom.30 no.5:64-65 'My: 156. (MIRA 9:9)  
(Packaging machinery)

YARMOLINSKIY, M.B.

Packaging granulated sugar in multiwall paper bags with side spout.  
Sakh. prom. 32 no.2:34 P '58. (MIRA 11:3)

1. Tsentral'nyy nauchno-issledovatel'skiy institut sakharnoy  
promyshlennosti.  
(Sugar) (Paper bags)

YARMOLINSKIY, M.; YARMOLINSKIY, A.

Our reply to professor I.F. Zelikman. Sakh. prom. 32 no.4:71-72

Ap '58.

(MIRA 11:6)

(Sugar industry)

YARMOLINSKIY, M.B.

Recovery of lime from filter-press cake. Sakh. prom. 32 no.11;40-41  
N '58. (MIRA 11:12)

1. Tsentral'nyy nauchno-issledovatel'skiy institut sakharney promysh-  
lennosti.

(Lime)

TARASOV, M.N.; YARMOLINSKIY, M.B.

Packing refined sugar into large bags. Sakh. prom. 32 no.12:48-49  
D '58. (MIRA 11:12)

1.Krasnopresnenskiy refinadnyy zavod (for Tarasov). 2.TSentral'nyy  
nauchno-issledovatel'skiy institut sakharnoy promyshlennosti (for  
Yarmolinskiy)  
(Sugar--Transportation)

YARMOLINSKIY, M.B.

Progressive technology of the production of refined sugar. Sakh.  
prom. 33 no.3:8-10 Mr '59. (MIRA 12:4)

1. Tsentral'nyy nauchno-issledovatel'skiy institut sakharnoy  
avekly.

(Sugar manufacture)

YARMOLINSKIY, M.B.; ZDANOVICH, I.L.; BRENMAN, M.A.; ALEKSEYENKO, F.P.

Use of granulated coal in the sugar refining industry. Sakh.  
prom. 35 no.12:21-26 D '61. (MIRA 15:1)

1. Tsentral'nyy nauchno-issledovatel'skiy institut sakharnoy  
promyshlennosti.

(Sugar manufacture)

(Coal)

YARMOLINSKIY, M.B.

Results of the work of sugar refining factories in the Ukrainian  
S.S.R. during 1962. Sakh.prom. 37 no.6:5-8 Je '63. (MIRA 16:5)

1. Tsentral'nyy nauchno-issledovatel'skiy institut sakharnoy  
promyshlennosti.

(Ukraine—Sugar factories)



YARMOLINSKIY, M.B.

Problems of equipment and technology in sugar refineries. Sakh.  
prom. 37 no.11:8-10 N '63. (MIRA 16:11)

1. Tsentral'nyy nauchno-issledovatel'skiy institut sakharnoy pro-  
myshlennosti.

TERESHCHENKO, I.P.; MOSKVIN, O.I.; DARAGAN, M.V.[Darahan, M.V.];  
ANISIMOV, V.P.; YARMOLINSKIY, M.R.[Iarmolyns'kyi, M.R.];  
BULGAKOV, P.S.[Bulbakov, P.S.]; KUTS, V.K.; KASHPUR, A.V.;  
VASILENKO, G.K.[Vasylenko, H.K.]; KUKOLEV, V.D.[Kukoliev,  
V.D.]; SIGOV, S.G.[Sihov, S.H., deceased]; NAGIRNYAK, P.A.  
[Nahirniak, P.A.]; VETCHINOV, I.A.[Vetchynov, I.A.];  
ZADOROZHNYI, V.K.; DROSOVSKAYA, L.I.[Drosovs'ka, L.I.];  
SHKITINA, M.I.; PROSHCHAKOV, O.M.; MOKIYENKO, B.F.  
[Mokiienko, B.F.]; GOLOVACH, A.V.[Holovach, A.V.];  
IVANITSKIY, I.V.[Ivanyts'kyi, I.V.]; KOZAK, V.Ye.;  
BORYAKIN, V.M., red.izd-va; NESTERENKO, O.O., glav. red.;  
DAKHNO, Yu.B., tekhn. red.

[National income of the Ukrainian S.S.R. during the period  
of the large-scale building of communism] Natsional'nyi  
dokhod Ukrains'koi RSR v period rozhornutoho budivnytstva  
kommunizmu. Red.kol.: O.O.Nesterenko ta inshi. Kyiv, Vyd-  
vo AN URSR, 1963. 333 p. (MIRA 16:12)

1. Akademiya nauk URSR, Kiev. Instytut ekonomiky.  
(Ukraine--Income)

SLINCHENKO, Ye.V.; YARMOLINSKIY, N.P.; KUDOYAROV, M.S.; AMAN'YEV, P.V.

Blast furnace operation with evaporation cooling. Metallurg  
7 no.7:9-11 JI '62. (MIRA 15:7)

1. Kuznetskiy metallurgicheskiy kombinat.  
(Blast furnaces---Cooling)

TEL'PUKHOVSKIY, N.A.; Prinimali uchastiye: MOROZ, A.M.; YARMOLINSKIY, S.Kh.;  
MALOMYZHEV, L.M.; BURKOV, V.I.

Electronic circuit of an apparatus for the emission of  
exact time signals. Trudy inst. Kom. stand., ser i izm.  
prib. no.58:108-119 '62. (MIRA 15:11)

1. Sotrudniki Irkutskoy laboratorii Vsesoyuznogo nauchno-  
issledovatel'skogo instituta fiziko-tehnicheskikh i  
radiotekhnicheskikh izmereniy (for Moroz, Yarmolinskiy,  
Malomyzhev, Burkov).

(Time signals)

MARKOVSKIY, A.V., inzh.; YARMOLINSKIY, V.D.

Sectional kitchen stove with infrared burners. Gor.khoz.  
Mosk. 34 no.4:36-37 Ap '60. (MIRA 13:8)  
(Infrared rays--Industrial applications)  
(Stoves)

RATTS, Emmanuil Genrikhovich, kand. tekhn. nauk; TSEYTLIN, Semen Yudovich, kand. tekhn. nauk; MELCOVIDOV, Konstantin Ivanovich, inzh.; YARMOLINSKIY, Vladimir Meiseyevich, inzh.; ANTONOVA, N.N., inzh., red.

[12 m reinforced concrete beams with rod reinforcement stressed by the electrothermal method; practices of the All-Union Research Institute for Reinforced Concrete and Plant No.18 for Reinforced Concrete Products of the Main Administration for Building Materials of the City of Moscow] Zhelezobetonnye podkrasovyye balki dlinoi 12 M so sterzhnevoi armaturoi, napriagaemoi elektrotermicheskim metodom; opyt raboty VNIIZhelezobetona zavoda zhelezobetonnykh izdelii no.18 Glavmospromstroimaterialov. Moskva, Strolizdat, 1964. 31 p. (MIRA 18:5)

1. Zaveduyushchiy laboratoriyey sbornykh zhelezobetonnykh konstruksiy Vsesoyuznogo nauchno-issledovatel'skogo instituta zavodskoy tekhnologii sbornykh zhelezobetonnykh konstruksiy i izdeliy (for Ratts). 2. Zaveduyushchiy sektorom inzhenernoy konstruksiy Vsesoyuznogo nauchno-issledovatel'skogo instituta zavodskoy tekhnologii sbornykh zhelezobetonnykh konstruksiy i izdeliy (for Tseytlin). 3. Glavnyy konstruktor sektora inzhenernykh konstruksiy Vsesoyuznogo nauchno-issledovatel'skogo instituta zavodskoy tekhnologii sbornykh zhelezobetonnykh konstruksiy i izdeliy (for Milovidev).

USSR/Soil Science - Physical and Chemical Properties of Soils. J

Abs Jour : Ref Zhur Biol., No 22, 1958, 100008

Author : ~~Yamaalinskiy, Ya. A.~~ Golovin, V.V.

Inst : Academy of Sciences TadzhSSR

Title : Depth of the Constant Yearly Temperature in the Sub-soil  
of Uzbekistan and Tadzhikistan.

Orig Pub : Dokl. AN TadzhSSR, 1956, No 19, 17-20

Abstract : No abstract.

Card 1/1

GOLOVIN, V.V.; YARMOLINSKIY, Ye.A.

Temperature conditions of Tajikistan rivers. Izv. Otd. est. nauk  
AN Tadzh.SSR no.19:57-66 '57. (MIRA 11:8)

1. Kafedra melioratsii Tadzhikskogo sel'khozinstituta.  
(Tajikistan--Rivers--Temperature)



Card 1/2

Development are developed.

L 59561-65

ACCESSION NR: AP5013849

0

... differential equations were solved on an electronic simulator;  
... experimental verification

NO REF COPY

GRIGOR'YEV, V.Ye.; CHURKIN, V.M.; YARMOLOVICH, E.S.

Calculation of the static characteristics of an electrohydraulic amplifier with pulse-width control. Avtom. 1 telem. 26 no.5: 918-923 My '65. (MIRA 18:12)

1. Submitted November 10, 1963.

L 1686-66 EPA/EWT(1)/EPA(s)-2/EWT(m)/FA/EPF(n)-2/ENG(m)/T-2/ETC(m). WW  
ACCESSION NR: AP5014818 UR/0209/65/000/006/0067/0072

AUTHOR: Yarmolovich, G. (Engineer, Lieutenant Colonel) 403

TITLE: Operational [and maintenance] characteristics of the L-29 aircraft engine

SOURCE: Aviatsiya i kosmonavtika, no. 6, 1965, 67-72

TOPIC TAGS: aircraft fuel system, fuel pump, fuel filters, aircraft engine/M701 engine

ABSTRACT: The author discusses the operational and maintenance characteristics of the M701 VS-150 and M701 S-250 engines, which are used on the L-29 aircraft. Emphasis is placed on the fuel system, including fuel supply, pumps, filters, and fuel atomizer. The different effects of the barostatic-control factor on M701 engines are compared to those of other aircraft engines, and safety and reliability factors are pointed out. Orig. art. has: 3 figures.

ASSOCIATION: none

Card 1/2

L 1686-66

ACCESSION NR: AP5014818

SUBMITTED: 00

ENCL: 00

SUB CODE: AC, PR <sup>0</sup>

NO REF SOV: 000

OTHER: 000

Card 2/2 *RP*

YARMOLOVICH, G., inzh.-pocpolkovnik

Characteristics of the operation of the i-29 airplane engine.

Av. 1 kosm. 47 no.6:67-72 Ia '65.

(MIR 4 18:5)

L 38417-66 ENP(τ)/ETI IJP(c) HW/JD

ACC NR: AP6018274

SOURCE CODE: PO/0038/65/010/004/0323/0336

48-1B

AUTHOR: Yarmolovich, G.--Jarmolowicz, H.

ORG: Institute of Physical Chemistry, Warsaw (Instytut Chemii Fizycznej PAN)

TITLE: Some magnetic properties and the structure of nickel electrodeposits obtained from chloride and sulfate solutions

SOURCE: Archiwum hutnictwa, v. 10, no. 4, 1965, 323-336

TOPIC TAGS: magnetic property, electrodeposition, chloride, ~~sulfate~~, sulfate, ~~solution~~, electrolysis, magnetic coercive force, magnetic saturation, annealing

ABSTRACT: In this paper the author presents the results of investigations of the effects of electrolysis conditions (pH, current density) on some magnetic properties (coercive force, residual magnetism, magnetic saturation) and on the structure of nickel electrodeposits obtained from chloride and sulfate baths. The sulfate bath contained CdCl<sub>2</sub> as a brightening agent. The deposits examined were received at pH 3.0-5.7 and at current densities 3-30 mA/cm<sup>2</sup>. It was stated that the coercive force of the deposits increases with increasing current density and pH value. The effect of annealing the deposits at 400 and 600C on their magnetic properties was

Card 1/2

L 38117-66

ACC NR: AP6018247

also the subject of studies. X-ray and electronographic analysis as well as electron micrographs of the deposits surfaces revealed distinct differences in the type of texture and in surface structures of the deposits received in different conditions of electrolysis. A relationship between the structure of the nickel deposits and their magnetic properties was stated. The author expresses his gratitude to Professor K. M. Gubrunova and senior scientific associate Yu. M. Polukanova for their attention and valuable advice during the study. Orig. art. has: 5 figures and 2 tables. [Author's abstract.]

2

SUB CODE: 07, ~~26~~ SUBM DATE: 01Dec64/ ORIG REF: 001/ SOV REF: 007/ OTH REF: 009 [KS]

Card 2/2 fb



YARMOLOVICH, I.P., dotsent

Estimate of errors in the joining of converging working faces  
and errors of orientation through two shafts. Izv. vyz. ucheb.  
zav.; gor. zhur. 7 no.10:27-32 '64.

(MIRA 18:1)

1. Vsesoyuznyy zaochnyy politekhnicheskyy institut. Rekomendovana  
kafedroy marksheyderskogo dela i geodezii.

YARMOLOVICH, I.P., dotsent

Analysis of mine surveying errors in traversing with the help  
of matrices. Izv.vys.ucheb.zav.:gor.zhur. 7 no. 1:68-75 '64.  
(MIRA 17:5)

1. Vsesoyuznyy zaochnyy politekhnicheskiy institut. Rekomendovana  
kafedroy marksheyderskogo dela i geodezii.

SVYATSKIY, Pavel Stanislavovich, inzh.; YARMOLOVICH, Konstantin Yulianovich, inzh.; SMIRNOV, N.A., prof., red.; FOMICHEV, A.G., red. izd-va; BELOGUROVA, I.A., tekhn. red.

[Methods of overall mechanization of the basic types of finishing work] Puti kompleksnoi mekhanizatsii osnovnykh vidov otdelochnykh robot. Pod obshelei red. N.A.Smirnova. Leningrad, Leningr. dom nauchno-tekhn. propagandy, 1961. 20 p. (Bibliotekha stroitel'ia po mekhanizatsii i avtomatizatsii stroitel'stva, no.14) (MIRA 15:7)

(Building--Details)

GALKIN, B.I.; BIRYUKOV, V.I.; KREYTER, V.M.; KULICHIKHIN, S.N.;  
ORLOVA, Ye.V.; POMERANTSEV, V.V.; RUSetskAYA, G.G.;  
YARMOLOVICH, N.V.; MAKEYEV, V.I., red. izd-va; BYKOVA,  
V.V., tekhn. red.

[Prospecting for stockwork deposits of nonferrous and rare  
metal ores] Razvedka shtokverkovykh mestorozhdeni tsvetnykh i  
redkikh metallov. [By] B.I.Galkin i dr. Moskva, Goageoltekh-  
izdat, 1962. 233 p. (MIRA 16:6)

1. Moscow. Vsesoyuznyy nauchno-issledovatel'skiy institut mine-  
ral'nogo syr'ya.

(Prospecting)

SHEVCHENKO, N.F., red.; AMELIN, F.S., red.; GRECHKO, V.Ye., red.; ISAYEV, V.I., red.; KUZUBOV, V.I., red.; LIBERMAN, Ye.G., prof., doktor ekonom.nauk, red.; MAKARENKO, V.P., red.; SHCHERBININ, I.P., red.; YARMOLOVICH, O.M., red.; KARDASH, G.I., red.; DONSKOY, Ya.Ye., red.; LIMANOVA, M.I., tekhn.red.

[First and foremost; ways to further increase labor productivity in machinery manufacturing enterprises of Kharkov] Samoe vazhnoe, samoe glavnoe; o putiyakh dal'neishogo povysheniya proizvoditel'nosti truda na mashinostroitel'nykh predpriyatiyakh Khar'kova. Khar'kov, Khar'kovskoe knizhnoe izd-vo, 1960. 205 p.

(MIRA 13:11)

1. Ukraine. Khar'kovskiy gorodskoy ekonomicheskoy administrativnyy rayon. Sovet narodnogo khozyaystva. 2. Nachal'nik tekhnicheskogo otdela Khar'kovskogo sovmarkhoza (for Kuzubov). 3. Khar'kovskiy inzhenerno-ekonomicheskoy institut (for Liberman).  
(Kharkov--Machinery industry--Labor productivity)

~~YARMOLYUK, V.A.~~  
~~YARMOLYUK, V.A.~~

The Tugur-Nemilen mesopotamia. Yop.geog.Dal'.Vost.no.3:92-101  
'57. (MIRA 10:12)  
(Tugur Valley) (Nemilen Valley)

AUTHOR: Yarmolyuk, V., Mining Engineer 30V/29-58-7-12/23

TITLE: Automatization in Coal Mines (V shakhte khozyaynichayut avtomaty)

PERIODICAL: Tekhnika molodezhi, 1958<sup>ib</sup> Nr 7, pp. 19-21 (USSR)

ABSTRACT: Industrial development is making ever increasing demands on science and technical engineering. This article deals with a new kind of coal mine projected by the "Ukrghproshakht" Institute for the L'vov-Volyn' basin. A factory operating without human labor is nothing new, but a coal mine without any miners working on its bottom seems to be a mere creation of fancy. Engineers of various branches participated in the task of working out a project for a fully automatized shaft of a mine. They abandoned the traditional manner of building and adopted the latest technical methods developed both in the USSR and in other countries. In order to reach the bed of coal in the pit, two vertical holes having a diameter of 6 m are bored by means of a special equipment; they are secured by means of re-inforced concrete stays. Adjoining these bore-holes under the ground there are additional auxiliary caverns and main tunnels, through which the coal is transported. The machines

Card 1/3

Automatization in Coal Mines

SOV/29-58-7-12/23

existing at present are very complicated and large, and they take up much room on the floors of pits. They do not, however, in all cases warrant normal working of solid layers. In spite of the experience gathered in the course of many years no success has as yet been attained with respect to the production of mechanized apparatus for long walls. The history of the development of the coal industry in the USSR shows that, as a rule, the endeavor was made to increase the length of the long wall. This method was abandoned in the case of the new shaft, and it was decided to shorten the long wall by half its length, i.e. from 100 to 50 m. Shortening of the long wall made it easier to have the conveyor band moving along the entire length of the floor without having to take sections apart in the L'vov-Volyn' basin. Also the construction, assembly, and dismantling of the hydraulic support as well as its operation were rendered considerably more simple. What is most important is, however, that coal will be conveyed from the long wall without the aid of human labor. In this way danger and diseases will be avoided. The entire machinery, which consists of 6 different parts, will work in 3 shifts of 6 hours each, and will dig 3000 t of coal out of the mine within 24 hours. Much attention

Card 2/3



Automatization in Coal Mines

SOV/29-58-7-12/23

was paid in the course of projecting also to the problem of transportation.

All these measures made it possible to treble production figures, to reduce the prime costs of a ton of coal by half its amount, and to reduce the number of workers to one third of its previous amount. There is 1 figure.

1. Coal mines--USSR
2. Coal--Applications

Card 3/3

~~YARMOLYUK, Viktor Andreyevich; SIMKHO, Kh.S., red.; KAYDALOVA, M.D.,  
tekhn.red.~~

[Put mineral resources of Khabarovsk Territory at the service  
of the seven-year plan] Poleznye iskopaemye kraia - na sluzhbu  
semiletki. Khabarovsk, Khabarovskoe knizhnoe izd-vo, 1959.  
39 p. (MIRA 12:12)  
(Khabarovsk Territory--Mines and mineral resources)

NUZHNOV, S.V.; YARMOLYUK, V.A.

Later Pre-Cambrian in the southeastern border of the Russian Platform. Sov.geol. 2 no.7:21-31 J1 '59. (MIRA 13:1)

1. Dal'ne-Vostochnoye geologicheskoye upravleniye.  
(Russian Platform--Geology, Stratigraphic)

SOV/20-127-1-42/65

3(5)  
AUTHORS:

Vlasov, G. M., Yarmolyuk, V. A.

the Peninsula

TITLE:

The Structural-tectonic Regions of Kamchatka (Strukturno-tektonicheskiye rayony Kamohatki)

PERIODICAL:

Doklady Akademii nauk SSSR, 1959, Vol 127, Nr 1, pp 156-158 (USSR)

ABSTRACT:

In recent years, important details helping to define the geological structure of Kamchatka have been detected by the geologists of the Dal'nevostochnoye (Far East) and Pyatoye (Fifth) geologicheskoye upravleniye (Geological Administrations). This has become evident with the composition of the new geological survey map (1957) (Fig 1). Kamchatka is a young folding area, chiefly consisting of cenozoic and closely connected Cretaceous formations. A middle massif of metamorphic rocks and of such risen from the depth by eruption lies in the south of the Sredinnyy chain. Two directions become evident in the chief structures of the peninsula: (a) a southern one corresponding to the prolongation of the middle massif, and (b) a north-eastern one connected with the range of two parallel fold arcs. One of the arcs (Vostochnyy Kamchatskiy chain) is the outer,

Card 1/4

The Structural-tectonic Regions of <sup>the</sup> Kamchatka Peninsula

SOV/20-127-1-42/65

and the other (the central and northern part of the Sredinnyy chain) is the inner and roughly follows the axis of the peninsula. The arcs are the most important structural-facial and -metallogenic zones. The outer arc consists of an anticlinal elevation from the Upper Cretaceous sediments and volcanic rocks, also ultrabasic ones among them. Pyrite-ore occurrences with copper, nickel, cobalt, occasionally with molybdenum are found here. Exclusively cenozoic volcanic rocks are developed in the inner range of the arc. There are ore occurrences of: mercury, antimony, gold, copper, and molybdenum, which have formed near the surface and rest in the midst of secondary quartzites and propylites. The arcs bend eastwards at the hitherto little investigated transversal south-eastern dislocation zone which crosses the peninsula near Petropavlovsk. The inner arc is continued as Bol'shaya Kuril'skaya insular chain (Great Kuriles), while the outer arc apparently follows a subaqueous projection eastwards, reaching the Malye Kuril'skiye (Little Kuriles) islands. Both arcs are continued in the north-east as Koryakskoye nagor'ye (Koryak mountains). The outer arc unites with the structures of Alaska. The Sredinnyy massif is prolonged northwards by young anticlinal folds. This

Card 2/4

The Structural-tectonic Regions of <sup>the</sup> Kamchatka Peninsula

SOV/20-127-1-42/65

massif and both the mentioned arcs form the chief anticlinoria of Kamchatka and the orographically marked mountain chains. A 4th smaller anticlinorium consisting of Paleogene and Miocene volcanic rocks and of flysch stretches along the eastern peninsula. Synclinoria lie between the anticlinoria: Western Kamchatskiy, Palanskiy, Parapol'skiy, and Penzhinskiy with Neogene coal- and petroleum bearing sediments. They represent a rear-downwarping (tylovoy progib) which is divided into two parts by the central elevation. The Tsentral'naya Kamchatskaya depression forms an inner (vnutrennyaya) downwarping between the fold arcs. Oceanic marginal (peredovyye) depressions are near the east coast and south of the Aleutians. Along with the two young fold arcs there are fractures in depth, along which powerful tertiary and Upper Cretaceous volcanic masses are spread. Three quaternary volcanic zones lie all over the mentioned structural elements. There is 1 figure.

ASSOCIATION:  
Card 3/4

Dal'nevostochnoye geologicheskoye upravleniye g. Khabarovsk  
(Far Eastern Geological Administration, City of Khabarovsk)

← YARMOLYUK, V.A., geolog

Mineral resources in the middle and lower Amur Basin within the  
Soviet Union. Amur sbor. no.2:218-240 '60. (MIRA 15:3)

1. Deystvitel'nyy chlen Geograficheskogo obshchestva SSSR.  
(Amur Province--Mines and mineral resources)  
(Khabarovsk Territory--Mines and mineral resources)

88907

S/026/60/000/011/004/009  
A166/A026

3.9000 (1041, 1109, 1327)

AUTHOR: Yarmolyuk, V.A. (Khabarovsk)

TITLE: The Dzhugdzhura Meteorite

PERIODICAL: Priroda, 1960, No. 11, pp. 100 - 101

TEXT: On December 22, 1959, at 10:30 h Moscow Time (or 17:30 h Local Time) a meteorite fell in the Dzhugdzhura Mountains somewhere near the upper reaches of the Northern Uy, Ul'ya, Etandzha and Ulkan Rivers. The article reproduces reports from eyewitnesses: I. Yuzefov, a teacher; F.V. Kostyuchek; I.Ye. She-mukova; P.A. Bredelov, Director of the Aldoma-7-Year School; Ye.V. Golub and G.I. Kozlov; G.Ya. Sorokin; L.F. Shokhirev; N.S. Yashkin, an engineer-econo-mist at the Aldomskiy rybozavod (Aldoma Fish Plant); Bondarenko, Chairman of the Aldomskiy sel'sovet (Aldoma Village Soviet); Mudariyeva; Volkova, Head of the Ayan Hydrometeorological Service; Tyutyunnikov; Fedayayev, Chief of Totta Radio Station; Amosov. The reports place the holid's fall in an area of more than 200,000 m<sup>2</sup>. The consensus of opinion describes the phenomenon as a fiery-white ball with a tail the same color, proceeding from the direction of Northern Sakh-alin (an azimuth of 345° from Aldoma). The meteorite exploded with a deafening roar which echoed for some 4 - 5 min. There is 1 map.

Card 1/1



YARMOLYUK, V.A.

Walled up "treasure trove.". Priroda 50 no.1:24, Ja '61.  
(MIRA 14:1)

1. Dal'nevostochnoye geologicheskoye upravleniye, Khabarovsk.  
(Trees)

ABDULLAYEV, Kh.M.; ALYAVDIN, V.F.; AMIRASLANOV, A.A.; ANIKEYEV, N.P.;  
ARAPOV, Yu.A.; BARSANOV, G.P.; BELYAYEVSKIY, N.A.; BOKIY, G.P.;  
BORODAYEVSKAYA, M.B.; GOVOROV, I.N.; GODLEVSKIY, M.N.; SECHZGLOV, A.D.;  
SHAKHOV, F.N.; SHILO, N.A.; YARMOLYUK, V.A.; DRAKIN, I.Ye.;  
YEROFEYEV, B.N.; YERSHOV, A.D.; IVANKIN, P.F.; ITSIKSON, M.I.;  
KARPOVA, Ye.D.; KASHIN, S.A.; KASHKAY, M.A.; KORZHINSKIY, D.S.;  
KOSOV, B.M.; KOTLYAR, V.N.; KREYTER, V.M.; KUZNETSOV, V.A.; LUGOV,  
S.F.; MAGAK'YAN, I.G.; MATÉRIKOV, M.P.; ODI NTSOV, M.M.; PAVLOV, Ye.S.;  
SATPAYEV, K.I.; SMIRNOV, V.I.; SOBOLEV, V.S.; SOKOLOV, G.A.; STRAKHOV,  
N.M.; TATARINOV, I.M.; KHRUSHCHOV, N.A.; TSAREGRADSKIY, V.A.;  
CHUKHROV, F.V.

In memory of Oleg Dmitrievich Levitskii; obituary. Sov.geol. 4  
no.5:156-158 My '61. (MIRA 14:6)  
(Levitskii, Oleg Dmitrievich, 1909-1961)

YARMOLYUK, V.A.

Achievements of the Far Eastern geological survey during  
the 40 years of Soviet power. Razved. i okhr. nedr 28  
no.10:1-6 0 '62. (MIRA 15:11)

1. Dal'nevostochnoye geologicheskoye upravleniye.  
(Soviet Far East--Geology, Economic)

VLASOV, G.M.; YARMOLYUK, V.A.; ZHEGALOV, Yu.V.

Some basis tectonic problems of Kamchatka. Sov. geol 6 no.6:  
32-50 Je '63. (MIRA 16:7)

1. Dal'nevostochnoye geologicheskoye upravleniye.  
(Kamchatka—Geology, Structural)

BELYAYEVSKIY, N.A., red.; ALI-ZADE, A.A., red.; ALIYEV, M.M., red.;  
BAKIROV, A.A., red.; BELOUSOV, V.V., red.; BEUS, A.A., red.;  
BOGDANOV, A.A., red.; BORISOV, A.A., red.; BRENNER, M.M.,  
red.; DYUKOV, A.I., red.; YERSHOV, A.D., red.; ZARIDZE, G.M.,  
red.; KALUGIN, A.S., red.; KOSOV, B.M., red.; KOPTEV-  
DVORNIKOV, V.S., red.; KOTLYAR, V.N., red.; LUGOV, S.F., red.;  
MAGAK'YAN, I.G., red.; MARINOV, N.A., red.; MARKOVSKIY, A.P.,  
red.; MALINOVSKIY, F.M., red.; PUSTOVALOV, L.V., red.; SATPAYEV,  
K.I., red.; SEMENENKO, N.P., red.; TYZHNOV, A.V., red.;  
KHRUSHCHOV, N.A., red.; SHCHEGOLEV, D.I., red.; YARMOLYUK, V.A.,  
red.

[Materials on regional tectonics of the U.S.S.R.] Materialy po  
regional'noi tektonike SSSR. Moskva, Izd-vo "Nedra," 1964. 193 p.  
(MIRA 17:4)

1. Russia (1923- U.S.S.R.) Gosudarstvennyy geologicheskii ko-  
mitet.

YARMOLYUK, V.A.; VARNAVSKIY, V.G.

Prospects for finding oil and gas in the *Ayan-Maya* region of  
the Khabarovsk Territory. Sov. geol. 7 no.6:23-31 Je '64  
(MIRA 18:1)

1. Dal'nevostochnoye geologicheskoye upravleniye.

KRYZHANOVSKAYA, T.A.; CHERNYI, G.I., kand.tekhn.nauk; YARMOLYUK, V.T.

Lining mine shafts in conditions of the Belozherka iron ore deposit. Mst. i gornorud. prom. no.2:49-51 Mr-Ap '64. (MIRA 17:9)

YARMOLYUK, V.T.

Wide and narrow face working under conditions of the Volyn'  
coal deposit. Ugol' Ukr. 3 no,12:11-12 D '59.  
(MIRA 13:4)

1. UkrNIIProyekt.  
(Ivov-Volyn' Basin--Coal mines and mining)



PESHIY, V.G., gornyy inzh.; YAEVOLYUK, V.T., gornyy inzh.

Shaft bottom with a high traffic capacity. Ugol' Ukr. 4  
no.8:24-26 Ag '60. (MIRA 13:9)  
(Lvov-Volyn Basin--Mine haulage)

YARMOLYUK, V.T., gornyy inzh.

Effect of seam thickness on the costs of coal mining. Ugol' Ukr.  
5 no.4:44-45 Ap '61. (MIRA 14:4)

1. UkrNIIProyekt. (Coal mines and mining--Costs)

SKOLOZDRA, R.V.; YARMOLYUK, Ya.I.; GLADYSHEVSKIY, Ye.I.

Compounds of the R-phase type in the systems Mo - Fe (Co, Ni) - Si (Ge).  
Zhur. struk. khim. 6 no.3:473-474 My-Je '65. (MIRA 18:8)

1. L'vovskiy gosudarstvennyy universitet imeni Iv. Franko.

GLADYSHEVSKIY, Ye.I.; KRIPYAKEVICH, P.I.; YARMOLYUK, Ya.P.

Crystalline structure of  $\text{Mn}_2\text{Si}$ . Izv. AN SSSR. Neorg. mat. 1 no.7:  
1086-1089 J1 '65. (MIRA 18:9)

1. L'vovskiy gosudarstvennyy universitet imeni I.Franko.

L 46769-66 EWT(m)/T/EWP(t)/FTI LJP(c) JD/HW/JG  
 ACC NR: AP6031724 SOURCE CODE: UR/0370/66/000/005/0148/0151

AUTHOR: Skolozdra, R. V. (L'vov); Gladyshevskiy, Ye. I. (L'vov); Yarmolyuk, Ya. P. (L'vov)

ORG: none

TITLE: Ternary Mo-Co-Si system

47  
B

SOURCE: AN SSSR. Izvestiya. Metally, no. 5, 1966, 148-151

TOPIC TAGS: *COBALT CONTAINING ALLOY, SILICON CONTAINING ALLOY,*  
 molybdenum cobalt silicon system, molybdenum cobalt silicon alloy, alloy  
 phase diagram, alloy phase composition, alloy structure, intermetallic compound,  
*TERNARY ALLOY, MOLYBDENUM CONTAINING ALLOY*

ABSTRACT: A study has been made of 120 alloys of the molybdenum-cobalt-silicon system. Alloys were melted from 99.9%-pure molybdenum, 99.98%-pure cobalt, and 99.99%-pure silicon. A phase equilibrium diagram of the system at 800C (see Fig. 1) was plotted on the basis of data obtained by physicochemical analysis. The existence of MoCoSi compound, with a homogeneity region extending from 15 to 30 at% silicon, was confirmed. Two previously unknown compounds,  $\text{Mo}_5\text{Co}_3\text{Si}_2$  ( $a = 11.06\text{\AA}$ ,  $c = 19.89\text{\AA}$ ) and  $\text{Mo}_3\text{CoSi}$  ( $a = 12.70\text{\AA}$ ,  $c = 4.85\text{\AA}$ ), were found. The solubility of cobalt in MoSi was about 3 at%

Card 1/3

UDC: 669.285'25'782

L 46769-66

ACC NR: AP6031724




Fig. 1. Phase equilibrium diagram of the Mo-Co-Si system at 800C

- 1 - Single-phase alloys;
- 2 - two-phase alloys;
- 3 - three-phase alloys.

Card 2/3

L 46769-66

ACC NR: AP6031724

and in  $\text{Mo}_5\text{Si}_3$  and  $\text{Mo}_3\text{Si}$ , about 5 at%. The solubility of silicon in  $\text{Mo}_6\text{Co}_7$  was 12 at% and in  $\text{MoCo}_3$ , less than 5 at%. The solubility of molybdenum in silicides was found to be insignificant. Orig. art. has: 2 figures and 2 tables. [TD]

SUB CODE: 11/ SUBM DATE: 21Dec64/ ORIG REF: 007/ OTH REF: 008/ ATD PRESS: 5090

hs

*YARMONENKO, S.P.*  
KRASNYKH, I.G.; YARMONENKO, S.P.

Concerning the article by B.M. Graevskaia and R.IA. Keilina "Decrease in the sensitivity of animals to fatal doses of X rays following irradiation with nonfatal doses." Biofizika 2 no.6:764-765 '57.

(X RAYS--PHYSIOLOGICAL EFFECT) (MIRA 10:12)  
(GRAEVSKAIA, B.M.) (KEILINA, R.IA.)



SOV/20-127-5-53/58

17(10).  
AUTHORS:

Yarmonenko, S. P., Kostyanovskiy, R. G.

TITLE:

The Effect of Methyl-bis-( $\beta$ -chloro Ethyl)-amine (HN2) on Frogs Under Hibernation Conditions

PERIODICAL:

Doklady Akademii nauk SSSR, 1959, Vol 127, Nr 5, pp 1125-1127 (USSR)

ABSTRACT:

The chloro ethyl amines, especially HN2 are typically radiomimetic substances (Refs 1,2). They reproduce distinctly the radiobiological effect (Refs 1-8). The analogy in the effect of the radiations and HN2 also concerns the preservatives against the damage caused by them (Ref 9), e.g. the radio preservatives of the mercamine type (Refs 10-14). All that proves the commonness of the concerning mechanisms in certain stages of the biological effect of the two mentioned factors. The authors assume that the radiobiological effect is caused not only by the short-lived radicals ( $HO_2$ , OH, H), but also by more stable intermediate products of the latter. Therefore, they say that HN2 imitates the effect of these intermediate products. In this connection it is important that the radiation- as well as the radiomimetic effect are based upon certain chemical reactions

Card 1/4

SOV/20-127-5-53/58

The Effect of Methyl-bis-( $\beta$ -chloro Ethyl)-amine (HN2) on Frogs Under Hibernation Conditions

the rate of which is bound to depend on the temperature. The radiation disease of frogs (Refs 5-17), mice (Refs 18-21), rats (Refs 18,22), Spermophilus (Refs 23,24), and Myoxis (Refs 25-27) develops extremely slowly at 0-12°. The latent period and mortality characteristic of the concerning dose occur at room temperature (Refs 16,25). The data on the temperature dependence of the alkylating agents are very rare; thorough investigations of the effect on the entire organisms are not known to the authors. In the present paper the intoxication course with HN2 at low temperatures was compared with analogous data on the radiation damage (Ref 16). The experiments were carried out with 100 female and male grass frogs (*Rana temporaria*). The experimental frogs were kept 24 hours before the introduction of HN2 at 0-1°, the control animals put into a thermo-chamber (18-20°). Both groups had a biological control (20 intact frogs each of them, kept under analogous conditions). HN2 was introduced as aqueous hydrochloride solution in doses of 40, 60, and 200 mg/kg into the leg muscles of the experimental animals. The control animals died according to the above doses

Card 2/4

SOV/20-127-5-53/58

The Effect of Methyl-bis-( $\beta$ -chloro Ethyl)-amine (HN2) on Frogs Under Hibernation Conditions

within the following days: 5-12, 1-3 days, 3-6 hours respectively; or 30 days (animals did not die), 5-12, 3-4 days respectively at low temperature. The frogs treated with 40 mg/kg HN2 showed under the control conditions (18-20<sup>o</sup>) after 2 days the symptoms of the radiation disease and died after 3-6 days. Figure 1 shows the dynamics of the dying. No satisfactory explanation exists for the time being for the deceleration of the radiation disease by low temperatures. Most of the research workers assumed a deceleration of the metabolism processes. The authors consider the mentioned phenomenon to be unequivocally clarified: the initially mentioned intermediate products react practically instantaneously with the biosubstrate at usual temperatures. The results obtained agree well with several experimental facts (Refs 16,25,30). There are 2 figures and 30 references, 7 of which are Soviet.

ASSOCIATION: Institut khimicheskoy fiziki Akademii nauk SSSR (Institute of Chemical Physics of the Academy of Sciences, USSR)

Card 3/4

17 (10)  
AUTHORS:

Zherebchenko, P. G., Krasnykh, I. G., SOV/20-129-6-63/69  
Lebkova, N. P., Yarmonenko, S. P.

TITLE:

The Influence of Local Asphyxia of the Bone Marrow on the Course and Result of the Radiation Disease

PERIODICAL:

Doklady Akademii nauk SSSR, 1959, Vol 129, Nr 6, pp 1427 - 1429 (USSR)

ABSTRACT:

The transplantation of the bone marrow of a donor is difficult. Since the sensitivity of individual organs to radiation can be reduced by local asphyxia caused in these organs, the authors tied up the hind legs of test animals. If the lower third of the upper thigh is tied up, complete hemostasia and, consequently, hypoxia occurs in all lower parts, even in the bone marrow of the lower leg. 263 white rats and 503 white mice were investigated. They were exposed to total irradiation of 700, 750, and 800 r (intensity of doses: 49 and 32 r per minute, respectively). A round rubber band was used as a tourniquet which was applied before irradiation and taken off immediately after irradiation. The tourniquet proved to be favorable for the course and results of the radiation damage in all experiments (Table 1).

Card 1/3

The Influence of Local Asphyxia of the Bone Marrow on the SOV/20-129-6-63/69  
Course and Result of the Radiation Disease

40-80% of the mice of the test groups were still alive on the twelfth day (doses: 700-750 r) whereas all control animals died on the 8th - 10th day. Only 10-50% of the mice were alive on the 30th day. The average life period of the test animals was considerably longer than that of the control mice. The results with rats were similar. The fact whether one or two legs had been tied up was not essential for the surviving of test animals. Novocaine was locally used in order to eliminate the effect of functional shifts (due to the pain reaction in applying the tourniquet). The effect of asphyxia was not reduced by this. This effect was determined by counting the degeneratively changed nuclei and the mitotic index on total preparations of mice and the micronecrotic centers of rats (according to M. N. Meysel', Ref 18). Figure 1 shows that local asphyxia considerably decreases the degeneration of the cells of the irradiated bone marrow, and considerably increases their mitotic activity. Rats had about 65% of micronecrotic centers less than the control animals (3 hours after irradiation). Local asphyxia of the bone marrow had no considerable effect on the intensity of leucopenia. The lower degree of bone-marrow injury is probably due to the

Card 2/3

The Influence of Local Asphyxia of the Bone Marrow on the Course and Result of the Radiation Disease SOV/20-129-6-63/69

reduced oxygen concentration. The effect of low metabolism has also to be taken into account. The effect determined combines with the protecting effect of mercamine. It can be maintained that the first effect will be increased by the complex of therapeutic methods used in the treatment of radiation diseases. Mercamine hydrochloride was synthesized by F. Yu. Rachinskiy. There are 1 figure, 1 table, and 19 references, 8 of which are Soviet.

PRESENTED: July 5, 1959, by I. L. Knunyants, Academician

SUBMITTED: July 5, 1959

Card 3/3

KRASNYKH, I.K.; LEBOVA, N.P.; YARMOHENKO, S.P.

Protection in early radiation injuries of the bone marrow.  
Med.rad. 5 no.4:35-3: Ap '60. (MIRA 13:12)  
(RADIATION PROTECTION) (ETHYLAMINE)  
(MARROW)

ZHEREBCHENKO, P.G.; KRASNYKH, I.G.; LEBKOVA, N.P.; YARMONENKO, S.P.

Protective action of local asphyxia of the bone marrow in acute radiation injury in animals. Med.rad. 5 no.10:28-35 '60.

(MIRA 14:2)

(RADIATION SICKNESS)

(MARROW)

(BLOOD--CIRCULATION, DISORDERS OF)



S/205/61/001/006/013/022  
D243/D305

27.2400 also 2209

AUTHOR: Yarmonenko, S.P.

TITLE: Influence of cystamine on endogenous tissue respiration

PERIODICAL: Radiobiologiya, v. 1, no. 6, 1961, 903 - 904

TEXT: The effect of cystamine dichlorhydrate on the endogenous tissue respiration of the spleen and kidney was investigated to try to clarify the mode of action of mercaptoamine radioprotective substances. 150 mg/kg of the preparation were introduced intraperitoneally into 60 sexually mature, healthy mice of both sexes. The mice were decapitated after 30 - 60 hours and the oxygen demands of spleen and kidney extracts over two hour periods determined with a Warburg apparatus. [Abstractor's note: No details given of the apparatus]. In both organs, tissue respiration increased by 20 % on average. The results are discussed. The author concludes that the protective effects of mercaptoamines is the result of interference in the energy exchange processes, the "oxygen effect" being

Card 1/2

32753

S/205/61/001/006/013/022

Influence of cystamine on endogenous ... D243/D305

of prime importance. There are 1 figure, 1 table and 10 references: 6 Soviet-bloc and 4 non-Soviet-bloc. The references to the English-language publications read as follows: B. Bergeret and H. Blashko, Brit. J. Pharmac., 12, 513, 1957; H. Brenk and R. Van-den Moor, Nature, 183, 4624, 1530, 1959. x

SUBMITTED: January 25, 1961

Card 2/2

YARMONENKO, S.P.; LESHKO, Yu.M.; MINEYEV, A.I.

Plastic cages for small laboratory animals. Lab. delo [7] no.4:  
55 Ap '61. (MIRA 14:3)

(LABORATORY ANIMALS)

LETAVET, A.A., prof.; KURLYANDSKAYA, E.B., prof., doktor biol.  
nauk; YARMONENKO, S.P., red.

[Materials on the biological effect of high-energy protons]  
Materialy po biologicheskomu deistviu protonov vysokikh  
energii. Moskva, Akad. med. nauk SSSR, 1962. 116 p.  
(MIRA 17:4)

1. Chlen-korrespondent AMN SSSR (for Letavet).

\*

S/365/62/002/000/037/042  
D405/D301

AUTHORS: Rapoport, I.A., Yarmonenko, S.P. and Avrunina, G.A.  
TITLE: Effect of high-energy protons on frequency of mutations  
SOURCE: Problemy kosmicheskoy biologii. v. 2. Ed. by N. Siskyan and V. Yazdovskiy. Moscow, Izd-vo AN SSSR, 1962, 370-386

TEXT: The study of the effect of high-energy protons on the frequency of mutations is important owing to the fact that protons of energy up to 1000 Mev constitute a major component of cosmic radiation at a radius of 100-1000 km from the Earth's surface, this component being concentrated in particular in the regions of magnetic anomaly. The proton beam was obtained from the synchrocyclotron of the Joint Institute for Nuclear Research. The characteristic of the beam and the method of calculation are described in the references. *Drosophila* males of the line yellow<sup>39</sup> were subjected to proton irradiation. Conclusions: Protons of energy 660 Mev in dose-rates of

Card 1/3

Effect of high-energy ...

S/865/62/002/000/037/042  
D405/D301

510-12140 r bring about an increase in the frequency of recessive lethal mutations in *Drosophila* sex chromosomes that is proportional to the dose, constituting  $\sim 2\%$  per each 1000 r. The curve mutation frequency vs. dose is very similar to the corresponding curve for gamma radiation. The measurement of the frequency of recessive lethal mutations for doses of 3980, 9165, 10745, 11485 and 12140 r during 6 experiments showed a sharp drop in the mutation yield of irradiated spermatogonia as compared to that of irradiated spermatozooids; the main reason for this is apparently the disproportionate rate of reduplication of gonial cells with chromosome aberrations and semi-dominant and dominant lethals with respect to those left normal, and not the low mutability of spermatogonia. On comparing the fertility in the first generation of crossed males, it is found that the frequency of dominant lethal mutations in the irradiated spermatozooids has a similar pattern to that of dominant lethal mutations due to X- and gamma radiation. The curves for the frequency of sterility and semi-sterility mutations are similar to those for other mutations. The fact that the mutability curves retain their linearity when the oxygen concentration increases suggests that, in

Card 2/3

Effect of high-energy ...

S/865/G2/002/000/037/042  
D405/D301

the case of irradiation by high-energy protons, the oxygen is a more effective component of the mutation background than organic peroxides and radicals. There are 3 figures and 5 tables.

Card 3/3

YARMONENKO, S.P.

Fluorescence microscopy of the bone marrow following subjection  
to radiomimetic substances. Med. rad. 4 no.3:52-57 Hr '59. (MIRA 12:7)  
(MARROW) (FLUORESCENCE MICROSCOPY)



ZHEREBCHENKO, P.G.; SUVOROV, N.N.; SHASHKOV, V.S.; YARMOHENKO, S.P.;  
MOROZOVSKAYA, L.M.

Mechanism of the radioprotective action of 5-hydroxytryptophan.  
Radiobiologia 1 no.5:789-791 '61. (MIRA 14:11)  
(RADIATION PROTECTION) (TRYPTOPHAN)

17 (10,12)

## AUTHORS:

Kostyanovskiy, R. G., Yarmonenko, S. P. SOV/20-127-6-42/51

## TITLE:

A Comparative Analysis of the Biological Effect of Ionizing Radiation and of Methyl-bis-( $\beta$ -chloroethyl)-amine (HN2) Within a Large Range of Doses

## PERIODICAL:

Doklady Akademii nauk SSSR, 1959, Vol 127, Nr 6, pp 1294 - 1296 (USSR)

## ABSTRACT:

There is a characteristic dependence of the lifetime of mammals on the irradiation dose (Refs 1,2). Rayewskiy et al distinguish 5 ranges of doses which, in their opinion, reflect independent detrimental mechanisms: they cause the death of test animals (100-1200; 1200-15,000; 15,000-30,000; 30,000-100,000; >100,000 r). In the second of these ranges, the lifetime is independent of the dose, and is 3-5 days on an average. The lifetime is rapidly reduced with an increase in the dose, and at 100,000 r death occurs during irradiation. From 30,000 r onward, spasms and other symptoms of a detrimental effect on the central nervous system occur. At doses of 20,000-50,000 r, an independent detrimental mechanism - the damage of the nerve centers - is said to act in contrast to the "peripheral" or "reflex" mechanism, the latter occurring with smaller doses (Refs 3,4).

Card 1/3

A Comparative Analysis of the Biological Effect of Ionizing Radiation and of Methyl-bis-( $\beta$ -chloroethyl)-amine (HN2) Within a Large Range of Doses SOV/20-127-6-42/51

In connection with the deliberations previously expressed (Ref 5) by the authors concerning the utility of radiomimetic representation of radiobiological effects, they investigated the detrimental effects by doses of from 1 to 2,000 mg/kg of HN2 administered intraperitoneally to 568 white mice in the form of an aqueous hydrochloride solution (0.1-0.5 ml). Figure 1 shows the dependence of the average lifetime on the HN2-dose as a logarithmic curve, as compared to the Rajewskiy-curve (Ref 1). Both curves coincide at one point which corresponds to the minimum, absolutely lethal, doses (750 r and 4 mg/kg). Already a casual comparison of these two curves excludes any doubt about the close relationship of the phenomena represented by them. The blood investigation showed that in the case of HN2-doses lying on the horizontal part of the curve, death occurs in connection with an extensive suppression of blood formation (Table 1). This agrees with the results obtained during the "acute" form of radiation disease (L. F. Semenov, Ref 3). Besides the above analogies in the character of the two curves, also very interesting differences are found. In

Card 2/3

A Comparative Analysis of the Biological Effect of Ionizing Radiation and of Methyl-bis-( $\beta$ -chloroethyl)-amine (HN2) Within a Large Range of Doses SOV/20-127-6-42/51

spite of these differences, the principal similarity of the two said dependences cannot be denied. Even if the complicity of the pathological process caused by the HN2-intoxication is considered, 2 leading detrimental mechanisms can be distinguished which bring about the death: a) In the range of the horizontal part of the curve, the detrimental effect on the rapidly dividing tissues, particularly the blood-forming ones, seems to be decisive; b) In the interspaces b and v, the animals die at symptoms of a detrimental effect to the central nervous system. The 5 interspaces by Rayewskiy only reflect 2 mechanisms of the radiation death. The two mechanisms must, however, not be opposed to each other. There are 1 figure, 1 table, and 7 references, 4 of which are Soviet.

ASSOCIATION: Institut khimicheskoy fiziki Akademii nauk SSSR (Institute of Chemical Physics of the Academy of Sciences, USSR)

PRESENTED: April 6, 1959, by N. N. Semenov, Academician

SUBMITTED: March 31, 1959  
Card 3/3