

Thermodynamics and Structure (Cont.)

SOV/2809

Chulanovskiy, V. M. Spectroscopic Methods for Studying the
Structure of Solutions

251

Belyy, M. U. Spectroscopic Methods for Studying Complexes in
Solution

258

Zelinskiy, V. V., V. P. Kolobkov, and I. I. Reznikova.
Relationship Between Electronic Absorption Spectra and
Radiation of Solutions of Organic Compounds and the Chemical
Nature of Solvents

262

Vaynshteyn, E. Ye., and I. I. Antipova-Karatayeva. Study of
Solvation of Ions in Solutions With the Aid of Optical
Absorption Spectra

266

Antipova-Karatayeva, I. I. Study of the Effect of the
Surrounding Medium on the State of the Chrome Ion by Means
of Absorption Spectra of Solutions and Alum Crystals

270

Card 9/10

Thermodynamics and Structure (Cont.)

SOV/2809

Vasenko, Ye. N., A. P. Chernyavskaya, and N. V. Chernaya.
Infrared Spectra of Electrolytic Solutions in Formamide

273

Levshin, V. L., Ye. G. Baranova, L. D. Derkacheva, and
L. V. Levshin. Study of Association in Concentrated
Solutions of Dyes by Means of Absorption and Luminescence
Spectra

275

Levshin, L. V. Effect of Ionization and Association on
Optical Properties of Complex Organic Molecules

285

AVAILABLE: Library of Congress

Card 10/10

TM/jmr
1-23-60

YUKHNOVSKIY, I.R. [Yukhnovs'kyi, I.R.]

Statistical theory of ionic systems. Ukr. fiz. zhurn. 4 no.2:167-176
Mr-Ap '59. (MIRA 13:1)

I.L'yovskiy gosudarstvennyy universitet.
(Ions) (Coulomb functions)

YUKHNOVSKIY, I.R. [Yukhnovs'kiy, I.R.]; RAKHIMOVA, I.Sh. VIADIMIROV, V.V.
[Vladymyrov, V.V.]

Contribution to the theory of systems of charged particles in an
external field. Ukr. fiz. zhur. 4 no.3:334-344 My-Je '59.
(MIRA 13:2)

I.L'vovskiy gosudarstvennyy universitet im. I. Franko.
(Plasma (Ionized gases))

24(5)

AUTHOR:

Yukhnovskiy, I. R.

SOV/20-126-3-27/69

TITLE:

The Free Energy of a System of Charged Particles (Svobodnaya energiya sistemy zaryazhennykh chastei)

PERIODICAL: Doklady Akademii nauk SSSR, 1959, Vol 126, Nr 3,
pp 557 - 560 (USSR)

ABSTRACT:

In the introduction to the present paper it is pointed out that the calculation of such a system must be carried out by means of a combined method, in which the effect of Coulomb reaction is calculated according to the method of the collective variable (D. N. Zubarev) (Ref 1), and the short interaction by Mayer's method of group integrals. In an already published paper by the same author (Ref 2) a cutoff Coulomb potential was used for a system of various ions. It is intended to be shown in the present paper that a cutoff is unnecessary for ion systems. The equation of free energy is obtained by calculating the thermodynamic quantities and the distribution functions. The author proceeds from a neutral system of ions, which is described by the equation (1). The formula for the free energy is then obtained from the configuration integral of the system by expansion in series, where the "short-range"

Card 1/2

The Free Energy of a System of Charged Particles

SOV/20-126-3-27/69

and the "Coulomb" particles are taken into account. In the case of a high concentration, the "short-range"-, and in the case of a weak concentration the Coulomb forces predominate. Finally, the free energy for weak concentrations is calculated. There are 4 Soviet references.

ASSOCIATION: L'vovskiy gosudarstvennyy universitet im. Ivana Franko
(L'vov State University imeni Ivan Franko)

PRESENTED: January 15, 1959, by N. N. Bogolyubov, Academician

SUBMITTED: January 9, 1959

Card 2/2

YUKHNOVSKIY, I.R. [I.Ukhnovs'kiy, I.R.]

Statistical theory of systems of interacting ions and dipole.particles.
Ukr. fiz. zhur. 6 no.3:333-339 My-Je '61. (MIRA 14:8)

1. Lvovskiy gosudarstvenny universitet im. IYa. Franka.
(Ions)
(Dipole moments)

YUKHNOVSKIY, I.R.

Statistical theory of mixed ion-dipole systems of interacting particles. Dokl. AN SSSR 136 no.6:1317-1320 F '61. (MIRA 14:3)

1. L'vovskiy gosudarstvennyy universitet im. Ivana Franko.
Predstavleno akademikom N.N. Bogolyubovym.
(Ions)
(Dipole moments)

YUKHNOMSKY

I. R.

S/185 36095
D299/D301/007/003/005/015

4500
AUTHOR:

TITLE:

PERIODICAL:

TEXT:

tial ϵ_{ab} and the dipole. Spheric-symmetrical functions are derived for the self-consistent potential and dipoles. On passing to a point distribution, one obtains a generalized expression for the self-consistent free energy F_c of a system of ions.

Ukrayins'kyy fizichnyy zhurnal, v. 1, no. 3, 1962

267 - 276

collective variables

spatial distribution of particles by the method of

integrating the distribution of charges. Collective variables are introduced by means of a Fourier representation for U ; (these variables are introduced by integrating the distribution of charges and dipoles). By in-

tegrating the expression for U with respect to the collective variables

one obtains:

Card 1 | 5

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001963120008-1

YUKHNOVSKYIY I. R.

36095

S/185/62/007/003/005/015
D299/D301

4,4500

AUTHOR:

Yukhnovs'kyy, I.R.

TITLE:

Spatial distribution of particles by the method of
collective variables

PERIODICAL:

Ukrayins'kyy fizychnyy zhurnal, v. 7, no. 3, 1962
267 - 276

TEXT: Formulas are derived for the self-consistent potential G_{ab} and the self-consistent free energy F_c of a system of ions and dipoles. Spheric-symmetrical functions are introduced, characterizing the space distribution of charges. On passing to a point distribution, one obtains a generalized expression for the energy of interaction U of point particles. Collective variables are introduced by means of a Fourier representation for U ; (these variables are Fourier images of the distribution functions of charges and dipoles). By integrating the expression for U with respect to the collective variables P_k , one obtains:

Card 1/5

S/185/62/007/003/005/015
D299/D301

Spatial distribution of particles ...

$$Z = \int \prod d\Omega \exp \left\{ -\frac{\bar{U}(0)}{\Theta} + \frac{1}{2} \sum_k \alpha(k) - \ln(\alpha(k)+1) \right\} \times \\ \times \left(1 + \sum_{a,b} \frac{1}{V^2} N_a N_b \int \delta_{ab} \frac{1}{3!} + \dots \right) dq_1 dq_2 + \dots \quad (9)$$

α_{ab} is the self-consistent potential:

$$\alpha_{ab} = - \sum_k \lambda_a \lambda_b \frac{\alpha(k\Omega)}{\alpha(k\Omega)+1} e^{i k R} \quad (10)$$

An approximate formula is obtained for a self-consistent potential of a space-distribution of charges, viz.:

$$\sum_k \frac{4\pi}{V} \frac{e^{ikr}}{(\alpha(k)+1) k^2} \approx \frac{1}{ikr} \cdot \frac{1}{1+x_3^2} \int \frac{k e^{ikr} dk}{k^2 + \frac{x_3^2(k)}{1+x_3^2}} \quad (11)$$

Card 2/5

S/185/62/007/003/005/015
D299/D301

Spatial distribution of particles ...

if point charges are assumed, then

$$\epsilon_{ab}(R_i - R_j) = -\frac{1}{\theta} \cdot \frac{1}{1 + \chi_d^2} \cdot \frac{\exp -\frac{\chi}{\sqrt{1 + \chi_d^2}} |R_i - R_j|}{|R_i - R_j|}, \quad (19)$$

where $1 + \chi_d^2$ has the meaning of dielectric constant of the medium. Further, the term $\frac{1}{k}$ in Eq. (9), is considered, which represents the self-consistent free energy F_c :

$$\begin{aligned} F_c &= -\frac{\theta}{2} \sum_k [\alpha(k) - \ln(\alpha(k) + 1)] = \\ &= \frac{1}{2} \frac{V\theta}{(2\pi)^3} \int d\Omega_A k^2 dk [\alpha(k\Omega) - \ln(\alpha(k\Omega) + 1)]. \end{aligned} \quad (20)$$

Card 3/5

Spatial distribution of particles ...

S/185/62/007/003/005/015
D299/D301

In order to simplify the computations, schematic operations are introduced. Thereby, formulas are obtained which elucidate the structure of the self-consistent free energy. An approximate formula is obtained:

$$F_c = -\frac{1}{2} \int \frac{g(h^2)}{h^2} \left\{ x^4(1+a^2k^2) + x^2x_d^2 \left[h^2 + \frac{a^2k^4}{2} + \frac{p^2k^4}{2} \right] + \frac{x_d^2p^2h^6}{2} \right\} dk, \quad (26a)$$

which can be integrated, (in Eq. (26a), F_c does not depend on the angles). Further, the group integrals in Eq. (9) are considered. By integration of these integrals, an exponential function is obtained. The free energy can be expressed as the sum of several terms, including the exponents of that function. After transformations, one obtains

$$F_c^i = \theta \sum_k \frac{1}{1 + \chi_d^2} \frac{\chi^2}{k^2} - \ln \left(1 + \frac{1}{1 + \chi_d^2} \frac{\chi^2}{k^2} \right). \quad (33)$$

Card 4/5

S/185/62/007/003/005/015

Spatial distribution of particles ... D299/D301

by setting in Eq. (33), α and β equal to zero, one obtains the exact Debye free-energy of self-consistent field theory with dielectric constant $\xi = 1 + \chi_2^2$. From Eq. (33) it is possible to pass to a point distribution by means of Bogolyubov's method of functional differentiation. From the final expression for the free energy, it is possible to obtain practically all the thermodynamic and static characteristics of the system; in particular, formulas are derived for the mean moment and for the dispersion of moments. The obtained formula for the free energy remains also valid for a system of associated ions. There are 2 references: 1 Soviet-bloc and 1 non-Soviet-bloc. The reference to the English-language publication reads as follows: E.F. Bertout, Phys. Rev., 91, 415, 1953.

ASSOCIATION: L'viv's'kyj derzhuniversytet im. Iv. Franka (L'viv State University im. Iv. Franko)

SUBMITTED: April 4, 1961

Card 5/5

f

L 18249-63

EWTS LIBRARY RECORD NUMBER

ACCESSION NO.: AP3002117

S 10184

AUTHOR: Yukhnovskyy I. R., Nekrot, A. O.

TITLE: Free energy and binary distribution functions of a mixed ion-dipole system.

SOURCE: Ukrains'kyy fizichnyy zhurnal, v. 8, no. 6, 1963, 637-644.

TOPIC TAGS: ion, molecule, dipole, ion-dipole interaction virial coefficient, third virial coefficient, free energy, ion-dipole plasma, binary distribution function

ABSTRACT: This paper is continuation of a study published previously by the authors in the same journal in issue 7, 1962 and issue 8, 1963. A classical system consisting of ions and dipoles distributed in space is taken, and equations leading to the determination of the free energy of this system are developed. The free energy is thus determined accurately up to the third complete virial coefficient. Binary distribution functions are also calculated. Particular cases of these functions are functions for ion systems described by Yukhnovskyy in this journal in issue 4, 1959, and functions for systems of molecules described by

Card 1/2

L 18249-63

ACCESSION NR: AP3002117

N. N. Bogolyubov in the 1946 OTTI publication entitled "Problems of dynamic theory in statistical physics". Orig. art. has: 39 formulas.

ASSOCIATION: L'viv's'ky'y Derzhuniversy'tet im. Iv. Franka
(Lvov State University im. I. Frank)

SUBMITTED: 29 Nov 62

DATE ACQ: 12 Jul 63

ENCL: 00

SUB CODE: NS, PH

NO REF Sov: 008

OTHER: 000

Card 2/2

GLAUBERMAN, A.Yu. [Hlauberman, A.IU.], prof., otv. red.; RYBALKA,
V.V., red.; SEN'KIV, M.T., dots., red.; VISHNEVSKIY, V.N.,
[Vishnev's'kyi, V.N.], dots., red.; YUKHOVSKIY, I.R.
[Iukhovs'kyi, I.R.], dots., red.; PALYUKH, B.M., dots.,
red.; KVITKO, I.S., red.

[Problems in solid state physics] Pytannia fizyky tverdoho
tila. L'viv, Vydz-vo L'viv's'koho univ., 1964. 117 p.
(MIRA 17:11)

L. Lvov. Universytet.

L 16366-65
ACCESSION NR: AP4043092

$\sigma_{\text{eff}} = 0.6 \times 10^{-24}$

ACCESSION NR: AP-4043092
AUTHOR: Yukhnovskiy, I. R. (Yukhnovskiy, I. R.)
and collective variabili

AUTHOR: Yukhnova'ky'y, I. N. Vrabec
TITLE: Statistical operator and collective variabil-
ation - as "Ukrayins'ky'y fizy*chnyy zhurnal."

TITLE: Statistical op-
eration
SOURCE: Ukrayins'kyi fizychnyy zhurnal.
operator, collection

SOURCE: Ukrayinskyj
TOPIC TAGS statistical operator, collective
action, fermion interaction

Ceramic

ACCESSION NR AP4043092

ASSOCIATION: L'vivs'kyy derzhuniv'sytet is.

ENCL: 00

SUBMITTED: 31Oct63

NO REF

SUB CODE: MA, NP

Card 2/2

L 16120-65 EWT(1) IJP(c)

S/0185151

ACCESSION NR: AP4044165

AUTHOR: Yukhnovskiy, L. R. (Yukhnovskiy, L. R.)

TITLE: Statistical quantum sum and collective variables
to collective variables

SOURCE: Ukrayins'kyi fizychnyy zhurnal, 1954, v. 10, no. 1

TOPIC TAGS: statistical quantum sum, collect variables
energy, union gas

ABSTRACT: As continuation of his previous article, the author computes the statistical sum of series obtained by means of collective variables and investigates its properties (D. N. Zabarev, DAN SSSR 45, #4(1954)). The series is derived for low and high temperatures. The result is a series, the first term of which gives the average energy. This art. has 54 equations

Card 1/2

L 16120-65
ACCESSION NR: AP4044165

ASSOCIATION: L'viv's'kyy derzhuniversytet im. I. I.

SUBMITTED: 15Nov63 ENCL: 00

SUB CODE: NP, MA NO REF Sov: 00

Card 2/2

L 38099-65

ACCESSION NR. AP3005908

operator of two-fermion interaction. The two terms for small values of k , and the diagrams for $k \rightarrow 0$ of the terms. Consequently the diagrams for the expansion also show that the small parameter for the expansion is Brueckner parameter r_B , not $r_B^{1/3}$. The exchange diagrams are calculated. The expression obtained for the diagrams are calculated. The expression obtained for the second-order diagrams in the series, is

$$R = \frac{221}{27} - \frac{0.016}{r_B} + \frac{0.016}{r_B^2},$$

as compared with the expression

$$R = \frac{221}{27} - \frac{0.016}{r_B} + \frac{0.016}{r_B^2},$$

given by M. Gell-Mann and K. A. Brueckner [Phys. Rev., 85, 1650, 1952]. This article has 3 figures and 16 references.

Card 2/3

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001963120008-1

L 38094-67

ACCESSION NR: AF5005908

ASSOCIATION: Lviv'skyj derzhuniversitet
(Lviv State University)

SUBMITTED: 14Feb64

INCL: 75

NR REF Sov: 604

OTRIR: 105

Conf. 3/3

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001963120008-1"

YUKHNOVSKIY, I.R.; NEKROT, A.A.

Some properties of binary functions of distribution of a
mixture of ions and dipole molecules. Ukr. fiz. zhur. 9
no.4:365-375 Ap '64. (MIRA 17:8)

1. Lvovskiy gosudarstvennyy universitet.

YUKHNOVSKIY, I.R., [IUkhnov's'kyi, I.R.]; TSYGANENKO, V.V. [TSyhanenko, V.V.];
VAVRUKH, M.V.

Mean energy of electron gas at absolute zero. Ukr. fiz. zhur. 10
(MIRA 18:4)
no.2:135-146 F '65.

1. L'vovskiy gosudarstvennyy universitat.

YUKHNOVSKIY, I.R. [Ukhnova'kyi, I.R.]

The statistical operator and collective variables. Ukr. fiz. zhur.
9 no.7:702-714 Jl '64. (MIRA 17:10)

1. Lvovskiy gosudarstvennyy universitet im. I. Franko.

YUKHNOVSKIY, I.R. [IUkhnov's'kyi, I.R.]

Quantum statistical sum and collective variables. Part 2.
Transition function to collective variables. Ukr. fiz. zhur.
9 no.8:827-838 Ag '64. (MIRA 17:11)

1. L'vovskiy gosudarstvennyy universitet im. I. Franko.

YUKHNOVSKIY, P.M.

Following Il'ich's advice, Sakh.prom. 36 no.4;6-9 Ap '62.

(MIRA 15:5)

1. Byvshiy presedatel' Tsentral'nogo komiteta soyuza rabochikh
sakharinoj promyshlennosti.
(Sugar industry)

S/058/53/000/002/012/070
A059/A101

AUTHOR: Yukhnovs'kyj, I. R.

TITLE: On the calculation of the quantum statistical sum of a system of charged particles

PERIODICAL: Referativnyy zhurnal, Fizika, no. 2, 1963, 6, abstract 2B44
("Visnyk L"vivs"ka. un-tu.Ser. fiz.", 1962, no. 1(8), 50 - 52,
Ukrainian).

TEXT: The quantum statistical sum is calculated for a system of interacting charged particles having free charges, in the general case. Difficulties connected with the presence of Laplacians appearing in the Hamiltonian in the quantized case are partly removed by way of applying the unitary displacement transformation. This procedure permits the elimination of operators in the exponents of integrands. The expression for the statistical sum is represented in the form of contour integral. The quantum corrections are investigated.

[Abstracter's note: Complete translation]

Card 1/1

S/058/63/000/002/013/070
A059/A101

AUTHOR: Yukhnovs'kyy, I. R.

TITLE: Application of the method of collective variables to calculating
the quantum statistical sum

PERIODICAL: Referativnyy zhurnal, Fizika, no. 2, 1963, 6, abstract 2B45
("Visnyk L"vivs"k. un-tu. Ser. fiz.", 1962, no. 1(8), 63 - 72,
Ukrainian).

TEXT: The expression for the statistical sum obtained in the previous
paper (abstract 2B44) is examined by introducing collective variables and func-
tional differentiation.

[Abstracter's note: Complete translation]

Card 1/1

YUKHNOVSKIY, Ye.

Expensive self-initiated activity. Grashd. av. 20 no. 1:21
(MIRA 16:4)
Ja '63.

1. Lineynaya ekspluatatsionno-remontnaya masterskaya
Khabarovskogo aeroporta.

(Aeronautics, Commercial--Management)

YUKHNOVSKIY, Yu.M.; PIRIAZEV, D.I.; VOLCHEK, F.R.

Rapid rolling conditions on blooming mills at the "Azovstal'" plant. Sbor. trud. UNIIM no.9:186-195 '64 (MIRA 18:1)

MELESJKO, A.M.; TVALICH, K.N.; YUKHNOVSKIY, Yu.M.

Studying the forward flow on continuous sheet rolling mills.
Met. i gornorud. prom. no.4:43-45 Jl-Ag '65. (MIRA 18:10)

ACC NR: AR6035036

SOURCE CODE: UR/0058/66/000/008/B007/B007

AUTHOR: Nevidoms'ka, L. A.; Tokar, S. S.; Yukhnovs'kyy, I. R.

TITLE: Chemical potential of a system of charged particles and the potential's parametric curves accurate precision to the second virial coefficient

SOURCE: Ref. zh. Fizika, Abs. 8B70

REF SOURCE: Visnyk L'vivs'k. un-tu. Ser. fiz, no. 2, 1965, 19-26

TOPIC TAGS: ~~charged particle system~~, charged particle, particle physics, chemical potential, ELECTROLYSIS, SOLUTION CONCENTRATION

ABSTRACT: The chemical potential is calculated for a neutral system of charged particles. Graphs for the relationship between the coefficient of electrolysis activity and the solution concentration are numerically plotted. [Translation of abstract] [NT]

SUB CODE: 20/

Card 1/1

YUZHIC
GALAUKO, A.A.; YUKHO, I.A.; MURHEU, A., redaktor; KALECHYTS, G., tekhnicheskiy
redaktor.

[The local soviets are the organizers of collective farm production;
work practices of local soviets of White Russia (1953-1956)]
Miastaovya Sovety organizatory kahnasni vytvorchastei; z vopystu
raboty minstsovykh Sovetau Belarusi (1953-1956 hh). Minsk,
Dziarzh.vyd-va BSSR, 1957. 194 p. (MIRA 10:11)
(White Russia--Soviets) (White Russia--Collective farms)

GANUSHCHAK, N.I.; YUKHOMENKO, M.M.; STANICHUK, M.D.; DOMBROVSKIY, A.V.

Haloarylation of unsaturated compounds with aromatic diazo
compounds. Part 18:Chloroarylation of diisopropenyl. Zhur. ob.
khim. 34 no.7:2238-2243 Jl '64 (MIRA 17:8)

1. Chernovitskiy gosudarstvennyy universitet i Leningradskiy
tekhnologicheskiy institut imeni Lensoveta.

ZOLOTUKHINA, K.G.; GANUSHCHAK, N.I.; YUKHOMENKO, M.M.; DOMBROWSKIY, A.V.

Tertiary amines and quaternary salts based on 4-chloro-1-aryl-2-butenes
of secondary and tertiary heterocyclic nitrogen bases. Zhur. ob. khim.
33 no.4:1222-1227 Ap '63. (MIRA 16:5)

1. Chernovitskiy gosudarstvennyy universitet.
(Amines) (Heterocyclic compounds)

YUKHOMENKO, M.N.; GANUSHCHAK, N.I.; DOMBROVSKIY, A.V.

Synthesis of 1-arylbuten-2-yl diethylmalonic esters and 1-arylbuten-2-ylacetic acids from chlorobaryl butenes and sodium malonic ester.
Zhur. ob. khim. 33 no.8:2528-2532 Ag '63. (MIRA 16:11)

1. Chernovitskiy gosudarstvennyy universitet.

GANUSHECHAK, N.I.; YUKHOMENKO, M.M.; ROZVAGA, R.I.; DOMBROVSKY, A.V.

Syntheses based on diene condensation. Part 3: 2-Methyl-arylyl-
and 2,3-dimethyl-arylanthraquinones. Zhur. ob. khim. 34 no.8:
2718-2721 Ag '64. (MIRA 17:9)

1. Chernovitskiy gosudarstvenny universitet.

YUKHOMENKO, M.M.; GANUSHCHAK, N.I.; DOMBROVSKIY, A.V.

Synthesis of 1-arylbuten-2-ylacetyleacetones. Ukr. khim. zhur.
30 no.6:616-618 '64. (MIRA 18:5)

1. Chernovitskiy gosudarstvennyy universitet.

I 06504-67 EWP(j)/EWT(m) RM
ACC NR: AP7000488

SOURCE CODE: UR/0079/66/036/006 /1153 /1153

AUTHOR: Ganushchak, N. I.; Yukhomenko, M. M.; Stadnichuk, M. D.; Shevchuk, A. I.

ORG: Chernovitskiy State University (Chernovitskiy gosudarstvennyy universitet);
Leningrad Technological Institute im. Lensoveta (Leningradskiy tekhnologicheskiy
institut)

TITLE: Synthesis of certain phosphonium salts and 1,5-diphenylpentadienes-1,3 on
the basis of chloroarylbutenes

SOURCE: Zhurnal obshchey khimii, v. 36, no. 6, 1966, 1150-1153

TOPIC TAGS: organic phosphorus compound, organic salt, organic synthetic process

ABSTRACT: The reaction of a number of chloroarylbutenes with triphenylphosphine yielded new triphenyl-(1-aryalkenyl-2)-phosphonium chlorides
 $[ArCH_2C(R)+C(R')CH_2P(C_6H_5)_3]Cl^-$. The phosphonium salts were converted to the

corresponding 1,5-diphenylpentadienes-1,3 by reaction with sodium ethylate and benzaldehyde. The infrared and nuclear magnetic resonance spectra of the products were studied. The diphenylpentadienes are oily, yellowish liquids, which are readily soluble in the usual organic solvents, decolorize bromine water and permanganate solution. They do not take part in diene synthesis reactions, even with such dienophiles as maleic anhydride with heating.

Orig. art. has: 2 figures and 1 table. [JPRS: 37, C23]

SUB CODE: 07 / SUEN DATE: 03Jun65 / ORIG REF: 010

UDC: 547.341

Cord 1/1 m/e

0923 1203

~~YUKHOV, I. starshiy prepodavatel'~~

Errors in radar tracking during rolling. Mor. flot 18 no.426
(MIRA 12:12)
Ap '58.

1. Akademiya imeni Krylova.
(Radar in navigation)

ACC NR: AM6036119

(N)

Monograph

UR/

Skvortsov, Mark Ivanovich; YUkhov, Ivan Vasil'yevich; Zemlyanov, Boris Ivanovich;
Abchuk, Vladimir Avramovich; Mrykhin, Oktyabr' Aleksandrovich

Principles of ship maneuvering (Osnovy manevrirovaniya korabley) Moscow,
Voyenizdat M-va obor. SSSR, 1966, 269 p. illus., bibli., 1 fold chart. Errata
slip inserted. Number of copies printed not given.

TOPIC TAGS: naval operation, marine engineering, ship navigation, naval tactic

PURPOSE AND COVERAGE: This book is intended for naval officers and students of
naval schools; it can be also used by the scientific and engineering staffs of
research institutes and the marine industry. Problems of ship navigation,
handling, and maneuvering at sea are discussed with particular application to
military purposes, such as approach to target or changing position of the ship
in relation to some specific object. Theories of probability, detection, and
errors are used extensively in the text, particularly for the theoretical and
practical analysis of problems of maneuverability. There are 16 references, all
Soviet.

TABLE OF CONTENTS [abridged]:

Introduction -- 3

Card 1/2

UDC: 359:656.61.052

ACC NR: AN6036119

- Ch. 1. Principles of the general theory of ship maneuverability -- 9
Ch. 2. Sea search for enemy -- 38
Ch. 3. Determining elements of target motion -- 91
Ch. 4. Changing distance and position in relation to an object -- 133
Ch. 5. Special cases of maneuvering by a single ship -- 170
Ch. 6. Principles for the evaluation of maneuvering accuracy -- 196
Ch. 7. Joint maneuver of ships -- 209
Ch. 8. Solving maneuvering problems with electronic computers -- 234
Appendix. Maneuvering tables (19 tables) -- 252

Bibliography -- 266

SUB CODE: 15,17/

SUBM DATE: 03May66/

ORIG REF: 027/

OTH REF: 001/

Card 2/2

8(5)

SOV/105-59-3-18/27

AUTHOR:

Yukhov, V. V., Engineer (Chelyabinsk)

TITLE:

Synchronous Motor Compounding With Directly Connected Exciter
(Kompaundirovaniye sinkronnykh dvigateley pri glukhom pod-
klyuchenii vozbuditelya)

PERIODICAL:

Elektrichestvo, 1959, Nr 3, pp 82 - 85 (USSR)

ABSTRACT:

Synchronous motors of the type MS-322-12/6 (2500 kva, 1920 kw, 6 kv, 1000 revs/min) with excitors of the type PN-40 (65 v, 320 a) or synchronous motors of the type MS-322-8/6 (1780 kva, 1350 kw, 6 kv, 1000 revs/min) with excitors of the type MP-543/2/5 (65 v, 360 a) are generally used to drive rotary pumps of the type 22-NDS. Circuits with conventional "magnetic stations" (magnitnaya stantsiya) are provided for starting and controlling. Test runs showed that such circuits are unreliable. For this reason they were re-designed to operate as direct starters with a direct connection of the exciter. These simplified units are notable for a manually regulated compounding device connected directly to the motor circuits. The compounding is carried out either according to the principle

Card 1/3

Synchronous Motor Compounding With Directly Connected
Exciter

SOV/105-59-3-18/27

ple used with generators, or by using the manually regulated exciter current, in which case the current passing through the excitation controller depends upon the compounding current. In order to vary the stability domain of the compounding unit, the connecting transformer is equipped with a movable winding, which is fed by the compounding current. Engineer A. V. Petukhov took part in the experiments. The advantages offered by the new starting device are as follows: 1) The circuits for starting, protection and control are very simple. 2) Powerful inertialess compounding devices with a uniform sensitivity markedly increase the dynamic stability of synchronous motors. 3) The use of the compounding unit accelerates the synchronization of the motors under load. 4) Such an installation somewhat accelerates the field degeneration in motors with a direct connection of the exciter and eliminated the possibility of inverting the sense of magnetization during starting operations. 5) If the parameters of the compounding unit are chosen correspondingly, there arises the possibility of abandoning a forced excitation by means of a relay. This unit, which operates with a compounding making

Card 2/3

Synchronous Motor Compounding With Directly Connected
Exciter

SOV/105-59-3-18/27

use of the current passing through the excitation controller, exhibits still other advantages; 6) The system for the excitation of the motor is independent of the voltage transformer circuits and can be used for a forced excitation of motors operating at rated supply voltage under shock load. 7) Under otherwise equal conditions the motor excitation current is less dependent upon the operational voltage fluctuations and upon the temperature of the exciter windings of the motor and of the exciter. The General Specification Nr E-5/54 of the Tekhnicheskoye upravleniye MES (Technical Administration of the MES) limits the field of application of units with a direct connection of the exciter machine by the motor load at the end of starting up and by the rate of field deterioration. Protests against this specification are advanced and their checking is required. There are 3 figures and 2 tables.

SUBMITTED: April 18, 1958

Card 3/3

8(5)
AUTHORS:

Yukhev, V. V., Engineer,
Petukhov, A. V. Engineer (Chelyabinsk)

sov/105-59-1-19/32

TITLE:

The Starting of a Compound Synchronous Motor in a Direct
Connection of the Exciter

PERIODICAL:

Elektrichestvo, 1959, Nr 11, p 81 (USSR)

ABSTRACT:

The synchronous motor of type MS-322-8/6 (1780 kva, 1350 kw,
1000 rpm) with the exciter of type MP-543/2/5 (65 v, 360 a)
provided for the drive with a centrifugal pump of type
22-NDS was reconstructed and provided with a starting circuit
with indirect connection of the exciter. In the present paper
experimental results are given on the automatic switching on
of a spare aggregate and of the starting of the motor after
the interruption of current supply. For all starting conditions
to which the motor was subject in the experiments the voltage
at the supply line were $0.86 U_n$ (U_n = rated voltage) at the
beginning, amperage $5.6 I_n$ (I_n = rated current) and the power
 $1.15 R_n$ (R_n = rated power). After a reconnection following an
interruption period of 3 or 2.45 seconds the voltages were

Card 1/2

The Starting of a Compound Synchronous Motor in
a Direct Connection of the Exciter

SOV/105-59-11-19/32

0.34 and 0.43 $\frac{U_n}{n}$ respectively. The further results of these experiments are summarized in table 1. In conclusion it is said that this circuit diagram guarantees not only a starting under load but also a starting after an interruption of the current supply. There are 1 table and 1 Soviet reference.

SUBMITTED: April 18, 1958

Card 2/2

YUKHOV, V.V., inzh. (Chelyabinsk)

Calculation of systems for compounding synchronous motors.
Elektrичество no.2:40-45 F '61.
(Electric motors, Synchronous) (MIRE 14:3)

YUKHOV, V.V., inzh.; KUZNETSOV, V.F., inzh.


Automatic device for regulating the voltage of storage batteries.
(MIRA 15:8)
Elek. sta. 33 no.8;85-86 Ag '62.
(Voltage regulators) (Storage batteries)


YUKHOVITSKIY, A.A.

"Application of Radio-active Isotopes in Solving Diffusion in Metals Theory Problems," A.A. Yukhovitskiy, M.E. Yanitskaya, Sotkov, A.D., Moscow, USSR

Paper submitted for presentation at the International Conference on Radioisotopes in Scientific Research, Paris, 9-20 Sep 1957.

Moscow Steel Inst, Min. Higher Education, Moscow USSR





YUKHTANOV, D. M.

ca


Melting difficultly fusible charges in reverberatory furnaces. D. M. Yukhtanov. Tsvetnoy Metal. 13, No. 12, 47-51 (1987). *Tsvetnoye i metalloobrabotka* 42, 659-660. The use of difficultly fusible charges lowers the efficiency of metalurgical furnaces and increases the fuel consumption. These disadvantages can be overcome by incorporating with the charge a small proportion of a powerful fluxing agent, fluor spar, Na and K salts, etc., which lower the melting point of the slag. A. Panineau-Couture

338-113 METALLURGICAL LITERATURE CLASSIFICATION

YUKHTANOV, D'M
CA

The influence of sodium oxide on the properties of slags of Karakpal copper smelter. D. M. Yukhnov, Tsvetnoy Metal', 1939, No. 10-11, 118-22. Typical composition of the Karakpal smelter slags is approx. SiO₂ 55, FeO 20, CaO 8 and Al₂O₃ 10%. The results of lab. expts. conducted with the purpose of lowering the melting temp., and viscosity of slags led the author to the following conclusions: (1) Introduction of alkali (Na₂O) into the charge lowers considerably the melting temp. of the slag. Addn. of 2% Na₂O lowers the melting temp. by about 100°. (2) The viscosity of the smelter slags, 420 poises at 1250°, is lowered to 120 poises by the introduction of 2% Na₂O. (3) The lowering of melting temp., and the increase of fluidity result in economy of fuel and in the decreased losses of Cu in the slags. Further work in the lab. and at the smelter is in progress with the purpose of detg. the optimum conditions of smelting with the use of Na₂O. 14 references. B. N. Daniloff

B. N. Daniloff



YUKHTANOV, Dmitriy Mikhaylovich

Cand. Technical Sci.

Deputy Dir., State Sci. Res. Inst. Nonferrous Metals, -1951-.

"Hydrometallurgy," Moscow, 1949.

Stalin 3rd Prize, 1950, publication.

YUKHTANOV, D.M.

DEMIE, G.V.; KAYVANOV, L.S.; SAKHANSKIY, N.A.; STERNIN, I.M.; YUKHTANOV,
D.M., kandidat tekhnicheskikh nauk, redaktor; PETROVA, N.S.,
tekhnicheskiy redaktor

[High-speed smelting in a reverberatory furnace; experience of
skilled workman A.A. Iarusev] Skorostnaya plavka v otzashatel'nykh
pechakh; opyt mastera A.A. Iaruseva. Moskva, Gos. nauchno-tekhn.
izd-vo lit-ry po chernoi i tsvetnoi metallurgii, 1952. 68 p.
(MIRA 9:12)
[Microfilm]

1. Russia (1923- U.S.S.R.) Ministerstvo tsvetnoy metallurgii.
Tekhnicheskoye upravleniye. TSentral'nyy institut informatsii.
2. Zamestitel' direktora instituta Gintsverstsent (for Yukhtanov)
(Smelting furnaces)

TUKHTANOV, Dmitriy Mikhaylovich; Suvorovskaya, N.A., redaktor;
~~Mikhaylova, V.V.~~, tekhnicheskiy redaktor.

[Production of selenium and tellurium] Proizvodstvo selen'a i
tellura, Moskva, Gos. nauchno-tekhn. izd-vo lit-ry po chernoi
i tsvetnoi metallurgii, 1955, 95 p. (MLRA 8:8)
(Selenium) (Tellurium)

AUTHOR

Yukhtanov, D.M.

TITLE

Production of Selenium From Anodic Slurries (Proizvodstvo selena iz anodnykh shlamov) *Bul. tsvetn. metallurgii*, 1957, Nr 9, pp 29-33
Translation from: Referativnyj zhurnal, Metallurgiya, 1958, Nr 6, p 114 (USSR)

PERIODICAL

A description of the technology employed at the Shen'yan metallurgical plant (China) in the production of Se from anodic slurries by means of sulfatization in a bubbler unit. Anodic slurry containing 11.5% Cu, 5.0% Ni, 1.3-1.4% Se, 0.14% Te, 3.0% As, and 10.0% Sb serve as the raw material from which the Se is extracted. The process involves the following operations: 1) sulfatization of anodic slurries in the sublimation products; 2) collection of SeO₂ in bubbler units; 3) collection of selenious acid with its subsequent acidification with H₂SO₄; 4) reduction to elemental Se through the formation of SO₂. The process is carried out in two cast iron vats filled with H₂SO₄ which is heated to a temperature of 80°C. Anodic

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001963120008-1"

Card 1/2

137-58-6-11988

Translation from: Referativnyy zhurnal, Metallurgiya, 1958, Nr 6, p 114 (USSR)

AUTHOR: Yukhtanov, D.M.

TITLE: Production of Selenium From Anodic Slurries (Proizvodstvo selena iz anodnykh shlamov)

PERIODICAL: Byul. tsvetn. metallurgii, 1957, Nr 9, pp 29-33

ABSTRACT: A description of the technology employed at the Shen'yan metallurgical plant (China) in the production of Se from anodic slurries by means of sulfatization in conjunction with collection of SeO_2 in a bubbler unit. Anodic slurries containing 11.5% Cu, 5.0% Ni, 1.3-1.4% Se, 0.14% Te, 3.0% As, and 10.0% Sb serve as the raw material from which the Se is extracted. The production of Se involves the following operations: 1) sulfatization roasting of anodic slurries; 2) sublimation of SeO_2 from the sulfatization products; 3) collection of furnace gases to be employed in the sublimation of SeO_2 in bubbler units, formation of a solution of selenious acid with its subsequent reduction to elemental Se through the formation of SO_2 ; 4) distillation of the raw Se. Sulfatization is carried out in two cast iron vats filled with H_2SO_4 which is heated to a temperature of 80°C. Anodic

Card 1/2

137-58-6-11988

Production of Selenium From Anodic Slurries

slurries containing 30-35% of moisture are poured into the vats under constant stirring. After all of the slurries have been transferred, the temperature is raised to 200-250° while the mixture is constantly stirred. The process of sulfatization requires 4-5 hours. The sublimation of SeO_2 is performed in a furnace lined with fireclay brick. Optimal condition for the operations are as follows: temperature, 400-500°; duration of the process, 4-5 hours; thickness of the layer, 33 mm. Up to 93-95% of Se are sublimated under these conditions. After sublimation only 0.05-0.1% of Se is contained in the residue. Gases from the furnace are captured in three bubbler units connected in series. Raw Se, containing 96-97% of Se, is obtained in these units. The raw Se is refined in a cast iron retort with dimensions of 150x200x500 mm installed in an electrical furnace with Nichrome windings. The process requires 6-9 hours at a temperature of 640°. The yield of Se amounts to 85-90%. The distillation residue is returned to the sulfatization stage. Taking into account all losses and the distillation residue that is returned from the retort for additional processing, the extraction of Se from anodic slurries constitutes 88-90%. Advantages of the technological procedure described are pointed out. 1. Selenium--Production 2. Selenium--Sublimation 3. Selenium oxides--Sublimation 4. Waste gases--Applications 5. Sulfuric acid--Applications

G.S.

Card 2/2

AZOS, S.; AREP'YEV, A.; ARTAMONOV, I.; BABINA, I.; BIRBOGOVSKIY, V.; BLOZHKO, V.; BRAVERMAN, A.; BYKHOVSKIY, Yu.; VINOGRADOVA, M.; GALANKINA, Ye.; GIL'DENGERSH, F.; GLOBA, T.; GREGVIE, N.; GORDON, G.; GUL'DIN, I.; GULYAYEVA, Ye.; GUSHCHINA, I.; DAVIDOVSKAYA, Ye.; DAMSKAYA, G.; DERKACHEV, D.; YEVDOKIMOVA, A.; YEGUNOV, V.; ZABELEVSHINSKIY, I.; ZAYDENBERG, B.; AZMOSHNIKOV, I.; ITKINA, S.; KARCHEVSKIY, V.; KLUSHIN, D.; KUVINOV, Ye.; KUZNETSOVA, G.; KURSHAKOV, I.; LAKERNIK, M.; LEYZEROVICH, G.; LISOVSKIY, D.; LOSKUTOV, F.; MALEVSKIY, Yu.; MASLYANITSKIY, I.; MAYANTS, A.; MILLER, L.; MITROFANOV, S.; MIKHAYLOV, A.; MYAKINENKOV, I.; NIKITINA, I.; NOVIN, R.; OGNEV, D.; OL'KHOV, N.; OSIPOVA, T.; OSTRONOV, N.; PAKHOMOVA, G.; PETKER, S.; PLAKSIN, I.; PLETENIKOVA, H.; POPOV, V.; PRIESS, Yu.; PROKOF'YEVA, Ye.; PUGHKOV, S.; REZKOVA, F.; RUMYANTSEV, M.; SAKHAROV, I.; SOBOL', S.; SPIVAKOV, Ya.; STRIGIN, I.; SPIRIDONOVA, V.; TIMKO, Ya.; TITOV, S.; TROITSKIY, A.; TOLOKONNIKOV, K.; TROFIMOVA, A.; YEDOROV, V.; CHIZHIKOV, D.; SHEYN, Ya.; YUKHTANOV, D.

Roman Lazarevich Veller; an obituary. TSvet. mat. 31 no.5:78-79
(NIRA 11:6)
My '58.
(Veller, Roman Lazarevich, 1897-1958)

AUTHORS:

Yukhtanov, D.M. and Burovoy, I.A. SOV/136-58-6-1/21

TITLE:

Production Automatic Process Units is the First Problem
in the Development of Integrated Automation of Production
(Sosdaniye avtomatizirovannykh tekhnologicheskikh
agregatov - pervoocherednaya zadacha v razvitiu kon-
pleksnoy avtomatizatsii proizvodstva)

PERIODICAL: Tsvetnyye Metally, 1958,³¹ Nr 6, pp 1 - 4 (USSR)

ABSTRACT: On May 12-16, 1958, an All-Union conference of industrial workers was held to discuss the development of integrated mechanisation and automation of production processes and the increased production of instruments and automation equipment. The authors suggest that efforts must now be made to progress from the automation of individual units to that of automation of complete plants and then works and to the automation of process control. They discuss the economic basis for the selection of priorities in automating, on the example of the copper industry where raw-material costs account for 60-65% and labour costs 15-17% of the total. Ancillary operations represent a fruitful field both for mechanisation and automation. The authors urge that the automation of units should be thoroughly studied and effected using as much proved

Card 1/2

SOV/136-58-6-1/21
Production Automatic Process Units is the First Problem in the
Development of Integrated Automation of Production

equipment and methods as possible. For shaft furnaces, efforts should be concentrated on a single experimental furnace. Leading technologists and designers should co-operate in automation work because of the importance of the plant aspect. Continuity of processes is of great importance for automation; therefore, regulating devices and direct measuring methods and automatic analysers (such as the automatic polarograph produced by K.B. Tsvetmetavtomatika, or the Gintsvetmet flame spectrophotometer), should be perfected. Correct design of automatic systems must also be studied. The authors conclude by urging that all non-ferrous metallurgical workers should concentrate their attention on automation.

Card 2/2

SOV/136-59-7-17/20

AUTHOR: B.N.S.

TITLE: Conference on Autoclave Processes

PERIODICAL: Tsvetnyye metally, 1959, Nr 7, pp 84-87 (USSR)

ABSTRACT: On 23-26 February 1959 a conference was held in Moscow for summing-up and coordinating work on autoclave processes in the metallurgy of heavy, non-ferrous, rare and noble metals. The conference was convened jointly by the Otdel Tsvetnoy metallurgii (Non-Ferrous Metallurgy Department) of the Gosplan SSSR (USSR), the GNTK of the USSR, the NTO for Non-Ferrous Metallurgy and the Gintsvetmet. The conference was opened by N.S. Seliverstov, GNTK of the USSR, who noted that applications of research on autoclave processes was being hampered in the USSR by lack of coordination. The conference heard reports as follows: D.M. Yukhtanov, Gintsvetmet, on progress throughout the world on the use of hydrometallurgical, particularly autoclave, methods for non-ferrous and rare metal production; G. N. Dobrokhotov, Gipronikel', on nickel leaching practice at some Soviet works; N. I. Onuchkina and G. N. Dobrokhotov

Card 1/5

Conference on Autoclave Processes

SOV/136-59-7-17/20

on the thermodynamics and kinetics of the selective reduction by hydrogen and carbon monoxide under pressure of nickel and cobalt from solution; I. Yu. Leshch and K. M. Shelepova, Gipronikel', on design decisions on the application of the flowsheets dealt with by G. N. Dobrokhotov at the Yuzhuralnikel' and Severonikel' Combines and the Ufaleyevskiy (Ufa) Nickel Works; I. N. Maslenitskiy, Leningradskiy gornyy institut (Leningrad Mining Institute) on the advantages of a combined flotation-autoclave method for nickel-electrolysis of slimes containing platinum-group metals; V. B. Zhilkin, Severonikel' combine, and S. I. Sobol', Gintsvermet, on the essentials of the neutral method of oxidizing leaching of nickel concentrate from converter-matte flotation; S. I. Sobol' on preliminary investigations on the development of a sulphurous-sulphuric method for leaching nickel and cobalt from oxidized nickel ores; N. N. Maslenitskiy, Mekhanobr, on the main results of investigations of the autoclave-soda process for treating tungsten-ore beneficiation products;

Card 2/5

SOV/136-59-7-17/20

Conference on Autoclave Processes

V. I. Poprukaylo, Mekhanobr, and D. A. Malakhov, Skopinskaya (Skopinsk) TsOF, separately, on problems in the application of an autoclave-soda flowsheet to scheelite and wolframite raw material; G. A. Meyerson, K. Ya. Shapiro, N. N. Khavskiy, R. A. Pavlyuk and A. P. Nadol'skiy, Krasnoyarskiy institut tsvetnykh metallov (Krasnoyarsk Non-Ferrous Metals Institute) on the treatment of tungsten concentrates in hermetical, heated ball-mills with acids or caustic alkalies; V. I. Spiridonova, S. I. Sobol', Ye. I. Gulyayeva, Z. L. Berlin, I. M. Nelen' and B. I. Rudenko, Gintsvetmet, on the treatment of prepared and unprepared sulphide molybdenum raw material by oxidizing autoclave alkaline leaching; I. M. Nelen' and S. I. Sobol' on the kinetics of oxidizing autoclave leaching; A. N. Zelikman and Z. M. Lyapina, Krasnoyarsk Non-Ferrous Metals Institute, on the results of a study of conditions for the selective separation of lower oxides of tungsten and molybdenum from their salt solutions by hydrogen under pressure; M. V. Darbinyan, Gorno-metallurgicheskiy institut (Mining-Metallurgical Institute)

Card 3/5

SOV/136-59-7-17/20

Conference on Autoclave Processes

of the Sovnarkhoz (economic council) of the Armyanskaya SSR (Armenian SSR), on his investigations of ammoniacal autoclave leaching under oxygen pressure of molybdenum concentrates; S. I. Sotol' on technical-economic factors of ammoniacal leaching; A. I. Sinel'nikova and I. N. Plakshin, Krasnoyarsk Non-Ferrous Metals Institute, on an oxidizing autoclave process for gold-containing raw material; N. G. Tyurin, Ural'skiy politekhnicheskiy institut (Ural Polytechnic Institute) on the behaviour of noble metals in oxidizing autoclave leaching in thiosulphate solutions; A. L. Tseft and D. A. Taraskin and A. Yu. Dadabayev. Institut metallurgii i obogashcheniya AN Kaz SSR (Metallurgy and Beneficiation Institute of the AS Kaz SSR), respectively, on the physicochemical fundamentals and on works' trials of autoclave salt leaching of polymetallic materials; I. Yu. Leshch, Gipronikel', on the unsuitability of autoclave leaching for lime-containing materials; V. A. Bernshteyn, VAMI, on industrial experience of a continuous autoclave leaching process for bauxites; V. G. Tronev, IONKh AN SSSR (IONKh AS USSR), on compounds of some rare elements in

Card 4/5

SOV/136-59-7-17/20

Conference on Autoclave Processes

various valency states under oxygen and hydrogen pressure in the presence of anhydrous ammonia; Z. L. Berlin, Gintsvetmet, on autoclave design and operation; P. G. Yakovlev, Gipronikel', and N. Ye. Vishnevskiy, VNII neftekhim, on model studies on autoclaves and the development of mixers; M. A. Polyanov, K. B. Giredmet, on the design of an experimental high-pressure pulp pump. G. L. Shvarts, NIIKhIMMASH, on the selection of steel for acid leaching of cobalt matte and matte-flotation concentrate; Yu. I. Archakov, VNIIneftekhim, on corrosion of types IKh18N9T, 12KhGN, 10KhSND and 10GND steels in soda and alkaline solutions in the presence of metal salts and oxygen at 5 - 15 kg/mm²; V. I. Deryabina and N. N. Kalgatin, VNIIneftekhim, separately, on mechanical properties of hydrogen-affected steels. The conference made recommendations aimed at the extension and improvement of autoclave practice.

Card 5/5

SVODTSEVA, G.V.; YUKHTANOV, D.M.

Simultaneous production of impregnated minerals during the recovery of heavy nonferrous metals in certain plants abroad.
(from foreign journals). TSvet. met. 33 no.7:98-102 J1 '60.

(MIRA 13:?)

(Nonferrous metals--Metallurgy)

S/080/60/033/009/003/021
A003/A001

AUTHORS: Yukhtanov, D.M., Plateneva, N.B.

TITLE: The Production of High-Purity Selenium

PERIODICAL: Zhurnal prikladnoy khimii, 1960, Vol. 33, No. 9, pp. 1951-1957

TEXT: The production of pure selenium from commercial selenium with 97.3% Se and from selenium with 99.99% Se was investigated. Commercial selenium contains a considerable amount of tellurium, the separation of which from selenium presents difficulties, because both elements are very similar. The separation is carried out by sublimation of the dioxides of the two metals. Commercial selenium is transformed to dioxide by burning in a flow of oxygen or a mixture of oxygen and nitrogen oxides. The burning temperatures used in the experiment were 500 and 560°C. The optimum conditions were found to be 560°C and an oxygen consumption of 1,000 ml/min. The stoichiometric oxygen consumption is 405 g per 1 kg of selenium. The actual consumption is 1 kg of oxygen, i.e., 250% of the theoretical. 99.99% selenium needs less oxygen and the burning is faster. At 560°C and a consumption of 500 ml/min the burning rate of 99.99% selenium is 100 g/hour, of commercial selenium 15 g/hour. The sublimation of selenium di-

Card 1/2

The Production of High-Purity Selenium

8/080/60/033/009/003/021
A003/A001

oxide was carried out in an air flow at 320-350°C. For a batch of 230-250 g selenium it lasted 2 hours. Selenium dioxide was washed with distilled water and elemental selenium was precipitated from selenious acid thus obtained. Hydrazine hydrate ($N_2H_4 \cdot H_2O$) is a good reducing agent. Its consumption is 72% based on the weight of selenium. Spectral analysis showed that high-purity selenium can be obtained from commercial selenium with 97.3% Se. There are 3 tables, 3 figures and 4 references: 3 Soviet, 1 German.

SUBMITTED: February 17, 1960

Card 2/2

MOLCHANOV, A.A.; YUKHTANOV, D.M.

Efficient utilization of copper and copper-zinc pyritic ores.
(MIRA 14:4)

TSvet. met. 33 no.6:21-25 Je '60.
(Copper ores) (Zinc ores)

S/137/61/000/006/088/092
A006/A101

AUTHORS: Yukhtanova, N.S., Gromova, V.S., Klark, G.B.

TITLE: Corrosion resistance of aluminum alloys with different galvanic coatings under atmospheric conditions

PERIODICAL: Referativnyy zhurnal. Metallurgiya, no. 6, 1961, 51, abstract 61401
("Tr. In-ta fiz. khimi, AN SSSR", 1960, no. 8, 173 - 180)

TEXT: During three years natural tests were made with Al-alloys of the following grades: A-1, A-2, L-1 (D1), D-16 (D16), Al-9. The tests were performed with alloys in delivery state and having galvanic coatings of Zn, Cd or the POC-40 (Pb-Sn) alloy (Pb-Sn). The tests were made under various climatic conditions. The thickness of the coatings was 40 μ . ✓

Ye. Layner

[Abstracter's note: Complete translation]

Card 1/1

YUKNA, R.D., Cand Tech Sci -- (diss) "Study of certain
problems of electromagnetic ~~resonating~~ shielding ^{of} ~~with~~
shields commensurable with ~~the~~ wave length."

Riga 1958, 1h pp with sketches (Latvian State Univ im

Petr Stuchka) 175 copies (KL, 39-58, 110)

- 50 -

6(4)

SOV/112-59-5-9989

Translation from: Referativnyy zhurnal. Elektrotehnika, 1959, Nr 5, p 223 (USSR)
AUTHOR: Yukna, R. D.
TITLE: Shielding a Vertical Grounded Radiator by a Cylindrical Shield With an End
or Ring Aperture

PERIODICAL: Uch. zap. Latv. un-t, 1958, Vol 21, Nr 7, pp 153-172

ABSTRACT: The problem of finding the shielding effect of a grounded vertical cylindrical shield with respect to a linear radiator placed in the shield axis is being solved. Two cases are considered: (1) incomplete shielding due to an open end of the shield; (2) incomplete shielding due to a round slit perpendicular to the shield axis and placed at an arbitrary height. Both cases are solved by means of introducing equivalent surface magnetic currents and using the specular-image method; both the ground and the shield are assumed to have perfect conductance. In the first case, calculations according to the first-approximation formulae show that the radiative capacity of the shield

Card 1/2

SOV/112-59-5-9989

~~-Shielding a Vertical Grounded Radiator by a Cylindrical Shield With an End or~~

currents amounts only to about 1% of the power radiated by the end aperture (the shield is not excited). In the second case, the shield is highly excited, so that the radiative power through the slit amounts only to a few per cent of the shield radiation power. An inference is drawn that the radiation of a large shield should be limited either by eliminating the coupling with the active element or by detuning the shield far away from the working frequency.

D.M.V.


Card 2/2

USSR/Chemistry - Polarographic analysis

Card 1/1 : Pub. 245 - 1/10
Author : Goldshteyn, Ya. P.; Sinyakova, S. I.; and
Title : Adaptation of oscillographic polarograph for
determination of Ti
Periodical : Zhur. anal. khim. 9/5, 255-264, Sept. 1954
Abstract : A method for polarographic or oscillographic
Ti in the presence of Fe, V, Cr, Ni and Mo
The mechanism of reduction of Ti compounds
factors of tartarate, citrate and oxalate
sulfuric acid, are explained. An effect of
sodium oxalate was found to be most marked.
The effect of Fe, V, Cr, Ni and Mo on the
Ti current, is elucidated. Eleven ref.
1-Belgian and 3-Czech (1932-1954).
Institution : Acad. of Sc. USSR, The V. I. Vernadsky
and Analytical Chemistry, Moscow
Received : March 13, 1954

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001963120008-1



APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001963120008-1"

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001963120008-1

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001963120008-1"

SOV / 20-120-1-37/63

AUTHOR: Yukhtanova, V. D.

TITLE:

The Determination of Polarographic Diffusion Coefficients
by Means of a Rotating Disk Electrode (Opredeleniye polaro-
graficheskikh koeffitsiyentov diffuzii s pomoshch'yu
vrashchayushchegosya diskovogo elektroda)

PERIODICAL: Doklady Akademii nauk SSSR, 1958, Vol. 120, Nr 1, pp. 137-139
(USSR)

ABSTRACT: At present only insufficient data are available on the so-called polarographic diffusion coefficient, i.e. on the so-called coefficients of diffusion in the case of an excess of the foreign electrolyte. Therefore the aim of the present paper is the determination of the diffusion coefficients by means of the reliable and relatively simple method of the rotating disk electrode. First the formula of V. G. Levich for the diffusion coefficient is given. By means of this formula the author calculates the diffusion coefficients of the ions Cd^{2+} , Tl^+ , Pb^{2+} and Zn^{2+} . The investigations were carried out by means of a copper-amalgamated disk electrode with a diameter of $4,997 \pm 0,002$ mm. These experi-

Card 1/3

304/20-120-1-57/67
Sov by Means of a

The Determination of Polarographic Diffusion Coefficients by Means of a Rotating Disk Electrode

experiments were carried out in a hermetically sealed cell in a hydrogen atmosphere at $t = 25 \pm 0.1$. The polarization according to the hydrogen ions Cd^{2+} , Tl^{+} , Pb^{2+} , plotted ranges of the limiting current, for the experiments, have also been calculated. In a diagram, within the same concentration of the experiment, the concentration of the ions Cd^{2+} , Tl^{+} , Pb^{2+} is shown as the author found a proportionality between the diffusion coefficient and the diffusion coefficient of the base. The diffusion coefficient of the base is determined by the diffusion coefficient of the ions Cd^{2+} , Tl^{+} , Pb^{2+} and the diffusion coefficient of the ions Zn^{2+} on a KCl base. The diffusion coefficient of the ions Cd^{2+} , Tl^{+} , Pb^{2+} and the diffusion coefficient of the ions Zn^{2+} on a KCl base are given. Thus the diffusion coefficients of other methods, various cases, with increasing concentrations of the base. There are 3 figures, 1 table, and 7 references, 3 of which are Soviet.

SOV2o-12o-1-37/63

The Determination of Polarographic Diffusion Coefficients by Means of a
Rotating Disk Electrode

PRESENTED: August 28, 1957, by A. N. Frumkin, Member, Academy of Sciences,
USSR

SUBMITTED: August 27, 1957

1. Cadmium ions--Diffusion
2. Lead ions--Diffusion
3. Thallium ions--Diffusion
4. Zinc ions--Diffusion
5. Polarization--Applications
6. Copper electrodes--
Applications

Card 3/3

SOA/20-124-2-38/71

Yukhtanova, V. D.

Yukhtanova, V. D.
The Migration Current on a Rotating Disk Electrode
(Migratsionnyy tok na vrashchayushchemsya diskovom elektrode)
Soviet science, 1959, Vol 124, No 2, pp 377-379

PERIODICALS

ABSTRACT:

If all ions in the solution have a simple charge and if the cation subjected to reduction has a common anion with the phonon, Euken's formula has the form $I = 2I_d [M - \sqrt{M(M-1)}]$. Here I_d denotes the boundary value of the diffusion current in the case of lacking migration; M - the ratio between the entire anion concentration and the concentration of the ion subjected to reduction. The present paper investigates Euken's formula at relatively low background concentrations carried out on a rotating amalgamated electrode. Measurements were 25 in hydrogen-atmosphere. The potential was measured relative to a 1n-calomel electrode. The experimental results were well reproducible within the entire measuring range. The polar-

card 1/3

APPROVED FOR RELEASE UNDER E.O. 14176 05 SEP 01 5312008-1

The Migration Current on a Rotating Disk Electrode SOV/20-124-2-38/71

rization curves of the Tl-ion are shown in a diagram. One of these curves was plotted over a background of sodium-perchlorate with a 50-fold surplus. In this case the boundary current amounts to $I = 5.4 \mu\text{A}$. The second polarization curve was plotted at the same concentration with a lacking indifferent electrolyte. This curve has a rather distinctly marked horizontal portion, from which the boundary current $I = 10.6 \mu\text{A}$ can be calculated. The curve agrees with Euken's formula, and by means of this formula it is possible to calculate the true diffusion currents in the case of highly diluted dilutions of the background. On the basis of the measuring series for the Tl-ion the authoress calculated the diffusion coefficients according to the formula by Levich. The values of these coefficients (as function of the square root of background concentration) suit a straight line, by the extrapolation of which to infinite dilutions, $D_{\text{extr}} = 19.90 \cdot 10^{-6}$ is obtained for the diffusion coefficient. This value agrees well with the value for infinitely diluted solutions ($D_{\lambda} = 19.94 \cdot 10^{-6} \text{ cm}^2/\text{sec}$) which was calculated by means of the Nernst formula from mobility. The authoress

Card 2/3

The Migration Current on a Rotating Disk Electrode SOV/20-124-2-38/71

I thank Academician A. N. Frumkin for his valuable advice in connection with this work. There are 2 figures, 1 table, and 7 references, 3 of which are Soviet.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet im. M. V. Lomonosova
(Moscow State University imeni M. V. Lomonosov)

PRESENTED: September 18, 1958, by A. N. Frumkin, Academician

SUBMITTED: September 12, 1958

Card 3/3

YU. TALOVA, V.D.

PHASE I BOOK EXPLOITATION

SOV/4443

Akademiya nauk SSSR. Komissiya po analiticheskoy khimii

Metody opredeleniya primesey v chistykh metallakh (Methods of Determining Admixtures In Pure Metals) Moscow, 1960. 411 p. (Series: Its: Trudy, 12) 3,500 copies printed.

Resp. Eds.: A.P. Vinogradov, Academician, and D.I. Ryabchikov, Doctor of Chemical Sciences; Ed. of Publishing House: M.P. Volymets; Tech. Ed.: T.V. Polyakova.

PURPOSE: This collection of articles is intended for chemists, metallurgists, and engineers.

COVERAGE: The articles describe methods for detecting and determining various admixtures and their traces in pure metals. Also discussed are many chemical, physicochemical, electrochemical, spectrochemical and luminescence methods of analyzing materials of high purity. The editors state that these methods have been developed within the last five or six years by various Soviet scientific institutes, and are now widely used in research and factory laboratories of the Soviet Union. No personalities are mentioned. References, mostly Soviet, accompany each article.

Card 1/2

Methods of Determining Admixtures(Cont.)

SCV/4443

TABLE OF CONTENTS:

Gokhshteyn, Ya. P., M.P. Volynets, and V.D. Yukhtanova. Determination of Admixtures of Copper, Lead, Zinc, Nickel, Iron, and Silver in High-Purity Metallic Germanium by the Oscillographic Polarization Method	5
Karabash, A.G., Sh. I. Peyzulayev, G.G. Morozova, and I.I. Smirenkina. Spectrochemical Method of Determining Admixtures in Metallic Germanium and Germanium Dioxide	25
Babko, A.K., and T.Ye. Get'man. Spectroscopic Detection of Small Quantities of Hydrogen in Metallic Germanium	36
Babko, A.K., and N.S. Kozachuk. Determination of Nitrogen Microadmixtures in Metallic Germanium	48
Babko, A.K. A.I. Volkova, and O.F. Drako. Determination of Small Quantities of Oxygen in Metallic Germanium	53
Melamed, Sh.G., A.K. Rusanov, and M.G. Zemskova. Determination of Tantalum and Niobium in the Pentoxide Mixture	65

Card 2/9

GGKHSHTEYN, Ya.P.; VOLNETS, M.P.; YUKHTANOVA, V.D.

Determining the presence of copper, lead, zinc, nickel, iron, cobalt, manganese, aluminum, tin, and antimony in high-purity metallic germanium by atomic absorption spectrometry
by E. N. Kuznetsov, Yu. P. Gokhshteyn,
(Germanium-Analysts) (P. I. S. T.)

YUKHTANOVA, V.D.

Testing Zucken's formula by means of a rotating disc electrode.
Zhur.fiz.khim. 35 no.12:2778-2779 D '61. (MIRA 14:12)

1. Akademiya nauk SSSR, Institut elektrokhimii.
(Electrochemistry)

YUKHTIN, V.I., kandidat meditsinskikh nauk

Tuberculous tumorous lesion of the large intestine. Sov. med.
(MIRA 7:11)
18 no.9:33-34 S '54.

1. Iz kliniki obshchey khirurgii pediatricheskogo fakul'teta
(direktor - professor G.P.Zaytsev) II Moskovskogo meditsinskogo
instituta im. I.V.Stalina

(TUBERCULOSIS, GASTROINTESTINAL
tuberculous tumoral swelling of large intestine)

TUKETIN, V.I.

Treatment of tetany by transplantation of the thyroid gland.
Vest.khir. 74 no.1:29-32 Ja-F '54. (MLRA 7:2)

1. Iz kliniki obshchey khirurgii (zavednyushchiy - professor G.P.Zaytsev) 2-go Moskovskogo gosudarstvennogo meditsinskogo instituta im. I.V.Stalina).
(Tetany) (Thyroid gland--Transplantation)

YUKHTIN, V.I., kandidat meditsinskikh nauk

Some peculiarities of clinical and anatomical changes in sepsis
in connection with compound therapy. Khirurgia no.6:53-58 Je '55.
(MLRA 8:10)

1. Is kliniki obshchey khirurgii pediatriceskogo fakul'teta
(zav.-prof. G.P.Zaytsev) II Moskovskogo meditsinskogo instituta
imeni I.V.Stalina)

(SEPTICEMIA AND BACTEREMIA, ther,
complex ther., clin. & anatomical aspects)

YUKHTIN, V. I., kandidat meditsinskikh nauk

Cases of melanoblastoma of the fingers. Nov.khir.arkh. no.3:74-75
My-Je '57.

(MIR 10:8)

1. Kafedra obshchey khirurgii (zav. - prof. G.P.Zaytsev) pediatriche-
skogo fakul'teta 2-go Moskovskogo meditsinskogo instituta. Adres
avtora: Moskva, Ul. Pavlovskaya, d.25, 4-ya gorodskaya klinicheeskaya
bol'ničsa.
(FINGERS--CANCER)

YUKHTIN, V.I., kand.med.nauk; SHALEVICH, M.A.

Tuberculosis of the stomach. Sov.med. 21 no.11:113-117 N '57.

(MIRA 11:3)

1. Iz kliniki obshchey khirurgii (dir.-prof. G.P.Zaytsev)
pediatricheskogo fakul'teta II Moskovskogo meditsinskogo instituta i
patologoanatomiceskogo otdeleniya 4-y gorodskoy klinicheskoy
bol'nitsy (zav.-prof. Ya.L.Rapoport).

TUBERCULOSIS, GASTROINTESTINAL, case reports)

YUKETIN, V. I., kandidat meditsinskikh nauk (Moskva, Pushkinskaya ul., d.16
kv. 8)

Gastrostomy with a musculo-aponeurotic press. Vest.Khir. 78 no. 6
136-139 Je '57.
(MLRA 10:8)

1. Iz kliniki obshchey khirurgii (zav. - prof. G.P.Zaytsev)
pediatriceskogo fakul'teta 2-go Moskovskogo meditsinskogo instituta
(STOMACH, surg.
gastrostomy with musculo-aponeurotic press)

YUKHTIN, V.I., dotsent (Moskva, Pushkinetskaya ul., d.16, kv.8)

Selection of the surgical method in cancer of the large intestine.
Vest.khir. 82 no.4:67-71 Ap '59. (MIRA 12:6)

1. Iz obshchey khirurgicheskoy kliniki (zav. - prof.G.P.Zaytsev)
2-go Moskovskogo meditsinskogo instituta im. N.I.Pirogova.
(INTESTINES--SURGERY)

YUKHTIN, V.I.

Single-stage ileocoloplasty in extensive resection of the large intestine in cancer. Vest. AMN SSSR 16 no.1:53-57 '61.
(MIRA 14:3)

1. Klinika obshchey khirurgii pediatricheskogo fakul'teta II
Moskovskogo meditsinskogo instituta imeni N.I.Pirogova.
(COLON-CANCER)