

ZIMINA, T.A.: KRYUKOVA, T.N.

Local cabbage variety of Sakhalin. Soob. Sakhal. kompl. nauch.-issl.
inst. AN SSSR no.4:102-106 '56. (MIRA 11:5)
(Sakhalin--Cabbage--Varieties)

ZIMINA, T. A.

5689. ZIMINA, T. A. Ogurtsy Na Sakhaline. Yuzhno-Sakhalinsk, Gaz (ob. Sakhalin)
1954. 23s. Ill. 20 sm. (Sakhalinskiy Filial "kad. Nauk SSR. Nauch.---Popul. Seriya).
1600 kuz 60 k.--(55-1473) 635.63(57.143.4)

SO: Knizhnaya, Letopis, Vol. 1, 1955

ZIMBA T.P

7

Monoclonal antibody system of analysis and synthesis of
carbohydrates and sulfonates. G. L. Alagashvili, P. S. Zil'berman,
I. S. Zil'berman, I. S. Zil'berman, I. S. Zil'berman, I. S. Zil'berman

ZIMINA, T.D.; BERGMAN, A.G.; NAGORNYI, G.I.

Reciprocal system consisting of chlorides and sulfates of sodium, calcium, and barium. Zhur. neorg. khim. 10 no.9:2145-2151 S '65.

(MIRA 18:10)

1. Irkutskiy gosudarstvennyy universitet i Rostovskiy-na-Donu institut sel'skokhozyaystvennogo mashinostroyeniya.

BELYAYEVA, V.A.; ZAKHVALINSKIY, M.N.; ZIMINA, T.D.; DEMINA, T.N.;
KALASHNIKOV, P.V.; NAGORNAYA, Ye.F.; NAGORNIY, G.I.; TITOVA, T.P.

Adsorption properties of Gynyl' argillites. Trudy DVEAN SSSR.
Ser.khim. no.7:18-25 '65.

(MIRA 18:12)

ZIMINA, Mariya Alekseyevna; EDEL'SHTAYN, V.I., prof., otvetstvennyy red.;
KRYLOV, S.V., red. izdatel'stva; POLESITSKAYA, S.M., tekhn. red.

[Vegetable gardening in Sakhalin] Ovoshevodstvo na Sakhaline.
Moskva, Izd-vo Akad. nauk SSSR, 1957. 241 p. (MIHA 10:11)
(Sakhalin--Vegetable gardening)

ZIMINA, T.D.; BERGMAN, A.G.; NAGORODNYI, G.I.

Diagonal sections of the quaternary reciprocal system consisting of sodium, calcium, and barium chlorides and sulfates. Ukr. khim. zhur. 31 no.10:1035-1040 '65. (MIRA 19:1)

1. Irkutskiy gosudarstvennyy universitat i Rostovskiy-na-Donu institut sel'skokhozyaystvennogo mashinostroyeniya. Submitted May 23, 1964.

ZIMINA, T.S.

Some characteristics of cholesterol metabolism in children with diseases of the blood system. Vop. gemat. v pediat. no.3:90-101 '64.

Dynamics of the content of general cholesterol and its fractions in children with acute and chronic leukemia. Ibid.:317-327

(MIRA 18:7)

POLYAKOV, S.M.; ZIMINA, T.S.

Obtaining replicas by the simultaneous evaporation of carbon and
preshadowed substance. Zav.lab. 29 no.8:973-974 '63.

(Electron microscopy)

(MIRA 16:9)

ZAKHVATKIN, M.O.; SAPIR, A.D.; SPIVAKOVSKIY, V.B.; ZIMINA, V.A.; MARGOLIS,
L.D.

Exchange of experience. Zav.lab. 28 no.3:290 '62. (MIRA 15:4)

1. Chelyabinskiy metallurgicheskiy zavod (for Zakhvatkin, Sapir).
2. Kiyevskiy gosudarstvennyy universitet (for Spivakovskiy, Zimina).
3. Dneprovskiy alyuminiyevyy zavod imeni S.M.Mirova (for Margolis).
(Metallurgical analysis)

SPIVAKOVSKIY, V.B.; ZIMINA, V.A.; GAVRILYUK, L.S.

Determination of uranium traces in rocks and natural waters. Zav.
lab. 27 no. 4:390-391 #61. (MIRA 14:4)

1. Kiyevskiy gosudarstvennyy universitet imeni T.G. Shevchenko.
(Uranium--Analysis) (Rocks--Analysis)
(Mineral waters)

ZIMINA, V.I.
AUTHOR: Zimina, V.I. ■

"Theory of Propagation of Electromagnetic Waves Along Tubes Filled with Ionized Gas,"
A-U Sci Conf dedicated to Radio Day," Moscow, 20-25 May 1957.

PERIODICAL; Radiotekhnika i Elektronika, Vol. 2, No. 9, pp. 1221-1224, 1957.
(USSR)

8053

9.4/20

S/109/60/005/06/008/021
E140/E163

AUTHOR: Zimina, V.I.

TITLE: Experimental Study of Electromagnetic Wave Propagation
Along a Cylinder of Ionised Gas

PERIODICAL: Radiotekhnika i elektronika, 1960, Vol 5, Nr 6,
pp 938-942 (USSE)

ABSTRACT: In Refs: 1 and 2 it was shown that electromagnetic waves may propagate along a cylinder formed by ionised gas in the presence of negative permittivity. The transfer of electromagnetic energy along the ionised gas cylinder has the following properties: the possibility of varying the propagation constant and phase velocity in wide limits by regulation of the number of electrons in the ionised gas; a high concentration of electromagnetic energy about the surface of the cylinder. The present article describes the results of experimental study and their comparison with theory. The following questions

S/109/60/005/06/008/021
E140/E163

Experimental Study of Electromagnetic Wave Propagation Along a
Cylinder of Ionised Gas

concentration; the attenuation of electromagnetic energy propagating along the gas-filled tube. It was found that mercury vapour is more advantageous for the experiments than the inert gases. The experiment confirmed the essential dependence of phase velocity on the dielectric permittivity of the ionised gas and the presence of a high delay of the electromagnetic waves at low values of the permittivity. The rapid decrease of field in the radial direction is also indicated. There are 4 figures and 2 references, of which 1 is Soviet and 1 is German.

Card
2/2

SUBMITTED: August 4, 1959

ZIMINA, V. I., Candidate Tech Sci (diss) -- "The propagation of electromagnetic waves along a dielectric cylinder filled with ionized gas". Moscow, 1959, published by the Acad Sci USSR. 16 pp (Acad Sci USSR, Inst of Radio Engineering and Electronics), 150 copies (KL, No 22, 1959, 115)

S/109/62/007/006/008/024
D266/D308

9.3700
AUTHOR:

Zimina, V. I.

TITLE:

Propagation of electromagnetic waves along a conducting cylinder surrounded by a layer of ionized air having a negative dielectric constant

PERIODICAL:

Radiotekhnika i elektronika, v. 7, no. 6, 1962, 988-994

TEXT: The analysis follows that of L. A. Vaynshteyn (Elektromagnitnye volny (Electromagnetic Waves), Izd. Sovetskoye radio, 1957) for an ordinary dielectric. The characteristic equation in terms of Bessel functions is obtained and solved graphically. It is shown that propagation is possible when the relevant dielectric constant of the plasma is negative. At $\epsilon_p = -1$ the axial propagation constant h is large, but as ϵ_p decreases h tends to its free space value. Integrating the Poynting vector.

✓
B

Propagation of electromagnetic ...

S/109/62/007/006/008/024
D266/D308

An analytical expression is obtained outside the plasma and a numerical solution inside the plasma. If $|\epsilon_p|$ is large most of the power is carried in the air outside the plasma. For a given ϵ_p the distribution of power in the plasma is nonuniform, concentrated mainly near the plasma-air boundary. Losses caused by collisions are generally very small but as ϵ_p tends to -1 they increase substantially. There are 6 figures.

SUBMITTED: July 3, 1961

GORODISSKAYA, G.Ya., prof., doktor med. nauk, otv. red.; BLOKHINA,
I.N., red.; GUSEVA, V.A., red.; DIKOVSKIY, F.F., red.;
ZIMINA, V.S., red.; LAZOVSKAYA, A.L., red.; PEROVA, R.S.,
red.

[Biochemistry of microbes] Biokhimiia mikrobov; sbornik
trudov. Gor'kii, 1964. 427 p. (MIRA 17:12)

1. Gorki. Gor'kovskiy nauchno-issledovatel'skiy institut
epidemiologii i mikrobiologii.

1. A. Z. SCRKIN, Prof., V. S. ZEMINA
2. USSR (600)
4. Bones - Tuberculosis
7. Lymph therapy of peripheral lymph node tuberculosis and osteoarticular tuberculosis. Probl. tub. no. 6. 1952.

1. SORKIN, A.Z., ZIMINA, V.S.
2. USSR (600)
4. Lymph
7. Lymph therapy of peripheral lymph node tuberculosis and osteoarticular tuberculosis.
Probl tub No. 6, 1952

SIMINA, V.S., sanitarnyy vrach

Work of the Khotin health and epidemiological division consolidated
with a district hospital. Gig. 1 san. 23 no. 6:41-43 Jo 158

(WZRA 11:7)

(SANITATION,

in Russia, cooperation with rural hosp. (Rus))

(HOSPITALS,

rural hosp. in Russia, cooperation with sanitation
serv. (Rus))

SORKIN, A.Z.; ZIMINA, V.S.

Therapy with lymph of peripheral lymph node tuberculosis and osteo-articular tuberculosis. Probl. tuberk. Moskva no. 6:25-29 Nov-Dec 1952. (CIME 2:15)

1. Professor for Sorkin. 2. Of Moscow Municipal Scientific-Research Tuberculosis Institute (Director -- Prof. V. L. Eynis) and of the Fourth Tuberculosis Dispensary of Moscow Municipal Public Health Department (Head Physician -- S. M. Zamukhovskiy).

1. SORKIN, A.Z., ZIMINA, V.S.
2. USSR (600)
4. Joints - Tuberculosis
7. Lymph therapy of peripheral lymph node tuberculosis and osteoarticular tuberculosis
Probl. tub. N_o. 6 1952.

BOGDANOV, G.A.; ZIMINA, V.V.

Automatic weight proportioning equipment. Ogneupory 30 no.3:15-16
'65. (MIRA 18:5)

1. Borovichskiy kombinat ogneuporov.

ZIMINA, V.Ye.

OLISUF'YEV, N.G.; TSVETKOVA, Ye.M.; BORODIN, V.P.; KOROLEVA, A.P.; SIL'CHENKO, V.S.; KHOROSHEV, I.G.; MYASHNIKOV, Yu.A.; PERFIL'YEVA, N.A.; KRATOKHIVIL' N.I.; YAYSTIKH, M.A.; RAVDONIKAS, O.V.; BARANOVA, N.K.; ZIMINA, V.Ye.; TORMASOVA, L.N.; USTIN-PETROVA, T.F.; ABEF'YEV, S.S.; KONKINA, N.S.; KUL'BA, A.P.; MAL'TSEVA, N.K.; SHELANOVA, G.M.; SORINA, A.M.; BRANITSKAYA, V.S.; PRUDNIKOVA, M.N.

Tularin from a vaccinal strain for epicutaneous use. Zhur. mikro-biol.epid. i immun. 27 no.9:22-28 S '56. (MLRA 9:10)

1. Iz Instituta epidemiologii i mikrobiologii im. N.F.Gamalei AMN SSSR i protivotuliaremiynykh stantsiy Stalingradskoy, Voronezhskoy, Tul'skoy, Plavskoy, Omskoy, Krasnodarskoy, Moskovskoy i Smolenskoy. (TULAREMIA, diagnosis, tularin epicutaneous test (Rus))

ZIMINA, Ye.A.; TARASENKO, Ye.N.

Study by the photoelastic method of the stressed state of
rocks around sublevel entries in thick steeply pitching seams.
Zap. LGI 48 no.1:3-11 '63. (MIRA 17:8)

ZIMINA, Ye. A.

"Study of Advanced Methods of Pushing Drifts Through Slightly Dipping Seams, as, for Example, in the Mines of the Donbais."
Cand Tech Sci, Chair of Construction of Mining Enterprises, Leningrad Order of Lenin and Order of Labor Red Banner Mining Inst, Min Higher Education USSR, Leningrad, 1955. (ML, No 11, Mar 55)

SO: Sum. No. 670, 29 Sep 55-Survey of Scientific and Technical Dissertations Defended at USSR Higher Educational Institutions (15)

ZIMINA, Ye.A., kand.tekhn.nauk

Temporary supports in making galleries in mined areas. Izv.vys.
ucheb.zav.; gor.shur. no.3:64-74 '59. (MIRA 13:4)

1. Leningradskiy ordena Lenina i ordena Trudovogo Krasnogo
Znameni gornyy institut imeni G.V.Plekhanova. Rekomendovana
kafedroy stroitel'stva gornyykh predpriyatiy.
(Mining engineering)

BOKIY, Boris Vyacheslavovich, prof.; ZININA Yekaterina Aleksandrovna,
dots.; SMIRNYAKOV, Vitaliy Vasil'yevich, dots.; TIMOFREYEV,
Oleg Vladimirovich, dots.; FEDOROV, S.A., prof., retsenzent;
SHELEEV, A.I., red.izd-va; LOMILINA, L.N., tekhn. red.

[Mining engineering and mine supports] Provedenie i kreplenie
gornykh vyrabotok. [By] B.V.Bokii i dr. Moskva, Gosgortekh-
izdat, 1963. 557 p. (MIRA 17:2)

ZIMINA, Ye.A., dotsent

Using the photoelastic method to study the rock of a mine roof
in the stressed state in supporting it with rod bolting. Izv.vys.
ucheb.zav.; gor.zhur. no.3:19-24 '61. (MIRA 15:4)

1. Leningradskiy ordena Lenina i ordena Trudovogo Krasnogo Znani
gornyy institut imeni G.V.Plekhanova; rekomendovana kafedroy
stroitel'stva gornykh predpriyatiy Leningradskogo gornogo
instituta.

(Moscow Basin--Mine roof bolting)
(Rock pressure) (Photoelasticity)

BOKIY, Boris Vyacheslavovich, prof.. Prinizala uchastiye ZIMINA, Ye.A.,
kand.tekhn.nauk. SHUSIKOVSKAYA, Ye.L., red.isd-va; VINOGRADOVA,
G.V., red.isd-va; BERESLAVSKAYA, L.Sh., tekhn.red.

[Mining engineering] Gornoe delo. Izd.3., ispr. i dop. Moskva,
Gos.nauchno-tekhn.isd-vo lit-ry po gornomu delu, 1959. 863 p.
(MIRA 13:3)

(Mining engineering)

ZIMINA, YE. A.

15.8580

L3537

S/196/62/000/023/004/006
E194/E155

AUTHORS: Vodop'yanov, M.A., Vorozhtsov, B.I.,
Potakhova, G.K., Lavrov, M.D., Nesmelova, Ye.S.,
Nesterov, V.M., Vorozhtsova, I.G., Ol'shanskaya, N.I.,
Zimina, Ye.A., Mikhaylova, T.G., Sitonhevskaya, G.V.,
and Filatov, I.S.

TITLE: The influence of betatron radiation on the
dielectric properties of certain electrical
insulating materials

PERIODICAL: Referativnyy zhurnal, Elektrotehnika i energetika,
no.23, 1962, 12-13, abstract 23 B 67. (In collection:
Elektron. uskoriteli (Electronic Accelerators),
Tomsk, Tomskiy un-t, 1961, 308-318)

TEXT: The temperature and frequency characteristics of
electrical insulating materials were investigated before and after
 γ -irradiation at dosages ranging from 10^4 to 2×10^5 rads with a
dosage rate ranging from 300 to 1500 rads/minute at temperatures
of -60, -20 and +60 °C and under tropical conditions (40 °C and
relative humidity of 98%); the source of radiation was a
Card 1/3

The influence of betatron radiation... S/196/62/000/023/004/006
E194/E155

15 MeV betatron. The characteristics of polyethylene were not altered by a radiation dose of 10^5 rads (the measurements were made at about 10^9 c/s). The low-frequency $\tan \delta$ of plastic АГ-4 (AG-4) increased (particularly after irradiation under tropical conditions and at -60°C) but the value in the frequency range $10^5 - 10^6$ c/s did not alter. Evidently irradiation increases the resistive component of loss by conductivity and does not alter the relaxation components. Similar results were obtained for plastics K-114-35, K-211-3 and ФКНМ-25 (FKPM-25). In the case of textolite with a silicoorganic binder СКМ-1 (SKM-1), a dosage rate of 500 rads/min first increases the low-frequency $\tan \delta$ only up to about 10^5 rads, and then diminishes it. Above 1200 rads/min the $\tan \delta$ steadily decreases. It is possible that with heavy dosages and high dosage rates a process of binding together reduces the $\tan \delta$. In the silicoorganic resins 14P-2 (14R-2), 14R-6 and 14R-15, dosage rates of 500 rads/min and a dosage of 10^5 rads cause a small increase in conductivity and $\tan \delta$ at low frequency, but this change disappears as temperature curves are being taken, so that the shape of the reverse temperature curve coincides with that

Card 2/3

The influence of betatron radiation.. S/196/62/000/023/004/006
E194/E155

for the non-irradiated material. Irradiation of varnishes K-47, 976-1, and MIM-16 (MGM-16) under various conditions caused no change in their electrical insulating properties. Irradiation of steatite ceramic (1% BaO, 91.6% Onot talc, 5.2% kaolin, 3.2% boracite) (with a dosage of 2×10^5 rads) did not alter the shape of the temperature curve of $\tan \delta$ (measured at 10^7 c/s) either in weak fields (945 V/cm) or in strong (1890 V/cm). With a dosage of 2.12×10^7 rads, $\tan \delta$ measured at 945 V/cm was not altered at low temperatures but increased appreciably at temperatures above 400 °C.

15 illustrations, 31 references.

[A tractor's note: Complete translation.]

Card 3/3

S/081/62/000/003/081/090
B160/B101

AUTHORS: Zimina, Ye. A., Senuk, D. P.

TITLE: Relationship of optical and mechanical properties of ED-6 epoxy resinbase materials to composition and production techniques

PERIODICAL: . Referativnyy zhurnal. Khimiya, no. 3, 1962, 563, abstract 3P40 (Zap. Leningr. gorn. in-ta, v. 44, no. 1, 1961, 59-63)

TEXT: A study is made of the optical and mechanical properties of ЭД-6 (ED-6) epoxy resin-base materials used as models for studying the distribution of stresses occurring during use in relation to their composition and production techniques. The optical constant $\sigma_0^{-1.0}$, Young's modulus E, Poisson's ratio μ and the quality factor $K = E/\sigma_0^{-1.0} \cdot 10^{-3}$ were determined at $\sim 20^\circ$ and "freezing" point. It was established that there is little change in elasticity at different percentage ratios of ED-6

Relationship of optical and...

S/081/62/000/003/081/090
B160/B101

modulus of elasticity of ED-6 resin-base optically active materials can be varied by a factor of 2-12 by changing the quantity ratio of all the material's components. The introduction of a plasticizer - dibutyl phthalate - at the rate of 1, 2, 3, 5, 7 and 10% of the weight of the resin at the optimum resin/hardener ratio of 100:30 changes E by a factor of 2-12 at "freezing" point (from 170-200 to 20 kg/cm²); there is little change in $\epsilon_0^{1.0}$. An increase in the plasticizer's percentage content in the resin leads to a certain reduction in the material's quality factor. [Abstracter's note: Complete translation.] ✓

ZIMINA, Ye.A.; SENUK, D.P.

Effect of the production techniques and composition of materials
from ED-6 epoxy resins on their opticomachanical properties. Zap.
LGI 44 no.1:59-63 '61. (MIRA 14:10)
(Epoxi resins) (Engineering models) (Rock pressure)

ZIMINA, Ye.A., kand.tekhn.nauk

Use of bolting for the control of rock swelling in Donets Basin
mines. Nauch.dokl.vys.shkoly; gor.delo. no.4:61-65 '58.

(MIRA. 12:1)

1. Predstavleno kafedroy stroitel'stva gornykh predpriyatiy Leningrad-
skogo gornogo instituta imeni G.V. Plekhanova.
(Donets Basin--Mine roof bolting)

ZIMINA, Ye. A.

AFANAS'YEVA, A.L., kand.biol.nauk; BAYMRTUYEV, A.A., kand.sel'skokhozyaystvennykh nauk; BAL'CHUGOV, A.V., kand.sel'skokhozyaystvennykh nauk; BELOZEROVA, M.A., agronom; BLOZOROV, A.T., kand.sel'skokhozyaystvennykh nauk; MAKSIMENKO, V.P., agronom; BERNIKOV, V.V., doktor sel'skokhozyaystvennykh nauk; BOGOMYAGKOV, S.T., kand.sel'skokhozyaystvennykh nauk; VOLYNETS, O.S., agronom; BODROV, M.S., kand.sel'skokhozyaystvennykh nauk; BOGOSIAVSKIY, V.P., kand.tekhn.nauk; KHRUPPA, I.F., kand.tekhn.nauk; VERBNER, A.R., doktor biol.nauk; VOZBUTSKAYA, A.Ye., kand.sel'skokhozyaystvennykh nauk; VOINOV, P.A., kand.sel'skokhozyaystvennykh nauk; VYSOKINS, G.P., kand.biol.nauk; GALDIN, M.V., inzhener-mekhanik; GERASIMOV, S.A., kand.tekhn.nauk; GORSHENIN, K.P., doktor sel'skokhozyaystvennykh nauk; YELENEV, A.V., inzhener-mekhanik; GERASKVICH, S.V., mekhanik [deceased]; ZHARIKOVA, L.D., kand.sel'skokhozyaystvennykh nauk; ZHEGALOV, I.S., kand.tekhn.nauk; ZIMINA, Ye. A., agronom; BARANOV, V.V., kand.tekhn.nauk; PAVLOV, V.D.; IVANOV, V.K., kand.sel'skokhozyaystvennykh nauk; KAPLAN, S.M., kand.sel'skokhozyaystvennykh nauk; KATIN-YARTSEV, L.V., kand.sel'skokhozyaystvennykh nauk; KOCHERGIN, V.I., doktor sel'skokhozyaystvennykh nauk; KOZHEVNIKOV, A.R., kand.A.Ye., kand.sel'skokhozyaystvennykh nauk; KOZHEVNIKOV, A.R., kand.sel'skokhozyaystvennykh nauk; KUZNETSOV, I.N., kand.sel'skokhozyaystvennykh nauk; LAMBIN, A.Z., doktor biol.nauk; LEONT'YEV, S.I., kand.sel'skokhozyaystvennykh nauk; MAYBORODA, M.M., kand.sel'skokhozyaystvennykh nauk; MAKAROVA, G.I., kand.sel'skokhozyaystvennykh nauk;

AFANAS'YEVA, A.L.... (continued) Card 2.

MIKIFOROV, P.Ye., kand.sel'skokhozyaystvennykh nauk; NEMASEEV, N.I.,
lesovod; PERVUSHINA, A.N., agronom; PLOTNIKOV, N.A., kand.biol.nauk;
L.G.; kand.sel'skokhozyaystvennykh nauk; PAVLOV, V.D., kand.tekhn.
nauk; PRUTSKOVA, M.G., kand.sel'skokhozyaystvennykh nauk; GURCHENKO,
V.S., agronom; POPOVA, G.I., kand. sel'skokhozyaystvennykh nauk;
PORTYANKO, A.F., agronom; RUCHKIN, V.N., prof.; RUSHKOVSKIY, T.V.,
agronom; SAVITSKIY, M.S., kand.sel'skokhozyaystvennykh nauk; BOLDIN,
D.T., agronom; NESTEROVA, A.V., agronom; SERAFIMOVICH, E.B., kand.
tekhn.nauk; SMIRNOV, I.N., kand.sel'skokhozyaystvennykh nauk;
SEREBRYANSKAYA, P.I., kand.tekhn.nauk; TOKHTUYAN, A.V., kand. sel'sko-
khozyaystvennykh nauk; PAL'KO, O.S., iznh.; FMDYUSHIN, A.V., doktor
biol.nauk; SHEVLYAGIN, A.I., kand.sel'skokhozyaystvennykh nauk;
YUFEROV, V.A., kand.sel'skokhozyaystvennykh nauk; YAKHTENFELD, P.A.,
kand.sel'skokhozyaystvennykh nauk; SEMENOVSKIY, A.A., red.; GOR'KOVA,
Z.D., tekhn.red.

[Handbook for Siberian agriculturists] Spravochnaya kniga agronoma
Sibiri. Moskva, Gos. izd-vo sel'khoz. lit-ry. Vol.1. 1957. 964 p.
(Siberia--Agriculture) (MIRA 11:2)

BERKUTOV, A.N., general-mayer meditsinsky aluzhby; KOROCHKINA, A.G.;
BOGACHOVA, D.I.; ZIMINA, Ye.P.

Direct blood transfusion in the treatment of acute radiation
sickness; an experimental study. Voen.-med. zhur. no.2:26-28
'65. (MIRA 18:11)

ZIMINA, Z.P. (Leningrad)

Changes in basal metabolism during burns. Pat. fiziol. i eksp.
terap. 4 no. 5:59 S-0 '60. (MIRA 13:12)

1. Iz patofiziologicheskoy ozhogovoy laboratorii (nachal'nik-
dotsent Ye.V. Gubler) kafedry gospital'noy khirurgii No.1
(nachal'nik - prof. I.S. Kolesnikov) Vbyenno-meditsinskoy
ordena Lenina akademii imeni S.M. Kirova.

(BURNS AND SCALDS) (BASAL METABOLISM)

ZIMINA, Z. V., Cand Med Sci -- (diss) "Disorders in reading in the case of focal diseases in the brain." Lutsk, 1960. 16 pp; (Ministry of Public Health Ukrainian SSR, Chernovtsy State Medical Inst); 200 copies; price not given; (KL, 17-60, 169)

~~ZIMINA, Z.V.~~

Certain characteristics of reading disorders in focal lesions of the parietal and occipital lobes. [with summary in French]. Zhur. nevr. i psikh. 58 no.8:929-933 '58 (MIRA 11:9)

1. Otdel nevrologii (nav. - prof. L.B. Litvak) i laboratoriya tsitoarkhite ktoniki Ukrainskogo nauchno-issledovatel'skogo psikhonevrologicheskogo instituta.

(ALEXIA, etiol. & pathogen.

occipital & parietal lesions (Rus))

(OCCIPITAL LOBE, dis.

causing alexia (Rus))

(PARIETAL LOBE, dis.

same (Rus))

ZIMINI I.A.

ZIMINI, I.A.; LINDIN, G.P.

Two cases of trichinosis in Sukhumi. Med.paraz. i paraz.bol.
supplement to no.1:66 #57. (MIRA 11:1)

1. Iz Abkhazskoy respublikanskoy protivomalyariynoy stantsii i
infektsionnoy bol'nitsy.
(SUKHUMI--TRICHINA AND TRICHINOSIS)

ZIMINOV, N.V.; SMIRNOV, Yu.F.; FAZLULLIN, M.I.

Results of the study of the dustiness of mine air in prospecting
drilling. Izv. vya. ucheb. zav.; geol. i razv. 6 no.5:140-145
My '65. (MIRA 18:10)

1. Sredneaziatskiy institut geologii i mineral'nogo syr'ya
(SAIGIMS).

ZIMINOVA, N.A.

Elements of the hydrological regimen and water balance of
Ivan'kovo Reservoir during 1951-1956. Trudy Inst.biol.
vodokhran. no.2:212-228 '59. (MIRA 13:5)
(Volga Reservoir--Hydrology)

ZIMINOVA, N.A.

Quantitative characteristics of suspended matter in Rybinsk
Reservoir. Trudy Inst. biol. vnutr. vod no.6:230-249 '63.

(MIRA 18:1)

157 and 158 (CCHN)

10

Catalytic properties of bentonite. I. Dehydration of isocyanic acid. M. N. Turrova-Pitlak and N. I. Zhurav (Moscow State Univ.). J. Applied Chem. (U.S.S.R.) 19, 2447-7 (1946) (in Russian); Bentonite (24), (in J. AL₂O₃ 13.50, Fe₂O₃ 1.30, CaO 0.80, MgO 0.51, K₂O 1. Na₂O 0.05, ignition loss 4.77, moisture 11.28), finely ground, was activated by 6 hrs. heating on a water bath with 30% H₂O₂ (30 g. H₂SO₄ sp. gr. 1.84, per 100 g. clay), washing and drying at 100°, then at 160-170°, and reheating 1 hr. at 400-450°; lower (300°) or higher (500°) temp. of heating results in less active catalyst. Dehydration expts. were made with 10 g. (30 ml.) catalyst in flowing isocyanic acid. (1) At const. rate of flow, 6 ml./hr. (vol. ratio 0.2), the yield, Y, in anylene, C₆H₅, increases with rising temp., from traces at 150°, to 30, 74, 91% at 250, 350, 450°, resp.; a max. is reached at the latter temp., followed by a decline (73% at 500°); expts. with fresh batches of the catalyst gave the same results. With nonactivated bentonite, Y = 13.2% at its max. at 350°. (2) At const. 450°, Y falls sharply with increasing rate of flow; e.g., for 0.05, 0.2, 0.5, 3.3 ml./min., Y = 90.5, 79.1, 71.1, 43.5%; optimum (Y = 91.0) is at 0.1 ml./min. (3) Under the optimum conditions (450°, 0.1 ml./min.), the activity of the catalyst remained unchanged 1 hr., then fell off gradually (Y = 55 and 40 after 90 and 120 min., resp.). Full regeneration was brought about by passing N at 500-550° to remove the C₆H₅, then air at the same temp. 1 hr. N. Thon

ASS. I.L.A. METALLOGICAL LITERATURE CLASSIFICATION

157 and 158 (CCHN)

CA

2

Promotion of nickel catalyst by hydrogen. L. Kh. Freidlin and N. I. Ziminova. *Izv. Akad. Nauk S.S.S.R., Otdel. Khim. Nauk* 1950, 650-61. — Studies on inactivity of pure deposed films of Ni as catalysts were continued (Kozlovskii, C.A. 26, 32*). Leaching of Al_2Ni_3 3 hrs. at 105° with NaOH, followed by washing with H_2O , HCl , and toluene gave the basic catalyst contg. 2.33 g. Ni per ml. After satu. with H_2 for 10 min., residual H_2 was displaced with

N_2 and the unsatd. compd. was introduced; then hydrogenation was carried at the expense only of H_2 retained by the catalyst. After washing with toluene the operation was repeated. $PhNO$, $CH_2=CH(OH)$, etc., were most satisfactory for the removal of the retained (adsorbed) H_2 from the catalyst. Removal of as much as 210 ml. adsorbed H_2 from 2.33 g. Ni still leaves certain activity (hydrogenating) in the catalyst, and the latter is pyrophoric; but removal of 321-4 ml. H_2 (26-hr. contact of the unsatd. compd.) completely inactivates the catalyst, which is then nonpyrophoric. The removal of H_2 is speeded by higher temp., and at 60° the max. amt. of H_2 removable from 2.33 g. Ni is about 224 ml., and only with removal of the last 3-4 ml. does the activity vanish. Thus, skeletal Ni is a catalyst that is promoted by adsorbed H_2 ; possibly other forms of Ni catalyst are similarly promoted. Some inhibitors (un-described) on hydrogenation over Ni lead to complete poisoning of the catalyst, which illustrates a case of the poison being either the reacting substance itself or an intermediate that reacts with the promoter. G. M. Kosolapoff

CA

2

Differentiated outgassing of nickel. Two forms of binding of hydrogen in a catalyst. L. Kh. Fridlin and N. I. Zimhova (Acad. Sci. U.S.S.R., Moscow). Doklady Akad. Nauk S.S.S.R. 74, 937-8 (1961). — Differentiation between the H₂ adsorbed on and the H₂ dissolved in H₂-saturated Ni was realized, and the nonequivalence of the 2 forms of H₂ was demonstrated by expts. of hydrogenation of org. compds. with the H₂ contained in Ni. The org. compds. were hydrogenated, in a Ni atom., with Ni satd. with H₂ for 10 min. prior to the expt. The compds. investigated fell into 3 groups: Group I, represented by Michler's ketone, withdraws from the Ni only a definite fraction of its H₂ content. Thus, 2.33 g. Ni gave up a total of 49-60 ml. H₂. On subsequent expts. with H₂, that amt. of Ni takes up exactly the

same amt. of H₂ as given up in the hydrogenation, and the process can be alternated repeatedly, with the amts. of H₂ given up and taken up again remaining the same. That 49-60 ml. H₂ is evidently the adsorbed H₂. With substances of group II, represented by styrene, cinnamic acid, or its ester, the 2.33 g. of Ni gives up 100-200 ml. H₂, and, on subsequent expts., takes up only 10-21 ml. If now the Ni is treated with a compd. of group I, it will give up only those 10-21 ml. Consequently, compds. of group II withdraw from the Ni, in addn. to the adsorbed H₂, also part of the dissolved H₂, with a concomitant partial destruction of the adsorptive capacity of the Ni, i.e. partial destruction of its active centers. Substances of group III, e.g. PhNO₂ and vinyl butyl ether, withdraw the total amt. of both the H₂ adsorbed and that dissolved in the Ni; the amts. withdrawn from 2.33 g. Ni by different compds. of that group were very close, 219-221, 223-224, and 223-234 ml. This is also the total amt. of H₂ that can be withdrawn from 2.33 g. of Ni by first treating it with a compd. of group I and then, after washing with toluene, by a substance of group III. After removal of all the H₂, the Ni becomes inactive for the hydrogenation of compds. of group I. The H₂ adsorbed on the surface evidently focuses no part of the active centers. In the Ni sample investigated, the ratio H₂(adsorbed):H₂(dissolved):Ni = 2:7:19, i.e. the sample contains 1 atom of dissolved H₂ per 2.7 atoms Ni. This coincides with the ratio H₂:Ni = 1:3 arrived at by Davison and Garner (C.A. 32, 350) by electron diffraction. The no. of H atoms dissolved in Ni is about 3.5 times the no. of H atoms adsorbed; i.e., the no. of potentially possible active centers is 3.5 times the no. of actual centers. The no. of the latter is roughly 1/3 the no. of Ni atoms.

N. Thon

CA

The atomic structure of the nickel-hydrogen catalyst. L. Kh. Franklin and N. I. Zhigonyi (Acad. Sci. U.S.S.R., Moscow). *Izv. Akad. Nauk S.S.S.R., Otdel. Khim. Nauk* 1961, 145-9. In a skeleton Ni catalyst, prepd. by leaching a 50% Ni-Al alloy at 100°, the atom ratio $H_{ads}:Ni$ was found = 1:3.5:9 (subscripts ads and dis referring to adsorbed and dissolved H, resp., detd. by the previously (C.A. 48, 16304) described method of selective hydrogenation of org. compds.). H_{ads} gives the measure of the active centers present at the surface. That this is directly related to the amt. of the structural H_{ads} is shown experimentally by progressive removal of H_{ads} with the aid of increasing amts. of 1-methyl-1-cyclopentene (I), in the absence of free H_2 . For complete removal of the H_{ads} corresponding to the given amt. of I, 90 min. is enough. The amt. of H_2 adsorbed by a given sample of the Ni catalyst (3.23 g., contg. 47.4 ml. H_{ads} and 177 ml. H_{dis}) proves to be a linear function of the remaining H_{ads} . Likewise, the catalytic activity in hydrogenation of allyl alc. is a linear function of H_{ads} , and both adsorption of H_2 and catalytic activity fall to zero when all the H_{ads} is removed by a sufficient amt. of I. The same behavior was observed with the use of $CH_2=CH(OEt)$ (II) as H-remover; allyl alc. had been previously shown to remove only H_{ads} , but not H_{dis} . The straight lines representing the change of adsorption and of catalytic activity, as a function of the remaining H_{ads} , have the same slope; in other words, the ratio H_{ads}/H_{dis} remains constant, and, in the given instance, is = 0.27. Both I and II are consumed in equimol. amts., equiv. to the amt. of H_{ads} removed. X-ray diagrams (Fe radiation, 30 kv., 10 ma.) show complete identity of the lattices of active Ni (contd. with H_{ads}), and of Ni inactivated by deprotonation with N. Thun

CA

2

Poisoning of hydrogenation-dehydrogenation catalysts in the light of the theory of their active structure. L. K. Frenkita and N. I. Ziminova (Inst. Org. Chem. Acad. Sci. U.S.S.R., Moscow). *Doklady Akad. Nauk S.S.S.R.* 76, 651-4 (1961).—Reasons are enumerated why poisoning of a great variety of hydrogenation-dehydrogenation catalysts (Pd, Pt, Ni, Co) with a great variety of catalyst "poisons" (O, S, Se, Te, P, As, Sb, Bi, Cl, Br, I, and their compounds) does not, as is commonly and generally assumed, consist in adsorptive "blocking" of active centers, but is due to removal of dissolved H that is an essential promoter of the catalysts. The poisons listed are all highly reactive towards H. The poisoning effect of this great variety of poisons cannot be due to a reaction with the metal, especially as I₂, which does not react with Pt-group metals at all, is a catalyst poison even at room temp. That, in all these instances, the "poisoning" consists simply in "depromotion" through removal of H, is further corroborated by the irreversible nature of that poisoning, and the necessity of renewed hydrogenation to restore the catalytic activity. Further proof is provided by the stoichiometric proportion between the consumption of the poison and the amt. of dissolved H, and the simple relation between the amt. of poison intro-

duced and the lowering of the activity. The very strong poisoning effect of H₂S can be explained by a chain reaction of the type $H_2S + H \rightarrow H_2 + HS$; $2HS \rightarrow H_2 + S_2$; $S + H \rightarrow HS$, etc. owing to which one mol. of H₂S can bring about the removal of a great no. of H atoms. Regeneration by a stream of H₂ has essentially the effect of bringing the catalyst anew with H. The observation of Shulkin, et al. (C. 1. 42, 4437a) that 1-ethyl-1-cyclopentene deactivates the Pt/C catalyst very rapidly, can be readily explained by a consumption of the dissolved H, in agreement with the easy zero-order hydrogenation of cyclopentene at room temp. under ordinary pressure; this hydrogenation takes place even in the absence of H₂, solely at the expense of the dissolved H, and results in complete poisoning of the catalyst. In the hydrogenation of 10 ml. of an 8.18% soln. of 1-methyl-1-cyclopentene (I) in C₆H₆, on 2.33 g. Ni, the initial rate of absorption of H₂ was 2 ml./min.; after 60 min., that rate fell to 0.5 ml./min., and, after another 90 min., to 0.2 ml./min. The I content of the soln. fell, at these stages, to 5 and to 1.1%, resp. Superficially adsorbed H is only loosely bound and, being consumed first, protects active centers against depromotion, as long as H₂ is supplied from without. Conditions favoring removal of the dissolved H, such as high temp., vacuum, or a stream of extraneous gas, depromote the catalyst. By practical experience in dehydrogenation and hydrogenation reactions, the strength of the bond between the dissolved-H promoter and the catalyst decreases in the order Pt > Ni > Pd. Acceptors of dissolved H have a depromoting action. The practice of carrying out dehydrogenation in a stream of H₂ is dictated by the advisability of keeping up a constant supply of the promoter.

ZIMINOVA, N. I.

ZIMINOVA, N. I. - "Investigation of the Nature of the Active Structure of a Skeleton Nickel Catalyst." Sub 5 Feb 52, Inst Organic Chemistry, Acad Sci USSR. (Dissertation for the Degree of Candidate in Chemical Sciences).

SO: Vechernaya Moskva January-December 1952

FREYDLIN, L.Kh.; ZHUKOVA, I.F.; ZIMIMOVA, N.I.; LAYNER, D.I.; KAGAN, H.M.

Deactivation of skeletal nickel catalyst by water vapor and enhancement of its stability by means of promoters. Kin. i kat. 2 no.1:112-117 Ja-F '61. (MIRA 14:3)

1. Institut organicheskoy khimii imeni N.D. Zelinskogo AN SSSR.
Institut giprotsvetmetobrabotka.
(Catalysts, Nickel)

TILGNER, Damazy (Gdansk-Wrzeszcz); ZIMINSKA, H., (Gdansk-Wrzeszcz); SZUBERT,
C., (Gdansk-Wrzeszcz)

Quality evaluation of 3 coffee extracts by means of sensory
analysis. Przem spoz 15 no.8:25-30 '61.

TILGNER, D.J. (Gdansk); ZIMINSKA, H. (Gdansk); SZUBERT, C. (Gdansk)

Detection of minute quality differences by the method of multiple comparisons. Przem spos 16 no.4:6-14 Ap '62

1. Katedra Technologii Zwierzecych Politechniki Gdanskiej Kierownik: prof. dr in. habil. D.J. Tilgner.

ZIMKIN, Andrey Vasilyevich; YUNCHENKO, L.I., red.; GORYACHEV, V.A.,
tekhn. red.

[At the sources of the Kolyma; notes of a geologist] U
istokov Kolymy; zapiski geologa. Magadan, Magadanskoe
knizhnoe izd-vo, 1963. 180 p. (MIRA 17:3)

ATLASOV, I.P.; VAKAR, V.A.; DIBNER, V.D.; YEGIAZAROV, B.Kh.; ZIMKIN, A.V.;
ROMANOVICH, B.S.

New tectonic map of the arctic regions. Dokl. AN SSSR 156
no.6:1341-1342 Je '64. (MIRA 17:8)

1. Nauchno-issledovatel'skiy institut geologii Arktiki.
Predstavleno akademikom D.V. Nalivkinym.

14-00000

L 19670-63

EWP(q)/EWT(m)/EWP(B)/BDS

AFFTC/ASD JD

ACCESSION NR: AR3006984

S/0058/63/000/008/E044/E044

SOURCE: RZh. Fizika, Abs. 8E314

59
58

AUTHORS: Zimkin, I. N.; Nadgornyy, E. M.; Smirnov, B. I.

TITLE: X-ray diffraction study of filament-like sodium chloride crystals

CITED SOURCE: Sb. shchelochnogaloidn. kristallov, Riga, 1962, 463-465

TOPIC TAGS: filament-like crystal , sodium chloride, X-ray dif-
fraction study

TRANSLATION: The method of diffraction microoentgenography (the Lang method) has been used to investigate the dislocation structure of filament-like crystals (FC) of NaCl. FC of NaCl grown by crystallization through a porous partition were investigated. It was

L 19670-63

ACCESSION NR: AR3006984

shown that in thin FC (10--20 μ) there are only dislocations, which are located along the growth axis (along the direction $\langle 100 \rangle$). Crystals of larger size have as a rule a more complicated dislocation structure. Heating of plastically bent FC leads to restoration of the dislocation structure existing prior to their bending. V. Regel'.
14

DATE ACQ: 06Sep63

SUB CODE: PH

ENCL: 00

ZIMKIN, I. N.; NADGORNYY, E. M.; SMIRNOV, B. I.

Studying whisker crystals of sodium chloride by the micro-
radiographic method. Fiz. tver. tela 5 no.1:170-176 Ja '63.
(MIRA 16:1)

1. Fiziko-tekhnicheskiy institut imeni A. F. Ioffe AN SSSR,
Leningrad.

(Microradiography) (Salt crystals)

L 42134-14 ENT(1)/ENP(2)/ENI(3)/ENP(4)/1/ENP(5)/ETI/ENP(6) TJP(c) NW/ID/HV
ACC NR: AP6027800 SOURCE CODE: UR/0126/66/022/001/0157/0158

AUTHOR: Konstantinov, B. P.; Zimkin, I. N.; Stepanov, M. I.; Shestopalov, L. M. 4/ B

ORG: Physicotechnical Institute im. A. F. Ioffe, AN SSSR (Fiziko-
tekhnicheskiy institut AN SSSR)

TITLE: Hardening of steel surface by wire explosion

SOURCE: Fizika metallov i metallovedeniye, v. 22, no. 1, 1966, 157-158

TOPIC TAGS: ~~steel~~ *metal* hardening, ~~steel~~ surface hardening, wire, ~~explosion~~, ~~explosive hardening~~ *steel*

ABSTRACT: Copper or steel wire, 0.38—0.4 mm in diameter and 40—50 mm long, placed 10 mm above the face of a cylindrical USA steel specimen was exploded by a current pulse produced by the discharge of a capacitor. As a result of this explosion, the surface microhardness increased from the original 170—200 kg/mm² to 950—1200 kg/mm². Although the average thickness of the hardened layer was 20—30 μ, it was uniform and varied from 0 to 60 μ. X-ray diffraction patterns showed that the ~~... 70% ...~~ the rest being mostly

L 42138-66

ACC NR: AP6027800

change to austenite, or the high rate of cooling preserved δ ferrite.
The block size in the hardened layer was about 450 Å. (ND)

SUB CODE: 11, 13/ SUBM DATE: 23 Sep65/ OTH REF: 001/ ATD PRESS:

5062

ZIMKIN, N., polkovnik meditsinskoy sluzhby, doktor med.nauk, prof.
KOROBKOV, A., podpolkovnik meditsiskoy sluzhby, doktor
meditsinskikh nauk, dotnent

Increasing the body's resistance. Voen.vest. no.9:92-95 S
'60. (MIRA 14:7)

(PHYSICAL EDUCATION AND TRAINING)

ZIMKIN, N. V.

PROCESSES AND PROPERTIES INDEX

Action of procaine on the effect produced on the pupil of a rabbit by stimulation of the trigeminal nerve. N. V. Zimkin and A. V. Lebedinskii. *J. Physiol. (U.S.S.R.)* 37: 103-104 (1958) (in Russian).—In expts. with rabbits, the effect of stimulation of the fifth pair of the trigeminal nerve in producing contraction of the pupil is confirmed. This effect persists when atropine, curare, nicotine, adrenaline or cocaine are administered either intravenously or directly under the conjunctiva. The administration of procaine (I) under the conjunctiva disturbed the transmission of excitation to the contractile elements of the pupil. In this way I completely prevents the development of the effect of antidromic excitation of the afferent nerve. This action of I develops along with disturbed transmission of other excitations along other innervation mechanisms of the contractile elements, thus showing that this action of I cannot be selective. After administration of I, the effect of acetylcholine on the pupil is greatly reduced. S. Gottlieb

Chair Physiology, Mil. Med Acad. im. Kirov.

ASB-51A METALLURGICAL LITERATURE CLASSIFICATION

CENTRAL LIBRARY

ZIMKIN, N.V.

Chair of Physiology, Military Med Acad of the Red Army ime. S.H.Kirov

On the functional structure of a reflex

So: Fiziologicheskly Zhurnal Vol 32, Nos 3⁵⁻⁶³ 1946

ZIMKIN, N. V.

114

Functional structure of reflexes. V. Facilitation of functional structure of reflexes under influence of poisons, which disturb the normal coordination (strychnine, alcohol), and soporific substances. N. V. Zimkin. *Vopr. Zhur. S.S.S.R. (J. Physiol.)* 23, 01-04 (1947) (Russian); *ibid.* 66, 300 (1949).—In studies on rabbits and frogs with strychnine (0.01-1.0 mg. for rabbit, 0.4-1.2 mg./kg. for frog), alc. (rabbits 2-6 ml./kg. injected rectally, frogs 4-20 ml./kg. injection into lymphatic sacs), and soporifics (barbital, chloral hydrate, and hexenal), it was shown that strychnine, in acting on the nerve centers, exhibits a considerable effect on the reflex mechanism in rabbits (shaking reflex) by increasing the threshold of some of the reflex components, and decreasing that of others; similar profound changes occurred in frog reflexes; the characteristic of strychnine action is the uneven effect on various nerve centers; the soporific substances listed above increase the stimulus thresholds and interrupted the transfer of stimuli from afferent to efferent neurons at low levels of excitation; at high excitation levels the reflexes occur with a fairly high order of coordination and movements. Alc. is between the strychnine and the soporifics in its effects; at medium dosage an uneven effect on various nerve centers occurs, similar to that of the strychnine effect but not as pronounced. In deeper alc. intoxication, however, the thresholds of excitability rise and the tonic reflexes disappear, showing similarity to the soporific action. G. M. Kozolapov

ASM 55.4 METALLURGICAL LITERATURE CLASSIFICATION

ZIMKIN, N. V.

See Also: ZIMKIN, A. M., and MIKHEL'SON, A. A.

"Problem of the variability of motor reflexes," Report 1, N. V. Zimkin, A. M. Zimkina and A. A. Mikhel'son, "Variability of the reflexogenic zones for the reflex of unbending the knee and the reflex of flexure of the foot from the rear," -- Report 2. N. V. Zimkin and A. A. Mikhel'son, "Peculiarities in the course of the knee-jerk reflex when the stimulus is a series of blows in close rhythm," Trudy Fiziol. in-ta Pavlova, Vol. III, 1949, p. 47-81 -- Bibliog: p. 60-61, 81

SO: U-3566, 15 March 53, (Letopis 'Zhurnal 'nykh Statey, No. 14, 1949).

ZIMKINA, A.H.; ZIMKIN, N.V.; KAPLAN, A.Ye.; MARENINA, A.I.; MIKHIL'SON, A.A.

Mobility of some reflex and sensory processes. Trudy fiziol. inst.
4:117-124 '49. (MIRA 9:5)

(REFLEXES) (SENSES AND SENSATION)

ZIMKIN, N.V. with Belen'kaya, Zelinkin and Kaplan

Chair of Physiol, Military Med Acad of the Red Army im S.M.Kirov

Lab of Physiol, Leningrad State Sci Inst im. Lesgaft

Inst Evolut Physiol and Pathol of the Higher Nervous Activity im. I.P.Pavlov, Acad Med Sci
USSR

Regulating the function of the spinal cord

So: Fiziologicheskii Zhurnal Vol 35, No 3, 1949

ZIMKIN, N. V.

Physiology

Problem of physiological characteristics of strength, speed, and endurance in
the light of I. P. Pavlov's teachings.
Teor. i prak. fizkul. 15 no. 4, 1952.

YELSHINA, M.A.; ZIMKIN, N.V.; MOISEVA, Z.Ye.

Formation of conditioned motor defense reflexes in mice following the application of an unconditioned stimulus before a conditioned stimulus. Zh. vys. nerv. deiat. 5 no.6:881-891 N-D '55. (MLRA 9:3)

1. Kafedra fiziologii Voenного instituta fizicheskoy kul'tury i sporta imeni V.I. Lenina.

(REFLEX, CONDITIONED,

defense & motor reflexes in mice, eff. of preliminary unconditioned stimulus)

ZIMKIN, N.V.

A. N. Krestovnikov, 1885-1955. Finiol. zhur. 41 no.3:459-460
My--Ja '55.

(MIRA 8:8)

(OBITUARIES,

Krestovnikov, Aleksei M)

ZIMKIN, N.V.; KAPLAN, A.Ye.; MEDVEDEV, V.I.

Change in the viscosity of saliva in dogs following disorders of the functional state of the central nervous system. *Fiziol. zhur.* 41 no.4:538-546 J1-Ag '55. (MLBA 8:10)

1. Kafedra fiziologii Voenno-meditsinskoy akademii im. S.M. Kirova i Institut evolyutsionnoy fiziologii i patologii vyushey nervnoy deyatel'nosti im. I.P.Pavlova, Leningrad.

(SALIVA,

viscosity, eff. of alcohol & strychnine in dogs)

(ALCOHOL, ETHYL, effects,
on saliva viscosity)

(STRYCHNINE, effect,
on saliva viscosity)

ZIMKIN, Nikolay Vasil'yevich; KHOTYANOVA, G.B., redaktor; SHALYGINA, G.A.,
tekhnicheskiiy redaktor

[Physiological aspects of strength, speed, and endurance; sketches]
Fiziologicheskaya kharakteristika sily, bystroty i vynoslivosti;
ocherki. Moskva, Gos. izd-vo "Fizkul'tura i sport," 1956. 205 p.
(PHYSICAL EDUCATION AND TRAINING) (MIRA 10:1)

Country : USSR
Category: Human and Animal Physiology. Nervous System.
Higher Nervous Activity. Behavior.

Abs Jour: RZhBiol., No 19, 1958, 89228

Author : Gakkel', L.B.; Zankina, A.M.; Zankin, N.V.;
Kaplan, A. Ye.; Kryshova, N.A.

Inst : -

Title : Afterimages in Patients with Brain Injuries.

Orig Pub: Zh. vyssh. nervn. deyat-sti, 1957, 7, No 4, 215-224

Abstract: In fifty patients with closed injuries of the brain, disappearance or marked shortening of afterimages (AI) in the optic, tactile and thermal analyzers was observed, which is considered a result of defensive inhibition. Asymmetry of AI was noted in patients with various degrees of damage of the right

Country : USSR
Category: Human and Animal Physiology. Nervous System.
Higher Nervous Activity. Behavior.
Abs Jour: RZhBiol., No 19, 1958, 89229
Author : Gol'danskaya, M.I.; Zinkina, A.M.; Zinkin, N.V.;
Ionisiani, G.L.; Kaplan, A. Ye.; Kryshova, N.A.
Inst : -
Title : The Thresholds of Sensitivity and Afterimages in the
Tactile, Thermal, Gustatory and Optic Analyzers in
Parkinsonians.
Orig Pub: Zh. vyssh. nervn. deyat-sti, 1957, 7, No 4, 225-234.
Abstract: The following was observed in 50 parkinsonian pa-
tients with average duration of their illness of
ten years: 1) elevation of the threshold in all
the investigated analyzers and their instability;

Country : USSR
Category: Human and Animal Physiology. Nervous System.
Higher Nervous Activity. Behavior.

T

Abs Jour: RZhBiol., No 19, 1958, 89229

2) paradoxical sensitivity in the gustatory analyzer; 3) the appearance (in the tactile and thermal) and prolongation (in the optic analyzer) of the latent period of the appearance of after-images (AI); 4) inconstancy of AI, and, frequently, prolongation of their duration; 5) more frequent appearance and greater quantity of secondary AI, than in normal; 6) apparent increase in the AI of the dimensions of the stimulated receptor zone; simultaneous manifestation of multiple and colored AI; changes in the form of optic AI.

ZIMKIN, N.V. (Leningrad)

Achievements in the physiology of physical education in the Soviet
Union. *Fiziol.shur.* 43 no.11:1037-1044 N '57. (MIRA 10:12)
(PHYSICAL EDUCATION AND TRAINING,
physiol. research in Russia (Bus))

ZIMKIN, N.V.

Arkadii Israelovich Bronshtein; 1896-1958; obituary. *Fiziol.shur.*
45 no.1:125-126 Ja '59. (MIRA 12:2)

(OBITUARIES,
Bronshtein, Arkadii I. (Rus))

ZIMKIN, H.V.

Physiology of work and sports. Fisiol.zhur. 45 no.11:1401-1402

N 159.

(MIRA 13:5)

(EXERTION physiol.)
(SPORTS)

ZIMKIN, N.V. (Leningrad)

Significance of the degree of exertion, duration and frequency of the exercises, and the intervals between exercises for the effectiveness of muscular training. Fiziol. Zhur. 46 no. 7:860-869 J1 '60. (MIRA 13:8)

(PHYSICAL EDUCATION AND TRAINING)

ZIMKIN, N.V. (Leningrad)

Congresses and symposia. Fiziol.zhur. 50 no.6:769-771. Je '64.
(MIRA 18:2)

SMIRNOV, K.M., prof., otv. red.; DAN'KO, Ya.I., prof., red.;
ZIMKIN, N.V., prof., red.

[Coördination of motor and vegetative functions in human
muscular activity] Koordinatsiia dvigatel'nykh i vegetativnykh
funktsii pri myshetschnoi deiatel'nosti cheloveka. Moskva,
Nauka, 1965. 137 p. (MIRA 18:12)

1. Akademiya nauk SSSR, Ob"yedinennyi nauchnyy sovet "Fiziol-
logiya cheloveka i zhivotnykh." 2. Gosudarstvennyy institut
fizicheskoy kultury im. P.F.Lesgafta, Leningrad (for Zimkin).
3. Pervyy Meditsinskiy ins'titut im. I.P.Pavlova, Leningrad
(for Dan'ko).
4. Gosudarstvennyy institut dlya usovershen-
stvovaniya vrachey im. S.M.Kirova, Leningrad (for Smirnov).

BIRYUKOV, Dmitriy Andreyevich, prof., otv. red.; GOLIKOV, N.V., red.;
ZIMKIN, N.V., red.; KARAMYAN, A.I., red.; KUPALOV, P.S., red.;
LAPINA, I.A., red.; VASIL'YEVA, Z.A., red.; KHARASH, G.A., tekhn.
red.

[Problems of the physiology and pathology of higher nervous activity]
Problemy fizologii i patologii vysshei nervnoi deiatel'nosti.
Pod obshchei red. D.A. Biriukova. Leningrad, Medgiz. No.2. 1963.
192 p. (MIRA 16:12)

1. Akademiya meditsinskikh nauk SSSR, Moscow. 2. Deystvitel'nyy
chlen AMN SSSR (for Biryukov).

(NERVOUS SYSTEM)

ZIMKIN, N.V. (Leningrad)

Symposium on the problems of fatigue and the restoration of
work capacity. Fiziol. zhur. 48 no.4:499-501 Ap '62. (MIRA 15:6)
(FATIGUE) (WORK)

ZIMKIN, N.V.

Stress during muscular exercise and the condition of non-specifically increased resistance of the body. Fiziol. zhur. 47 no.6:741-751 Ja '61. (MIRA 15:1)

1. From the P.F.Leshaft Institute of Physical Culture, Leningrad.
(EXERCISE) (STRESS)

BARYSHNIKOV, I.A.; BIRYUKOV, D.A.; ZIMKIN, N.V.

Twenty-second Congresses of the CPSU and some important problems
in physiology. Fiziol. zhur, 48 no.1:I-VIII Ja '62. (MIRA 15:2)
(COMMUNISM) (PHYSIOLOGY)

IORZH, K.P., kand.tekhn.nauk; MIMIREV, V.P., inzh; PREOBRAZHENSKIY, V.N.,
inzh.

Use of induction generators on ships. Sudostroenie no.7:32-35
J1 '60. (MIRA 13:7)
(Electricity on ships) (Induction (Electricity))

ZIMKIN, A.V.

Geology of northeastern Yakutia. Trudy IAFAN SSSR. Ser. geol.
no.3:3-28 '59. (MIRA 13:6)

(Yakutia--Geology)

ZIMKIN, A.V.

Verkhoyansk complex of sediments in the Yana Valley. Trudy
IAFAN SSSR.Ser.geol. no.3:29-55 '59. (MIRA 13:6)
(Yana Valley (Yakutia)—Geology, Stratigraphic)

ANIKEYEV, N.P.; BISKE, S.F.; VERESHCHAGIN, V.N.; ZINKIN, A.V.; LARIN, N.I.

Interdepartmental conference on the preparation of unified stratigraphic plans of the northeastern part of the U.S.S.R.
Sov. geol. no.62:182-188 '57. (MIRA 11:6)

1. Severo-Vostochnoye geologicheskoye upravleniye Ministerstva geologii i okhrany nedr SSSR i Vsesoyuznyy nauchno-issledovatel'skiy geologicheskiy institut.
(Siberia, Eastern--Geology, Stratigraphic)

21

4

AMPEROMETRIC DETERMINATION OF SULPHATES IN NICKEL PLATING BATHS.
 VF Torppova BA Zimkin and AA Popel. Zavodskaya Laboratoriya 1949,
 vol. 15, Apr., pp 404-407. In Russian. An account is given of an ex-
 perimental investigation carried out to determine the influence of
 chlorides on the results of amperometric determinations of sulphates.
 Two methods are proposed for the determination of the sulphate
 ion in the presence of chlorides in the second method acetate ions
 are added in sufficient concentration to prevent the precipitation
 of lead chloride while the precipitation of the sulphate takes
 place quantitatively. The results obtained are compared with those
 of gravimetric analysis.

SK

ASB.SLA METALLURGICAL LITERATURE CLASSIFICATION

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30
 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60
 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90
 91 92 93 94 95 96 97 98 99 100


21

7

Determination of Sulphates in Nickel Plating Solutions.
 V. F. Turpova, E. A. Zhukin, and A. A. Popel. (Electro-
 plating, 1948, vol. 3, Sec. pp. 122-123). This is an English
 summary of the paper which appeared in *Zavodskaya*
Laboratoriya, 1948, vol. 15, pp. 404-407. (See *Journ. I. and*
S.I., 1949, vol. 182, Nov., p. 246).

A 58-51A METALLURGICAL LITERATURE CLASSIFICATION

SEARCHED	SERIALIZED	INDEXED	FILED	MAY 1949	SERIALS	SECTION	SUBJECT	AUTHOR	TITLE	NUMBER	DATE
●	●	●	●	●	●	●	●	●	●	●	●



B

Amperometric Determination of Sulfates in Nickel-Flating Baths. (In Russian.) V. F. Toropova, E. A. Zimkin, and A. A. Popel. *Zavodskaya Laboratoriya* (Factory Laboratory), v. 15, Apr. 1949, p. 404-407.

On the basis of experiments, two methods for determining SO₄ ions in the presence of Cl ions are proposed: precipitation of Cl by AgNO₃, followed by titration of SO₄ in the presence of AgCl; and formation of a Pb acetate complex which impedes precipitation of PbCl₂.

458.554 METALLURGICAL LITERATURE CLASSIFICATION

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----

34

7

Amperometric determination of sulfates in nickel-plating baths. V. F. Torgova, B. A. Zimkin, and A. A. Popel. *Zashchita Lab.* 15, 604-7(1949). -- The sulfate in Ni-plating baths can be detd. amperometrically, with the dropping-Hg cathode and satd. calomel electrode anode, with agar-KNO₃ bridge, and 0.8 v. potential im-

pressed on the system from an external source and with a galvanometer to record the current. The soln. is titrated with 0.2 N Pb(NO₂)₂ which gives curves with sharp breaks at the equiv. points.
G. M. Kosolapoff

ASB-51A METALLURGICAL LITERATURE CLASSIFICATION

EXPERIMENTAL

SEARCHED	INDEXED	SERIALIZED	FILED
NOV 19 1949	NOV 19 1949	NOV 19 1949	NOV 19 1949

